A Partially Defined Game with Payments

Satoshi Masuya

Faculty of Business Administration, Daito Bunka University. 1-9-1, Takashimadaira Itabashi-ku TOKYO 175-8571 (Japan) masuya@ic.daito.ac.jp

Abstract. We investigate a new problem that can be solved by using the theory of a partially defined game. We consider the situation described below: first, we assume that the worth of the grand and singleton coalitions is only known. It take some amount of costs to obtain worth of larger coalitions. If it is performed, then players make a payment from the worth of the grand coalition. That is, the worth of the grand coalition is reduced by examinations of coalitional worth. The problem of a partially defined game with payments is finding the solution of partially defined games at each point and the best exiting rule of examinations of coalitional worth.

Keywords: Cooperative game, Partially defined cooperative game, Payment, Shapley value, Exiting rule

AMS Classification Code: 91A12

Competing interests: The authors declare no competing interests.

1 Introduction

Cooperative game theory provides a useful tool to analyze various cost and/or surplus allocation problems, the distribution of voting power in a parliament, and so on. This theory is employed to analyze problems that involve n entities called players which are usually expressed by characteristic functions that map each subset of players to a real number. The solutions are given by a set of n-dimensional real numbers or a value function that assigns a real number to each player. Such a real number can represent the cost borne by the player, the power of influence, allocation of shared profits, and so on.

We deal with a cooperative game that is called a partially defined cooperative game in this paper. A partially defined cooperative game (a PDG, in short) is a cooperative game in which the worth of some coalitions is unknown. A cooperative game is called full if all worth of coalitions is known.

Willson [12] first considered partially defined cooperative games. He introduced a generalized Shapley value [11] derived solely from the known worth of coalitions in a game.

After that, many results are obtained in this area. Recently, Yu [13] studied a cooperative game with a coalition structure under limited feasible coalitions.

Here, a PDG with a coalition structure was considered, and he developed and axiomatized its Owen value [10]. Aguilera et al. [1] and Calvo and Gutiérrez [5] proposed independently the same extension of the Shapley value for PDGs from different points of view. This extended value was characterized for games with restricted cooperation in Albizuri et al. [2] with three axioms which are more elementary than that proposed in [5]. That is, this type of extended Shapley value has been often studied. On the other hand, Černý and Grabisch [6] studies so called player-centered PDGs and they derived the collection of monotonic full games which can be obtained from a PDG within the class of such games.

Partially defined games and restricted games are same concepts mathematically although the reasons in which they have unknown coalitions are different. PDGs have unknown worth of coalitions since examinations of all worth of coalitions are very expensive while restricted games have unknown worth of coalitions since those coalitions cannot form because of the difference of ideologies among players, and so on.

Myerson [8] first considered restricted games using the set of feasible coalitions that is called communication situations. Subsequently, he proposed and axiomatized the Shapley value for restricted games which is called the Myerson value. In the line of this research, many studies generalize the set of feasible coalitions of a restricted game. As representative studies, conference structures by Myerson [9] and union stable systems by Algaba et al. [3] can be mentioned.

In this paper, we investigate a new problem which can be solved by using a concept of a PDG but cannot by using that of a restricted game.

We consider the situation which is described below: first, we assume that the worth of the grand and singleton coalitions is only known. It take some amount of costs to obtain worth of larger coalitions. If it is performed, then players make a payment from the worth of the grand coalition. That is, the worth of the grand coalition is reduced by examinations of coalitional worth.

With the view point of fairness of an allocation of payoffs, we should examine coalitional worth as many as possible. However, we should stop examining coalitional worth at some point since total payoff is reduced by continuing the examinations. We name the new decision making problem a partially defined game with payments. The problem of a PDG with payments is finding the solution of PDGs at each point and the best exiting rule of examinations of coalitional worth.

We extend the Shapley value for PDGs that is proposed in Aguilera et al. [1], Calvo and Gutiérrez [5], and Albizuri et al. [2] to PDGs with payments and axiomatize the proposed value. Furthermore, we propose an rule to finish examinations of coalitional worth for the problems and axiomatize the proposed rule.

This paper is organized as follows. In Section 2, we present the definitions of partially defined games and their Shapley value. In Section 3, we define a partially defined game with payments and the assumptions a cost function has in this paper. In Section 4, we propose and axiomatize the Shapley value for PDGs with payments. In Section 5, we propose a rule of stopping examinations

of coalitional worth and axiomatize the proposed rule. In Section 6, concluding remarks are given.

2 The Shapley Value and Partially Defined Games

In this section, we provide the definition and related concepts of partially defined games and the Shapley value for PDGs that is proposed by Albizuri et al. [2], which is used to define the Shapley value for PDGs with payments.

Let $N = \{1, 2, \dots, n\}$ be the set of players. A non-empty set of players $S \subseteq N$ is called a coalition. Then the pair (N, v) where $v : 2^N \to \mathbb{R}$ is called a TU-game. For every coalition $S \subseteq N$, v(S) is called the worth of the coalition S.

A game (N, v) is superadditive if and only if:

$$v(S \cup T) \ge v(S) + v(T), \ \forall S, \ T \subseteq N \text{ such that } S \cap T = \emptyset.$$
 (1)

Superadditivity is a natural property that gives each player an incentive to form a larger coalition. The set of superadditive games is denoted Γ^N .

The Shapley value [11] is a well-known one-point solution concept for TU-games and its explicit form is:

$$\phi_i(v) = \sum_{\substack{S \subseteq N \\ S \ni i}} \frac{(|S| - 1)!(n - |S|)!}{n!} (v(S) - v(S \setminus i)), \ \forall i \in N.$$
 (2)

A PDG, on the set of known coalitions \mathcal{K} is a set-function v which maps every set $S \in \mathcal{K}$ a real number v(S), such that $v(\emptyset) = 0$. A triple (N, \mathcal{K}, v) identifies a PDG. Usually, it is assumed that the worth of grand coalition is known.

Moreover, we assume that v is superadditive in the following sense:

$$v(S) \ge \sum_{i=1}^{s} v(T_i), \ \forall S, T_i \in \mathcal{K}, \ i = 1, 2, \dots, s \text{ such that } \bigcup_{i=1, 2, \dots, s} T_i = S$$
 and $T_i, \ i = 1, 2, \dots, s$ are disjoint. (3)

We provide the definition of an extension of the Shapley value for PDGs by Albizuri et al. [2]. According to Harsanyi's procedure [7], all the members of a coalition S receive a dividend from S. We will distinguish two cases depending on whether the coalitional worth is known or not: (i) if $S \in \mathcal{K}$ the total amount of the dividends allocated by all the subcoalitions of S is v(S), (ii) otherwise, i.e. if $S \notin \mathcal{K}$, the dividend of S is zero. Formally the procedure can be described as follows. If v is a PDG, define recursively a function $d_v: 2^N \to \mathbb{R}$ by:

$$d_v(\emptyset) = 0$$
; and (4)

$$d_v(S) = \begin{cases} v(S) - \sum_{T \subseteq S} d_v(T), & \text{if } S \in \mathcal{K}, \\ 0, & \text{if } S \notin \mathcal{K}. \end{cases}$$
 (5)

The real number $\frac{d_v(S)}{|S|}$ is usually called the Harsanyi dividend of coalition S in v. The Shapley value for PDGs by [1][5][2] $\hat{\phi}$ is defined as follows:

$$\hat{\phi}_i(v) = \sum_{S \subseteq N, S \ni i} \frac{d(v, S)}{|S|} \quad \forall i \in N.$$
 (6)

In this study, we assume a property of K that is defined as follows:

$$\mathcal{K} = \{ S \subseteq N \mid |S| \le k \} \cup \{ N \}, \text{ for some } 1 \le k \le n - 1.$$
 (7)

We can establish $k=\max\{|S|\mid S\in\mathcal{K},S\neq N\}$. Therefore, a PDG (N,\mathcal{K},v) satisfying equation (7) is called an (N,k)-PDG. Thus, $1\leq k\leq n-1$ holds. When k=n-1, (N,\mathcal{K},v) is a full game and when k=1, the game is a PDG where only the worth of the grand and singleton coalitions is known. An (N,k)-PDG (N,\mathcal{K},v) could be written as (N,k,v). The set of (N,k)-superadditive PDGs is denoted $\Gamma^{N,k}$. The relation between (N,k+l,v) $(l\geq 1)$ and (N,k,v) is as follows:

$$(N, k+l, v)(S) = (N, k, v)(S)$$
 for all $S \subseteq N$ such that $|S| \le k$ or $|S| = n$. (8)

3 Superadditive Partially Defined Games with Payments

In this section, we define a superadditive PDG with payments and related concepts.

We consider the situation which is described below: On the first stage, we assume that the worth of the grand and singleton coalitions is only known. It takes some amount of costs to obtain worth of larger coalitions. The cost function of $v \in \Gamma^N$ is defined by $c_v(s)$ for every $2 \le s \le n-1$ since all worth of coalitions whose cardinality is the same is obtained at one examination. If it is performed, then players pay the fee from the worth of the grand coalition. That is, if the examination is performed for s=2, then v(N) is reduced to $v(N)-c_v(2)$. If $c_v(s)=0$ for all $2 \le s \le n-1$, then all coalitional worth can be obtained with no costs.

In this situation, if players are not satisfied with the allocated payoffs of the first game (that is, the game in which the worth of the grand and singleton coalitions is only known) and agree to the payment to examine the coalitional worth whose cardinality is two, then the worth of coalitions is obtained. Then the same dialog is performed. If they are not satisfied with the allocated payoffs even if they consider the cost, then the worth of coalitions whose cardinality is three is obtained. If they obtain the worth of coalitions whose cardinality is n-1, then the examination is finished and they are allocated the payoffs of the last game. A triple (N,k,v) and c_v identifies a PDG with payments. We assume that a game satisfies superadditivity and $v(S) \geq 0$ for all $S \subseteq N$ such that $|S| \leq k$ or |S| = n. Since N is fixed in this study, we simply write v^k and c_v to represent a PDG with payments if there is no confusion. The problem of

a PDG with payments is finding the solution of PDGs and the best exiting rule of examinations of coalitional worth.

Here, we give several assumptions with respect to a cost function in this study.

Assumption 1 A cost function is known by all players.

Assumption 2 Let $v \in \Gamma^N$ and $k \in \{2, ..., n-1\}$. If the examination is proceeded to stage k, then the following holds:

$$v(N) - \sum_{l=2}^{k} c_v(l) \ge v(N \setminus S) \text{ for all } S \subset N \text{ such that } |N \setminus S| = k.$$
 (9)

Assumption 2 states that examinations of coalitional worth are performed if a game satisfies monotonicity only.

Assumption 3 Let $v \in \Gamma^N$ and $k \in \{1, ..., n-2\}$. If v(S) has been examined for all $S \subset N$ such that $|S| \leq k$, and $c_v(k+1) = 0$ holds, then v(T) is also examined for all $T \subset N$ such that |T| = k+1.

Assumption 3 states that the examination of coalitional worth is surely performed if the cost of it is zero. That is, if $c_v(s) = 0$ for all $s \in \{2, ..., n-1\}$, then all coalitional worth is examined.

Assumption 4 Let $v \in \Gamma^N$ and $k \in \{2, ..., n-1\}$. If $d_v(S) = 0$ for all $S \subset N$ such that |S| = k, then $c_v(k) = 0$ holds.

If the examination of coalitional worth yields $d_v(S) = 0$ for all $S \subset N$ with |S| = k, then employing the solution $\hat{\phi}$ provides no information to players. Thus, Assumption 4 states that $c_v(s) = 0$ for such a result.

In the following, we show the relation between $v \in \Gamma^N$ and $v^k \in \Gamma^{N,k}$ as follows:

$$v^{k}(S) = v(S) \text{ if } |S| \le k \text{ or } |S| = n.$$
 (10)

for every $k \in \{1, ..., n-1\}$.

A solution for PDGs with payments is represented by $\sigma(v) \in \mathbb{R}^{n \times (n-1)}$. That is, $\sigma(v)$ is represented by a set of n-dimensional real numbers for every stage from one to n-1. The solution of Stage k is represented by $\sigma(v^k)$ for all $k \in \{1, \ldots, n-1\}$.

Assume that we obtained a solution for (N,k)-superadditive PDGs $\sigma: \bigcup_{k=1}^{n-1} \Gamma^{N,k} \to \mathbb{R}^{n\times (n-1)}$. Let $\sigma(\bigcup_{k=1}^{n-1} \Gamma^{N,k})$ is the set of solutions that is obtained from its domain $\bigcup_{k=1}^{n-1} \Gamma^{N,k}$. Then we define an indicator function $\pi: \sigma(\bigcup_{k=1}^{n-1} \Gamma^{N,k}) \to \{0,1\}^{n-1}$ which indicates the exiting rule of the examination of the coalitional worth. For each $k \in \{1,\ldots,n-1\}$, if $\pi(\sigma(v^k))=1$, then the next examination is performed toward Stage k+1. If $\pi(\sigma(v^k))=0$, then the examination is terminated at Stage k. If k=n-1, then $\pi(\sigma(v^k))=0$ for any solutions. In this study, we propose a solution for (N,k)-superadditive PDGs with payments and an indicator function for the solution and axiomatize them.

4 The Shapley Value for Superadditive Partially Defined Games with Payments and Its Axiomatization

In this study, we use the Shapley value for partially defined games by [2] to define the Shapley value for (N, k)-superadditive PDGs with payments. Let $v \in \Gamma^N$. The Shapley value $\tilde{\phi} : \bigcup_{k=1}^{n-1} \Gamma^{N,k} \to \mathbb{R}^{n \times (n-1)}$ is defined as follows:

$$\tilde{\phi}_{i}(v^{k}) = \sum_{\substack{S \subset N \\ S \ni i, 1 \le |S| \le k}} \frac{d_{v^{k}}(S)}{|S|} + \frac{d_{v^{k}}(N) - \sum_{s=2}^{k} c_{v}(s)}{n} \quad \forall i \in N.$$
 (11)

for every $k \in \{1, \dots n-1\}$.

The proposed value is essentially the same as the value proposed by [2]. The difference between them is considering the cost to examine the coalitional worth.

We axiomatize the proposed value $\tilde{\phi}$. Let $\sigma: \bigcup_{k=1}^{n-1} \Gamma^{N,k} \to \mathbb{R}^{n \times (n-1)}$. In the axiomatization of the proposed value, we use the concept of the degree of increase with respect to a marginal contribution.

Let $v \in \Gamma^N$. Then we define $M_i^S(v^k) = v^k(S) - v^k(S \setminus i)$ for every $S \subseteq N$ such that $1 \leq |S| \leq k$. In addition, let $v, w \in \Gamma^N$ and $d \in \mathbb{R}$. Then we define $(v^k + w^k)(S) = v^k(S) + w^k(S)$ and $(dv^k)(S) = d \cdot v^k(S)$ for every $S \subseteq N$ such that $1 \leq |S| \leq k$ or |S| = n.

Axiom 1 (Efficiency) Let $v \in \Gamma^N$. Then the following holds:

$$\sum_{i \in N} \sigma_i(v^k) = v(N) - \sum_{s=2}^k c_v(s), \text{ for all } k \in \{2, \dots, n\}.$$
 (12)

Axiom 2 (Linearity) Let $v_1, v_2 \in \Gamma^N$ and $d_1, d_2 \in \mathbb{R}$. Then the following holds:

$$\sigma(d_1v_1^k + d_2v_2^k) = d_1\sigma(v_1^k) + d_2\sigma(v_2^k), \text{ for all } k \in \{1, \dots, n\}.$$
 (13)

Axiom 2 is the adaptation of the axiom of Linearity for the solution of TUgames to PDGs with payments.

Axiom 3 (Dummy player) Let $v \in \Gamma^N$, $k \in \{2, ..., n\}$, and $i \in N$. Assume that $v(N) - \sum_{S \subset N, |S| = k} v(S) = v(N) - \sum_{S \subset N, |S| = k-1} v(S)$. If $M_i^S(v^k) - M_i^{S \setminus j}(v^{k-1}) = 0 \ \forall S \subseteq N \ such \ that \ S \ni i \ and \ 1 \le |S| \le k, \ \forall j \in S \setminus i, \ then \ the following \ holds:$

$$\sigma_i(v^k) = \sigma_i(v^{k-1}) - \frac{c_v(k)}{n}, \text{ for all } k \in \{2, \dots, n\}.$$
 (14)

Axiom 3 states that if marginal contributions of a player do not increase with respect to set-theoretic inclusion, then the allocated payoff to him does not increase. That is, he is a dummy player.

Axiom 4 (Players with Same contributions) Let $v \in \Gamma^N$, $k \in \{2, ..., n\}$, and $i, l \in N$. If $M_i^S(v^k) - M_i^{S \setminus j}(v^{k-1}) = M_l^S(v^k) - M_l^{S \setminus j}(v^{k-1}) \quad \forall S \subseteq N \text{ such that } S \supseteq \{i, l\} \text{ and } |S| = k, \quad \forall j \in S \setminus \{i, l\}, \text{ then the following holds:}$

$$\sigma_i(v^k) - \sigma_i(v^{k-1}) = \sigma_l(v^k) - \sigma_l(v^{k-1}), \text{ for all } k \in \{2, \dots, n\}.$$
 (15)

Axiom 4 states that if the degrees of increases with respect to marginal contributions of two players coincide, then the difference of allocated payoffs to them between two stages are equal.

Axiom 5 (Fairness of first stage) Let $v \in \Gamma^N$. Then the following holds:

$$\sigma_i(v^1) - \sigma_j(v^1) = v(i) - v(j), \text{ for all } i, j \in N.$$

$$\tag{16}$$

Lemma 1. Let $v \in \Gamma^N$, $k \in \{2, ..., n-1\}$, and $i \in N$. If $M_i^S(v^k) = M_i^{S \setminus j}(v^{k-1})$ holds for any $S \subset N$ such that $S \ni i$ and |S| = k for all $j \in S \setminus i$, the following holds:

$$\sum_{\substack{S \subset N \\ S \ni i, |S| = k}} d_{v^k}(S) = 0. \tag{17}$$

Proof. We have:

$$M_{i}^{S}(v^{k}) - M_{i}^{S\setminus j}(v^{k-1}) = v^{k}(S) - v^{k}(S\setminus i) - (v^{k-1}(S\setminus j) - v^{k-1}((S\setminus j)\setminus i)) = \sum_{T\subseteq S, |T|\leq k} d_{v^{k}}(T) - \sum_{T\subseteq S\setminus i, |T|\leq k} d_{v^{k}}(T) - \sum_{T\subseteq S\setminus j, |T|\leq k-1} d_{v^{k-1}}(T) + \sum_{T\subseteq S\setminus j\setminus i, |T|\leq k-1} d_{v^{k-1}}(T) = \sum_{T\subseteq S\atop T\ni i, |T|\leq k} d_{v^{k}}(T) - \sum_{T\subseteq S\setminus j\atop T\ni i, |T|\leq k-1} d_{v^{k-1}}(T) = 0.$$

$$(18)$$

Using the induction with respect to k, we show that equation (17) holds. When k = 2, we have:

$$d_{v^{k-1}}(S \setminus j) = d_{v^{k-1}}(i) = v(i). \tag{19}$$

On the other hand,

$$\sum_{\substack{T \subseteq S \\ T \ni i, |T| \le k}} d_{v^k}(T) = v(i) + \sum_{\substack{T \subseteq S \\ T \ni i, |T| = k}} d_{v^k}(T). \tag{20}$$

Using equation (18) and calculating equation (19) - (20), $\sum_{T \ni i, |T| = k} d_{v^k}(T) = 0$ holds.

Assume that $\sum_{\substack{S \subset N \\ S \ni i, |S| = k}} d_{v^k}(S) = 0$ holds when k = l.

When k = l + 1, from equation (18), the following holds:

$$\sum_{\substack{T \subseteq S \\ T \ni i, |T| \le l+1}} d_{v^k}(T) = \sum_{\substack{T \subseteq S \setminus j \\ T \ni i, |T| \le l}} d_{v^{k-1}}(T)$$

$$\Leftrightarrow \sum_{\substack{T \subseteq S \\ T \ni i, l \le |T| \le l+1}} d_{v^k}(T) = \sum_{\substack{T \subseteq S \setminus j \\ T \ni i, |T| = l}} d_{v^{k-1}}(T).$$

Thus
$$\sum_{\substack{T \subseteq S \\ T \ni i, |T| = l}} d_{v^k}(T) = \sum_{\substack{T \subseteq I, |T| = l+1 \\ T \ni i, |T| = l+1}} d_{v^k}(T) = 0.$$

From induction hypothesis, $\sum_{\substack{T\subseteq S\backslash j\\T\ni i,|T|=l}}d_{v^{k-1}}(T)=0 \text{ holds for all } j\in S.$ Thus $\sum_{\substack{T\subseteq S\\T\ni i,l\leq |T|\leq l+1\\S\ni i,|S|=k}}d_{v^k}(T)=\sum_{\substack{T\subseteq S\\T\ni i,|T|=l+1\\S\ni i,|S|=k}}d_{v^k}(T)=0.$ That is, $\sum_{\substack{S\subseteq N\\S\ni i,|S|=k}}d_{v^k}(S)=0 \text{ holds for all } k\in\{2,\ldots,n-1\}.$ This completes a proof the proof.

Theorem 1. $\tilde{\phi}$ is the unique function on $\bigcup_{k=1}^{n-1} \Gamma^{N,k}$ that satisfies Axioms 1 through 5.

Proof. First, we show that $\tilde{\phi}$ satisfies Axioms 1 through 5. From the definition of $\tilde{\phi}$, it is clear that $\tilde{\phi}$ satisfies Axiom 1 and 2. Moreover, it is straightforward to show that ϕ satisfies Axiom 5.

We show that $\tilde{\phi}$ satisfies Axiom 3. Let $v \in \Gamma^N$, $k \in \{2, \dots, n-1\}$, and $i \in N$. Let $M_i^S(v^k) = M_i^{S\setminus j}(v^{k-1})$ holds for any $S \subset N$ such that $S \ni i$ and |S| = k for all $j \in S \setminus i$. We have:

$$\begin{split} \tilde{\phi}_{i}(v^{k}) - \tilde{\phi}_{i}(v^{k-1}) \\ &= \sum_{\substack{S \subset N \\ S \ni i, |S| \leq k}} \frac{d_{v^{k}}(S)}{|S|} + \frac{d_{v^{k}}(N) - c(k)}{n} - \sum_{\substack{S \subset N \\ S \ni i, |S| \leq k-1}} \frac{d_{v^{k-1}}(S)}{|S|} + \frac{d_{v^{k-1}}(N)}{n} \\ &= \sum_{\substack{S \subset N \\ S \ni i, |S| = k}} \frac{d_{v^{k}}(S)}{|S|} + \frac{-\sum_{S \subset N, |S| = k}}{n} \frac{d_{v^{k}}(S) - c(k)}{n} \\ &= \sum_{\substack{S \subset N \\ S \ni i, |S| = k}} \frac{d_{v^{k}}(S)}{|S|} - \frac{c(k)}{n} \\ &= -\frac{c(k)}{n}. \end{split}$$

The second equality follows from the definition of the Harsanyi dividend d_v . The third equality follows since $\sum_{S \subset N, |S| = k} d_{v^k}(S) = 0$ holds from v(N) $\sum_{S\subset N,|S|=k}v(S)=v(N)-\sum_{S\subset N,|S|=k-1}v(S)$. The last equality follows from That is, $\tilde{\phi}$ satisfies Axiom 3.

We show that $\tilde{\phi}$ satisfies Axiom 4. Let $v \in \Gamma^N$, $k \in \{2, ..., n-1\}$, and $i, l \in N$. From the proof of the satisfaction of Axiom 3 above, the following holds:

$$\tilde{\phi}_{i}(v^{k}) - \tilde{\phi}_{i}(v^{k-1}) = \sum_{\substack{S \subset N \\ S \ni i, |S| = k}} \frac{d_{v^{k}}(S)}{k} - \frac{c(k)}{n}$$
(21)

$$\tilde{\phi}_{l}(v^{k}) - \tilde{\phi}_{l}(v^{k-1}) = \sum_{\substack{S \subset N \\ S \ni l, |S| = k}} \frac{d_{v^{k}}(S)}{k} - \frac{c(k)}{n}$$
(22)

From $M_i^S(v^k)-M_i^{S\backslash j}(v^{k-1})=M_l^S(v^k)-M_l^{S\backslash j}(v^{k-1})$ holds from the assumption of Axiom 4, $\sum_{\substack{S\subset N\\S\ni i,|S|=k}}\frac{d_{v^k}(S)}{k}=\sum_{\substack{S\subset N\\S\ni l,|S|=k}}\frac{d_{v^k}(S)}{k} \text{ holds. Thus, } \tilde{\phi} \text{ satisfies Axiom 4.}$

Next, we show the uniqueness. Let $\sigma: \Gamma^N \to \mathbb{R}^{n \times (n-1)}$. Let u_S be a unanimity game for every $S \subseteq N$, and let u_S^k a PDG of u_S of each stage $k \in \{1, \dots, n-1\}$.

We show the uniqueness of $\sigma(u_S^k)$ for all $k \in \{1, \dots, n-1\}$ for every $S \subset N$ using the induction with respect to k.

When k=1, $\sigma_i(u_S^1)=\sigma_j(u_S^1)$ holds $\forall i,j\in N$ from Axiom 5 if $|S|\geq 2$. That is, $\sigma_i(u_S^1)=\frac{1}{n}$ for all $i\in N$ from Axiom 1. Similarly, if |S|=1, the following holds:

$$\sigma_i(u_S^1) = \begin{cases} 1, & \text{if } i \in S, \\ 0, & \text{otherwise.} \end{cases}$$
 (23)

That is, $\sigma(u_S^k)$ is obtained uniquely when k=1.

Assume that $\sigma(u_S^k)$ is obtained uniquely when k = l. Let k = l + 1. We consider the case of |S| = l + 1. Let $i_1, i_2, j \in T$ such that |T| = |S|, and let $i_1, i_2 \in S$ and $j \notin S$.

Then the following holds:

$$M_{i_1}^T(u_S^{l+1}) - M_{i_1}^{T \setminus j}(u_S^l) = M_{i_2}^T(u_S^{l+1}) - M_{i_2}^{T \setminus j}(u_S^l).$$
 (24)

Also, for $i_1, i_2, j \in N$ such that $i_1, i_2, j \in S$, equation (24) holds. Therefore, from Axiom 4, the following holds:

$$\sigma_{i_1}(u_S^{l+1}) - \sigma_{i_2}(u_S^{l+1}) = \sigma_{i_1}(u_S^l) - \sigma_{i_2}(u_S^l). \tag{25}$$

for all $i_1, i_2 \in S$.

Let $i_3 \in N \setminus S$, and $i_3 \in R$ such that $R \neq S$ and |R| = l+1. Then $M_{i_3}^R(u_S^{l+1}) - M_{i_3}^{R\setminus j}(u_S^l) = 0$ holds. Moreover, $u_S(N) - \sum_{T\subset N, |T| = l+1} u_S(T) = u_S(N) - \sum_{T\subset N, |T| = l} u_S(T)$ also holds. Therefore, $\sigma_{i_3}(u_S^{l+1}) = \sigma_{i_3}(u_S^l) - \frac{c_{u_S}(l+1)}{n}$ holds from Axiom 3.

Thus, from the induction hypothesis, $\sigma_i(u_S^{l+1})$ is obtained uniquely for all $i \in N \setminus S$.

Moreover, from equation (25), we obtain (|S|-1) linearly independent equations. Additionally, from Axiom 1, we obtain (|S|) linearly independent equations. That is, we obtain $\sigma_i(u_S^{l+1})$ is obtained uniquely for all $i \in S$. Namely, we obtain $\sigma(u_S^k)$ uniquely when k = l + 1.

Finally, we consider the case |S|=n. In this case, $\sigma_i(u_S^l)=\frac{1-\sum_{m=2}^l c_{u_S}(m)}{n} \ \forall i\in N$ holds for any $l\in\{2,\ldots,n-1\}$ from Axiom 1, 4, and 5 using a similar manner used above.

Finally, using Axiom 2, we have:

$$\sigma(v^k) = \sigma(\sum_{S \subseteq N} d_S u_S^k) = \sum_{S \subseteq N} d_S \sigma(u_S^k)$$
 (26)

for all $k \in \{1, ..., n-1\}$

This completes the proof.

Example 1. Let $N = \{1, 2, 3, 4\}$ and $v \in \Gamma^N$, and let $c_v(2) = 1$, $c_v(3) = 4$. v is defined as follows:

$$\begin{split} v(\{1\}) &= 8, \ v(\{2\}) = 4, \ v(\{3\}) = 2, \ v(\{4\}) = 0, \\ v(\{1,2\}) &= 15, \ v(\{1,3\}) = 12, \ v(\{1,4\}) = 8, \ v(\{2,3\}) = 10, \\ v(\{2,4\}) &= 4, \ v(\{3,4\}) = 2, \ v(\{1,2,3\}) = 20, \ v(\{1,2,4\}) = 15, \\ v(\{1,3,4\}) &= 12, \ v(\{2,3,4\}) = 10, \ v(\{1,2,3,4\}) = 25. \end{split}$$

The Harsanyi dividend d_v can be obtained as follows:

$$\begin{split} &d_v(\{1\})=8,\ d_v(\{2\})=4,\ d_v(\{3\})=2,\ d_v(\{4\})=0,\\ &d_v(\{1,2\})=3,\ d_v(\{1,3\})=2,\ d_v(\{1,4\})=0,\ d_v(\{2,3\})=4,\\ &d_v(\{2,4\})=0,\ d_v(\{3,4\})=0,\ d_v(\{1,2,3\})=-3,\ d_v(\{1,2,4\})=0,\\ &d_v(\{1,3,4\})=0,\ d_v(\{2,3,4\})=0,\ d_v(\{1,2,3,4\})=5. \end{split}$$

Then, $\tilde{\phi}(v^k)$ for all $k \in \{1, 2, 3\}$ is obtained from equation (11) as follows:

$$\begin{split} \tilde{\phi}_1(v^1) &= \frac{43}{4}, \ \tilde{\phi}_2(v^1) = \frac{27}{4}, \ \tilde{\phi}_3(v^1) = \frac{19}{4}, \ \tilde{\phi}_4(v^1) = \frac{11}{4}. \\ \tilde{\phi}_1(v^2) &= \frac{43}{4}, \ \tilde{\phi}_2(v^2) = \frac{31}{4}, \ \tilde{\phi}_3(v^2) = \frac{21}{4}, \ \tilde{\phi}_4(v^2) = \frac{1}{4}. \\ \tilde{\phi}_1(v^3) &= \frac{38}{4}, \ \tilde{\phi}_2(v^3) = \frac{26}{4}, \ \tilde{\phi}_3(v^3) = \frac{16}{4}, \ \tilde{\phi}_4(v^3) = 0. \end{split}$$

5 An Indicator Function for the Exit Point of Examinations of Coalitional Worth

Assume that we obtained a solution for (N,k)-superadditive PDGs $\sigma: \bigcup_{k=1}^{n-1} \Gamma^{N,k} \to \mathbb{R}^{n\times(n-1)}$. Then we define an indicator function $\pi: \sigma(\bigcup_{k=1}^{n-1} \Gamma^{N,k}) \to \{0,1\}^{n-1}$ which indicates the exit point of examinations of the coalitional worth. For some $k \in \{1,\ldots,n-1\}$, if $\pi(\sigma(v^k))=1$, then the next examination is performed toward Stage k+1. If $\pi(\sigma(v^k))=0$, then the examination is terminated at Stage k.

We consider an optimal indicator function of a PDG using the solution $\tilde{\phi}$ to determine allocated payoffs for players. If we terminate examinations of coalitional worth early, then the cost for examinations is low. However, we have to determine the allocated payoffs under uncertain environment.

We may propose several indicator functions for PDGs with payments. In this paper, we propose an indicator function and axiomatize it.

Definition 1 (Null player). Let $v \in \Gamma^N$ and c_v a cost function of v. Let $k \in \{1, \ldots, n-1\}$. Then player $i \in N$ is said to be a null player of v^k under c_v if $v^k(S) - v^k(S \setminus i) = 0$ for all $S \subset N$ such that $S \ni i$ and $|S| \le k$, and $d_{v^k}(N) - \sum_{l=2}^k c_v(l) = 0$.

Let $\tilde{\phi}(\bigcup_{k=1}^{n-1} \Gamma^{N,k}) = \{A \in \mathbb{R}^{n \times (n-1)} | A = [\tilde{\phi}(v^1), \dots, \tilde{\phi}(v^{n-1})], v \in \Gamma^N \}$, and $NP(v^k)$ the set of null players in $v^k \in \Gamma^{N,k}$ under c_v for every $k \in \{1, \dots, n-1\}$.

Then, we define an indicator function $\gamma:\pm\tilde{\phi}(\bigcup_{k=1}^{n-1}\Gamma^{N,k})\to\{0,1\}^{n-1}$ as follows:

$$\gamma(\tilde{\phi}(v^p)) = \begin{cases} 0, & \text{if } \left(NP(v^{p-1}) = \emptyset \text{ and } NP(v^p) \neq \emptyset\right), \\ \text{or } \left(\gamma(\tilde{\phi}(v^{p-1})) = 0 \text{ if } p \geq 2\right) \\ 1, & \text{otherwise.} \end{cases}$$
 (27)

for all $p \in \{1, ..., n-1\}$.

The proposed function γ states that examinations of coalitional worth of a game are terminated if some players are turned out to be null players. Assume that we use $\tilde{\phi}$ as the allocation rule. From the definition of a null player, if null players appear in a game at some stage, then worth of the grand coalition equals to worth of some coalition whose cardinality is k. That is, if we use the proposed stopping rule, then we examine coalitional worth as many as possible. Next, we axiomatize the proposed indicator function. Let $\pi: \sigma(\bigcup_{k=1}^{n-1} \Gamma^{N,k}) \to \mathbb{R}$

Next, we axiomatize the proposed indicator function. Let $\pi : \sigma(\bigcup_{k=1}^{n-1} \Gamma^{N,k}) \to \{0,1\}^{n-1}$. Let \overline{x} be the negation of a boolean value x and $x \vee y$ represents $\max\{x,y\}$ where x and y are boolean values.

Assume that
$$\sigma(v^0) = (\frac{v(N)}{n}, \dots, \frac{v(N)}{n}).$$

Axiom 6 Let $v \in \Gamma^N$. If $\sigma_i(v^k) = \sigma_i(v^{k-1})$ for all $i \in N$, then $\pi(\sigma(v^k)) = 1$ for every $k \in \{1, \ldots, n-1\}$.

Axiom 6 states that if the values of all players at Stage k are same as those at Stage k-1, then an examination of coalitional worth continues.

Axiom 7 Let $v_1, v_2 \in \Gamma^N$ and $k \in \{1, ..., n-2\}$. If $NP(v_1^k \pm v_2^k) = NP(v_1^k) \cap$ $NP(v_2^k)$, then the following holds:

$$\pi(\sigma(v_1^k \pm v_2^k)) = \begin{cases} \overline{\pi(\sigma(v_1^k))} \vee \overline{\pi(\pm \sigma(v_2^k))}, & \text{if } NP(v_1^k) \cap NP(v_2^k) = \emptyset, \\ \pi(\sigma(v_1^k)) \vee \pi(\pm \sigma(v_2^k)), & \text{otherwise.} \end{cases}$$
(28)

Axiom 7 states that an indicator function of a solution for a game can be represented by the indicator functions of its component games using logical operations.

Axiom 8 Let $v_1, v_2 \in \Gamma^N$ and $k \in \{1, ..., n-1\}$. If $NP(v_1^k - v_2^k) \supset NP(v_1^k) \cap$ $NP(v_2^k)$, then the following holds:

$$\pi(\sigma(v_1^k - v_2^k)) = 0. (29)$$

Axiom 8 states that if new null players appear in $v_1^k - v_2^k$, then examinations of coalitional worth are finished.

Axiom 9 Let
$$v \in \Gamma^N$$
. If $\pi(\sigma(v^{k-1})) = 0$, then $\pi(\sigma(v^k)) = 0$.

Axiom 9 states that if examinations are finished, then there is no possibility to resume it.

Theorem 2. Let $\sigma = \pm \tilde{\phi}$. Then, γ is the unique function on $\sigma(\bigcup_{k=1}^{n-1} \Gamma^{N,k})$ that satisfies Axioms 6 through 9.

Proof. First, we show that γ satisfies Axioms 6 through 9. It is clear that γ satisfies Axiom 6, 8, and 9.

We show that γ satisfies Axiom 7. Let $v_1, v_2 \in \Gamma^N$.

(i) When $NP(v_1^k) \cap NP(v_2^k) = \emptyset$ such that $k \in \{1, \ldots, n-2\}$:

We consider the case $\gamma(\tilde{\phi}(v_1^k)) = 0$ and $\gamma(\tilde{\phi}(v_2^k)) = 0$ holds. In this case,
$$\begin{split} \tilde{\phi}_i(v_1^k) &= 0 \text{ holds for all } i \in NP(v_1^k), \text{ and } \tilde{\phi}_i(v_2^k) = 0 \text{ holds for all } i \in NP(v_2^k). \\ \text{However, since } NP(v_1^k \pm v_2^k) &= NP(v_1^k) \cap NP(v_2^k) = \emptyset, \ \tilde{\phi}_i(\underline{v_1^k \pm v_2^k})) \neq 0 \text{ for all } i \in N. \text{ That is, } \gamma(\tilde{\phi}(v_1^k \pm v_2^k)) = 1 \text{ holds and } \gamma(\tilde{\phi}(v_1^k \pm v_2^k)) = \overline{\gamma}(\sigma(v_1^k)) \vee \overline{\gamma}(\pm \sigma(v_2^k)) \end{split}$$
holds.

Next, we consider the case $\gamma(\tilde{\phi}(v_1^k)) = 1$ and $\gamma(\tilde{\phi}(v_2^k)) = 0$ holds. In this case, from $NP(v_1^k) = \emptyset$, $\tilde{\phi}_i(v_1^k) \neq 0$ for all $i \in N$. That is, $\tilde{\phi}_i(v_1^k \pm v_2^k) =$ $\tilde{\phi}_i(v_1^k) \pm \tilde{\phi}_i(v_2^k) \neq 0$ for all $i \in N$ and $\gamma(\tilde{\phi}(v_1^k \pm v_2^k)) = 1$ holds. (ii) When $NP(v_1^k) \cap NP(v_2^k) \neq \emptyset$ or $NP(v_1^k) = NP(v_2^k) = \emptyset$ such that

 $k \in \{1, \dots, n-2\}$:

We consider the case $\gamma(\tilde{\phi}(v_1^k)) = 0$ and $\gamma(\tilde{\phi}(v_2^k)) = 0$ holds. In this case, from $NP(v_1^k \pm v_2^k) = NP(v_1^k) \cap NP(v_2^k) \neq \emptyset, \ \gamma(\tilde{\phi}(v_1^{k'} \pm v_2^k)) = 0 \text{ holds. That is,}$ $\gamma(\tilde{\phi}(v_1^k \pm v_2^k)) = \gamma(\sigma(v_1^k)) \vee \gamma(\pm \sigma(v_2^k))$ holds.

Next, we consider the case $\gamma(\tilde{\phi}(v_1^k)) = 1$ and $\gamma(\tilde{\phi}(v_2^k)) = 1$ holds. In this case, since $NP(v_1^k) = NP(v_2^k) = \emptyset$, $NP(v_1^k \pm v_2^k) = \emptyset$ holds. That is, $\gamma(\tilde{\phi}(v_1^k \pm v_2^k)) = 1$

From (i) and (ii), γ satisfies Axiom 7.

Next, we show the uniqueness. Let $\pi: \pm \tilde{\phi}(\bigcup_{k=1}^{n-1} \Gamma^{N,k}) \to \{0,1\}^{n-1}$. For every $S \subseteq N$, let $u_S \in \Gamma^N$ be a unanimity game and $c_S \in \mathbb{R} \setminus \{0\}$. Let k < |S|. Then from the definition of $\tilde{\phi}$, the following holds:

$$\tilde{\phi}_i(c_S u_S^k) = \frac{c_S}{n} \quad \forall i \in N.$$
(30)

Otherwise, that is $k \geq |S|$, the following holds:

$$\tilde{\phi}_i(c_S u_S^k) = \begin{cases} \frac{c_S - c_{c_S u_S}(k)}{|S|}, & \text{if } i \in S, \\ 0, & \text{otherwise.} \end{cases}$$
 (31)

Here, notice that $c_{c_{SUS}}(k) = 0$ holds if $S \neq N$ from Assumption 2 and 4 of a

Therefore, from Axiom 6 and Assumption 2, 3, and 4 of a cost function, the following holds:

$$\pi(\tilde{\phi}(c_S u_S^k)) = \begin{cases} 1, & \text{if } k \in \{1, 2, \dots, n-2\}, \\ 0, & \text{if } k = n-1. \end{cases}$$
 (32)

Let $v \in \Gamma^N$. Since $\tilde{\phi}$ satisfies Additivity, the following holds:

$$\pi(\tilde{\phi}(v^k)) = \pi(\sum_{S \subseteq N} \tilde{\phi}(c_S u_S^k))$$
(33)

$$= \pi \left(\sum_{S \subseteq N, c_S > 0} \tilde{\phi}(c_S u_S^k) + \sum_{S \subseteq N, c_S < 0} \tilde{\phi}(c_S u_S^k) \right)$$
(34)

such that $c_S \in \mathbb{R} \setminus \{0\} \ \forall S \subseteq N$, for all $k \in \{1, 2, ..., n-1\}$.

From Axiom 7, we have:

$$\pi(\sum_{S\subseteq N, c_S>0} \tilde{\phi}(c_S u_S^k)) = \begin{cases} \overline{\pi(\sum_{S\in P(N)\backslash S_1, c_S>0} \tilde{\phi}(c_S u_S^k))} \vee \overline{\pi(\tilde{\phi}(c_{S_1} u_{S_1}^k))}, \\ \text{if } NP(\sum_{S\in P(N)\backslash S_1, c_S>0} c_S u_S^k) \cap NP(c_{S_1} u_{S_1}) = \emptyset, \\ \pi(\sum_{S\in P(N)\backslash S_1, c_S>0} \tilde{\phi}(c_S u_S^k)) \vee \pi(\tilde{\phi}(c_{S_1} u_{S_1}^k)) \\ \text{otherwise.} \end{cases}$$

Continuing this calculation, $\pi(\sum_{S\subseteq N, c_S>0} \tilde{\phi}(c_S u_S^k))$ can be represented by using \forall of $\pi(\tilde{\phi}(c_S u_S^k))$ for all $S\subseteq N$ such that $c_S>0$. Similarly, $\pi(\sum_{S\subseteq N, c_S<0} \tilde{\phi}(c_S u_S^k))$

can be represented by using \vee of $\pi(\tilde{\phi}(c_S u_S^k))$ for all $S \subseteq N$ such that $c_S < 0$.

Finally, if
$$NP(\sum_{S\subseteq N, c_S>0} c_S u_S^k) \cap NP(\sum_{S\subseteq N, c_S<0} c_S u_S^k) = NP(\sum_{S\subseteq N, c_S>0} c_S u_S^k + \sum_{S\subseteq N, c_S<0} c_S u_S^k)$$
, then $\pi(\sum_{S\subseteq N, c_S>0} \tilde{\phi}(c_S u_S^k) + \sum_{S\subseteq N, c_S<0} \tilde{\phi}(c_S u_S^k))$ is obtained uniquely from Axiom 7.

If $NP(\sum_{S\subseteq N, c_S>0} c_S u_S^k) \cap NP(\sum_{S\subseteq N, c_S<0} c_S u_S^k) \neq NP(\sum_{S\subseteq N, c_S>0} c_S u_S^k)$

$$-\sum_{S\subseteq N, c_S<0} -c_S u_S^k)$$
, then $NP(\sum_{S\subseteq N, c_S>0} c_S u_S^k) \cap NP(\sum_{S\subseteq N, c_S>0} c_S u_S^k)$

$$-\sum_{S\subseteq N, c_S<0} -c_S u_S^k) \supset NP(\sum_{S\subseteq N, c_S>0} c_S u_S^k) \cap NP(\sum_{S\subseteq N, c_S<0} c_S u_S^k)$$
 holds since a player is a null player in $v_1^k - v_2^k$ if he is a null player in v_1^k and v_2^k for any $v_1^k, v_2^k \in \Gamma^{N,k}$.

Therefore, $\pi(\sum_{S\subseteq N, c_S>0} \tilde{\phi}(c_S u_S^k) - \sum_{S\subseteq N, c_S<0} \tilde{\phi}(-c_S u_S^k)) = 0$ from Axiom 8.

That is, $\pi(\tilde{\phi}(v^k))$ is obtained uniquely for any $v^k \in \Gamma^{N,k}$ for all $k \in \{1, \dots, n-1\}$.

This completes the proof. \square

Example 2. Using the result of calculation of Example 1, the $\gamma(\tilde{\phi}(v^k))$ for all $k \in \{1, 2, 3\}$ is obtained as follows:

$$\gamma(\tilde{\phi}(v^1)) = 1, \ \gamma(\tilde{\phi}(v^2)) = 1, \ \gamma(\tilde{\phi}(v^3)) = 0.$$
 (36)

6 Concluding Remarks

In this paper, we considered a new problem of cooperative game theory that is called a partially defined game with payments. We proposed and axiomatized the Shapley value for PDGs with payments and an exiting rule which indicates when we should stop examinations of coalitional worth. It might be possible to propose exiting rules other than the rule we propose in this paper. For instance, it can be proposed that examinations of coalitional worth are not performed at all. That is, if we use this exiting rule, then we have to decide allocations of payoffs in the situation that only the worth of the grand and singleton coalitions are known. If we use $p\tilde{h}i$ as the solution for PDGs with payments, then the allocation rule coincides with the CIS value [4]. Many rules other than this rule and our rule may be proposed in future research with their axiom system.

References

- 1. Aguilera, NE and Di Marco, SC and Escalante, MS (2010). The Shapley value for arbitrary families of coalitions. European Journal of Operational Research 204: 125-138.
- 2. Albizuri, MJ and Masuya, S and Zarzuelo, JM (2022) Characterization of a value for games under restricted cooperation. Ann. Oper. Res. 318: 773–785.

- 3. Algaba, E and Bilbao, JM and Borm, P and López, JJ (2001) The Myerson Value for Union Stable Structures. Mathematical Methods of Operations Research 54: 359–371.
- 4. Driessen, TSH and Funaki, Y (1991) Coincidence of and collinearity between game theoretic solutions. OR Spektrum 13:15–30.
- 5. Calvo, E and Gutiérrez-López, E (2015) The value in games with restricted cooperation. Discussion Papers in Economic Behavior 01/15. University of Valencia.
- Černý, M and Grabisch, M (2024) Incomplete cooperative games with playercentered information. Discrete Applied Mathematics 346: 62–79.
- 7. Harsanyi, JC (1963) A simplified bargaining model for the n-person cooperative game. Int. Econ. Rev. 4: 194–220.
- 8. Myerson, RB (1977) Graphs and Cooperation in Games. Mathematics of Operations Research 2: 225–229.
- 9. Myerson, RB (1980) Conference Structures and Fair Allocation Rules. International Journal of Game Theory 9: 169–182.
- Owen, G (1977) Values of games with a priori unions. In: Henn, R and Moeschlin,
 O (eds) Mathematical Economics and Game Theory. Springer 76–88.
- Shapley, LS (1953) A value for n-person games. In: Kuhn, H and Tucker, A (eds) Contributions to the Theory of Games II. Princeton University Press 307–317.
- Willson, SJ (1993) A value for partially defined cooperative games. Int. J. Game Theory 21: 371–384.
- 13. Yu, X (2021) Extension of Owen value for the game with a coalition structure under the limited feasible coalition. Soft Comput. 25: 6139–6156.