

A SPARSITY TEST FOR MULTIVARIATE HAWKES PROCESSES

ANTOINE LOTZ

PARIS-DAUPHINE UNIVERSITY, PSL, Paris, France
EDF-LAB, EDF R&D, Palaiseau, France

ABSTRACT. Multivariate Hawkes processes (MHP) are a class of point processes in which events at different coordinates interact through mutual excitation. The weighted adjacency matrix of the MHP encodes the strength of the relations, and shares its support with the causal graph of interactions of the process. We consider the problem of testing for causal relationships across the dimensions of a marked MHP. The null hypothesis is that a joint group of adjacency coefficients are null, corresponding to the absence of interactions. The alternative is that they are positive, and the associated interactions do exist. To this end, we introduce a novel estimation procedure in the context of a large sample of independent event sequences. We construct the associated likelihood ratio test and derive the asymptotic distribution of the test statistic as a mixture of χ^2 laws. We offer two applications on financial datasets to illustrate the performance of our method. In the first one, our test reveals a deviation from a static equilibrium in bidders' strategies on retail online auctions. In the second one, we uncover some factors at play in the dynamics of German intraday power prices.

Mathematics Subject Classification (2020): 62F03, 62F12, 62F30, 60G55

Keywords: Parametric estimation; Non standard likelihood theory, Likelihood ratio test; Multivariate Hawkes processes

1. INTRODUCTION

1.1. Motivation. Hawkes processes are self-exciting point processes in which the probability of an event appearing increases in the aftermath of past occurrences. Multivariate Hawkes processes (MHP) extend this property to interdependent sequences, wherein events in a coordinate of the process are also contingent upon the past of other coordinates. They provide an ideal model for the dynamics of interacting systems and have demonstrated this capacity in diverse applications ranging from neuroscience [39] [7], to high-frequency financial data [5], or information propagation on media networks [10]. In this work, we consider the problem of testing for temporal influence between the coordinates of a marked MHP. The hypotheses for our test express naturally in terms of characteristics of the process, and we informally introduce the relevant features of the model hereafter.

Suppose one observes a K -th dimensional counting process $(\mathbf{N}_t) = (N_{k,t})$ over a fixed time-frame $[0, T]$. The jumps of the k -th coordinate occur at distinct times (t_i^k) with $1 \leq i \leq N_T^k$, and to each jump time t_i^k is associated a mark X_i^k with values in \mathbb{R}^m . The law of (\mathbf{N}_t) is characterised by its predictable intensity process $(\lambda_{1,t}, \dots, \lambda_{K,t})$, where each $\lambda_{k,t}$ can be thought of as the instantaneous probability of an event arriving at coordinate k at time t . In this paper, we are interested in the class of linear marked MHPs, which describes processes with intensity of the form

E-mail address: antoine.lotz@dauphine.eu.

$$(1) \quad \lambda_{k,t} = \mu_k \left(\frac{t}{T} \right) + \sum_{k=1}^K \sum_{t_i^l < t} \varphi_{kl}(t - t_i^l, X_i^l),$$

where the $\varphi_{kl}: [0, \infty) \times \mathbb{R}^m \mapsto [0, \infty)$ are integrable functions referred to as kernels of the Hawkes process. The intensity process (1) is constructed so that an event at coordinate l at time s increases the intensity at coordinate k at time t by $\varphi_{kl}(t - s)$. The influence between different coordinates of the process are thus encoded into $\varphi = (\varphi_{kl})$, and selecting interactions simply amounts to selecting null kernels with the question typically phrased in terms of the L^1 norms of the kernels. To this aim, a significant effort has been dedicated to penalization-based approaches, with a variety of criteria being successfully developed for least-square (Hansen *et al.* [23], Bacry *et al.* [5]), and maximum likelihood estimation (Zhou *et al.* [52], Xu *et al.* [51], Salehi [41], Sulem *et al.* [46] and Goda [20]). Testing for causal influence is a natural adjacent question to variable selection for (possibly marked) MHPs. Yet, the topic seems to have attracted a more modest interest. The only available test is a multiple comparison procedure introduced by Bonnet *et al.* [7] as a by-product of their estimation method for inhibiting MHPs. Such methods however incur a change in statistical power and the full theoretical background for their results remains an open question.

1.2. Contribution and organisation of the paper. Given a subset $\mathcal{J} \subset \llbracket 1, K \rrbracket^2$ of coordinate pairs, we construct the likelihood-ratio test for the null hypothesis

$$(2) \quad \int_0^\infty \varphi_{kl}(s) ds = 0, \text{ for every } (k, l) \in \mathcal{J},$$

against the one-sided alternative

$$(3) \quad \int_0^\infty \varphi_{kl}(s) ds > 0, \text{ for every } (k, l) \in \mathcal{J}.$$

Our configuration differs from the usual situation where a single long time trajectory of the MHP is recorded. We observe instead n independent realisations of (N_t) over $[0, T]$, where T is fixed and n large. Such data structures are commonly found in biology (see for example Reynaud-Bouret [39]) and they are well fitted to the financial applications we have in mind, which involve short-maturity illiquid products (as will be detailed further down this section). Following an aggregation procedure, this setting results in the large baseline regime introduced by Chen & Hall [9] in the univariate case. More precisely, we work in the parametric setting where μ and φ depend on some $\theta \in \Theta \subset \mathbb{R}^p$ and we are concerned with processes which intensities take the form

$$\lambda_{k,t}^{(n)}(\theta) = n\mu_k \left(\frac{t}{T}, \theta \right) + \sum_{k=1}^K \sum_{t_i^l < t} \varphi_{kl}(t - t_i^l, X_i^l, \theta).$$

The necessary assumptions are detailed in section 2 along a rigorous definition of the MHP.

Our main result is Theorem 1 in section 3, stating the convergence of the likelihood ratio toward a difference of squared distances from a normal variable to two sub-spaces depending on (2) and (3). Going from the univariate to the multivariate incurs some specific new difficulties, as the kernels possibly being null translate into the true parameter lying at a boundary of Θ . In turn, this gives rise to singularities in the statistics of interest. The asymptotic distribution of the likelihood ratio is in particular defined as a chi-bar ($\bar{\chi}^2$) law, as introduced by Kudo [33] and Shapiro [43]. The $\bar{\chi}^2$ arises naturally from constraints on a parameter space and is a recurrent occurrence in

statistics of random processes, see for instance Andrews [2] [3] for an application to GARCH models. We refer also to Silvapulle & Sen [45] for a comprehensive review on constrained statistical inference. In our case, the $\bar{\chi}^2$ expresses as a mixture of standard χ^2 laws. Though the weights are generally intractable, we show they have a simple expression in most of the low-dimensional use cases. In particular, our procedure yields a straightforward test for the Poisson *versus* self-exciting problem introduced by Dachian & Kutoyants [13] in a different setting. In the general case, the evaluation of the weights can be entirely foregone in favour of a conditional test based on Susko's [47] method for $\bar{\chi}^2$ -tests, with negligible power loss. As a corollary of our results and still in section 3, we recover the consistency and asymptotic distribution of the maximum likelihood estimator (MLE), answering a question of Chen & Hall [9] regarding the extension of their work to MHPs. Hypotheses of the type (2) describe the most general forms for the parameter space, and are in many cases a desirable property. The non-regular distribution of the MLE is in this sense a generic feature of the linear MHP. While the boundary parameter issue has been raised in other works (see Goda [20] for the particular example of an exponential MHP in a long time regime), the derivation of the entire asymptotic distribution of the MLE and likelihood ratio is a seemingly new result.

In section 4 we provide ample numerical illustrations of our procedure at work, including both simulation studies and applications. In a first example, we show how our test may be used to infer agents' behaviour in online auctions. Specifically, we provide evidence that interactions between agents are a significant factor in bid arrivals, therefore shifting the transaction price away from the static Vickrey equilibrium. In a second example, we study the price formation of hourly power futures on the German intraday market. We find that in addition to the reversion classically accounted for by MHP-based models in finance, the price mechanics display a form of inertia. We offer an explanation to our finding in terms of liquidity constraints. Our two real-world examples pertain to illiquid assets. The configuration of sparsely populated event sequences has been the subject of a relatively limited body of work (see Salehi [41]), and this type of application had consequently remained at the margin of the range of MHPs-based model. Our examples demonstrate the good performance of our approach in this context.

In section 5 are the proofs. Before moving to our assumptions, we provide a brief overview of our strategy. We follow a classical M-estimator approach. The initial steps of the proof involve showing the convergence of the log-likelihood after appropriate normalisation and centering. This requires proving the law of large number

$$(4) \quad \frac{1}{n} \{ \lambda_t^{(n)}(\theta) - \mathbb{E}[\lambda_t^{(n)}(\theta)] \} = o_{\mathbb{P}}(1)$$

holds uniformly in t . Up to some technicalities, the proof of (4) does not require specific adjustments to deal with the specificities of the multivariate case. The main complication stems instead from satisfying the conditions for Geyer's [19][Theorem 4.4] constrained M-estimator master theorem, which is the key to the weak convergence of the likelihood ratio. In particular, we require a stochastic equicontinuity condition for the second order remainder of the log-likelihood, as classically introduced by Pollard [37][Chapter VII, section I]. We show that verifying this assumption can be reduced to precisising the rate of convergence in (4), in the sense that it suffices to have

$$(5) \quad \frac{1}{\sqrt{n}} \{ \lambda_t^{(n)}(\theta) - \mathbb{E}[\lambda_t^{(n)}(\theta)] \} = O_{\mathbb{P}}(1)$$

uniformly in t and θ . The proof for (5) is slightly more involved than that of (4). We proceed in the manner of Pollard [37][Chapter VII, section II] and use a chaining technique to achieve the uniformity in θ . To this end, we need a sub-exponential bound on the probability that the left-hand side of (5) deviates as θ moves. A fitting bound is achieved using a martingale concentration inequality due to van de Geer [49][Lemma 2.1] among others. We can then apply Geyer's [19] result, yielding theorem 1 and the rest of our results.

2. MODEL AND ASSUMPTIONS

2.1. Setting and notation. Consider a rich enough probability space on which, for each $k \in \llbracket 1, K \rrbracket$, the event times (t_i^k) introduced above are defined as a sequence of increasing positive random variables and required to be distinct, meaning that jumps as different coordinates may not occur simultaneously. To each event time t_i^k is associated a mark (X_i^k) with values in \mathbb{R}^m and law $F^k(dx)$. The (X_i^k) form an i.i.d collection of random variables. The multivariate counting process $(\mathbf{N}_t) = (N_{1,t}, \dots, N_{K,t})$ for the (t_i^k) is then defined for any $t \geq 0$ and $k \in \llbracket 1, K \rrbracket$ as

$$N_{k,t} = \sum_{i \in \mathbb{N}} \mathbb{1}_{\{t_i^k \leq t\}},$$

and the counting measure $N_k(ds, dx)$ is the random Borel measure from $\mathcal{B}(\mathbb{R}^+) \otimes \mathcal{B}(\mathbb{R})^{\otimes m}$ to \mathbb{R}^+

$$N_k : A \times B \mapsto \sum_{i \in \mathbb{N}} \mathbb{1}_{\{t_i^k \in A\} \cap \{X_i^k \in B\}}.$$

Denote by $\mathbb{F} = (\mathcal{F}_t)$ the natural filtration of (\mathbf{N}_t) and consider a \mathbb{F} -predictable coordinate-wise positive process $\lambda_t = (\lambda_{1,t}, \dots, \lambda_{K,t})$. From Jacod [27], under mild technical assumptions we may choose our probability space such that (\mathbf{N}_t) admits $(\mathbf{\Lambda}_t) = (\int_0^t \lambda_s ds)$ as its compensator, that is, as the unique predictable increasing process such that $(\mathbf{N}_t - \mathbf{\Lambda}_t)$ is a \mathbb{F} -local martingale. We refer to the comprehensive textbook of Daley & Vere-Jones [14] for more details on processes with stochastic intensity. This equips us with a characterisation of the (marked) MHP.

Definition 1 (multivariate marked Hawkes process). *Let $T > 0$. The process $(\mathbf{N}_t)_{t \in [0, T]}$ is said to be a multivariate marked Hawkes process if, for any $t \in [0, T]$, its intensity takes the form*

$$(\lambda_{k,t})_k = \left(\mu_k \left(\frac{t}{T} \right) + \sum_{l=1}^K \int_{[0,t) \times \mathbb{R}^m} \varphi_{kl}(t-s, x) N_l(ds, dx) \right)_k,$$

concisely written,

$$\lambda_t = \mu \left(\frac{t}{T} \right) + \int_{[0,t) \times \mathbb{R}^m} \varphi(t-s, x) \mathbf{N}(ds, dx),$$

where $(\mu_k) : [0, 1] \mapsto (0, \infty)^K$ is the baseline, and $(\varphi_{kl}) : \mathbb{R}^+ \times \mathbb{R}^m \mapsto \mathcal{M}_K(\mathbb{R}^+)$ is the kernel of the process.

Now denote by $\mathbf{M}(ds, dx) = (M_k(ds, dx))$ the compensated measure

$$\mathbf{M}(ds, dx) := N_k(ds, dx) - \lambda_{k,s} ds F_k(dx).$$

Define also $d\mathbf{N}_s = \mathbf{N}(ds, \mathbb{R}^m)$, and $d\mathbf{M}_s = \mathbf{M}(ds, \mathbb{R}^m)$. Then, from Definition 1,

$$(\mathbf{M}_t) = \left(\int_0^t d\mathbf{M}_s \right), \quad t \in [0, T],$$

is a local martingale. While Jacod's [27] theorem suffices to guarantee the existence of (\mathbf{N}_t) , note that one can always explicitly construct the MHP such that the no simultaneous jump condition

we required above is satisfied. See for instance Hawkes & Oakes [26] and Delattre *et al.* [15] for two equivalent representations, respectively as a cluster process and a Poisson SDE solution. The jumps being distinct entails some useful simplification for the variation of the local martingale (\mathbf{M}_t) , as transcribed in remark 1 below.

Remark 1. *The quadratic variation $[\mathbf{M}, \mathbf{M}]$ and predictable variation $\langle \mathbf{M}, \mathbf{M} \rangle$ of (\mathbf{M}_t) are both diagonal and respectively given by $\text{Diag}(N_{k,t})$ and $\text{Diag}(\int_0^t \lambda_k(s) ds)$.*

We need more notation. Throughout, we denote by $\|\cdot\|_p$ the ℓ^p -norm on \mathbb{R}^K and by $\|\cdot\|_{L^1}$ the $L^1(\mathbb{R}^+)$ -norm on the positive integrable functions $\|\phi\|_{L^1} = \int_0^\infty \phi(s) ds$. The set of squared real $K \times K$ matrices is labelled $\mathcal{M}_K(\mathbb{R})$ and for any $M \in \mathcal{M}_K(\mathbb{R})$, $\rho(M)$ is the spectral radius of M . For any open set $U \subset \mathbb{R}$, the set $\mathcal{C}^0(U)$ designates the continuous functions from U to \mathbb{R} , and likewise $\mathcal{C}^1(U)$ designates the differentiable functions from U to \mathbb{R} . Finally, for any $f = (f_{kl}): \mathbb{R}^m \mapsto \mathcal{M}_K(\mathbb{R}^+)$, we employ the notation

$$F(dx) = \text{Diag}(F^j(dx)) \quad \text{and} \quad \langle F, f \rangle = \int_{\mathbb{R}^m} f(\cdot, x) F(dx),$$

and write $f \in L^2(F(dx))$ when $\forall i, j \in \llbracket 1, K \rrbracket^2 \int_{\mathbb{R}^m} f_{ij}^2(x) F^j(dx) < \infty$.

2.2. Estimation procedure and assumptions. As mentioned in the introduction, our setting differs from the usual observation of a single Hawkes process. We suppose instead that n independent realisations $(\mathbf{N}_t^1), \dots, (\mathbf{N}_t^n)$ of the same Hawkes process with baseline μ and kernel φ are recorded on a fixed time interval $[0, T]$. Now, by linearity of the intensity the sum process

$$(6) \quad (\mathbf{N}_t^{(n)}) := \left(\sum_{i=1}^n \mathbf{N}_t^i \right), \quad t \in [0, T],$$

is also a Hawkes process and admits $n\mu$ and φ as its baseline and kernel.

Remark 2. *This specificity of linear MHPs can be linked to the cluster representation of Hawkes & Oakes [26]. Summing up linear Hawkes processes is equivalent to aggregating on the same timeline independent clusters from different realisations. Since cluster arrivals are Poissonian and thus additive, this leads to the baseline of the resulting process being the sum of the baseline of each realisation.*

We therefore recover Chen & Hall [9]'s framework of a baseline tending to infinity. Though their work does not appear to take advantage of the previous observation, the univariate results they introduce extend immediately to our configuration through the aggregation device (6), and we construct our assumptions as minimal extension of their own. Following the preceding remarks, we work in a parametric setting and we consider a collection $\mathbf{N}^{(n)}$ of marked MHPs indexed by $n \in \mathbb{N}^*$, with intensities

$$(7) \quad \lambda_t^{(n)}(\theta) = n\mu \left(\frac{t}{T}, \theta \right) + \int_0^t \varphi(t-s, x, \theta) \mathbf{N}^{(n)}(ds, dx), \quad t \in [0, T].$$

under the probability measure $\mathbb{P}(\theta)$ where θ lies in a compact subset $\Theta \subset \mathbb{R}^d$ with non empty interior. To simplify the expression for our test, we assume θ separates into $\theta = (\gamma, \alpha, \beta)$ where μ depends only on $\gamma \in \mathbb{R}^K$ and φ expresses as

$$(8) \quad \varphi_{kl}(\cdot, \cdot, \theta) = \alpha_{kl} \Phi_{kl}(\cdot, \cdot, \beta),$$

where $\alpha \in \mathcal{M}_K(\mathbb{R}^+)$ and for k, l and any value of β

$$(9) \quad \int_{[0, \infty) \times \mathbb{R}^m} \Phi_{kl}(s, x, \beta) F_l(dx) ds = 1.$$

Expression (8) and (9) are consistent with parameterisations found in the literature (see for instance Ogata [36] or Hardiman [25]). The support of the matrix α , that is, the matrix $h = (h_{kl}) \in \mathcal{M}_K(\{0, 1\})$ with $h_{kl} = 0 \iff \alpha_{kl} = 0$, is then known as the adjacency of the MHP (see for instance Sulem [46]). Accordingly, α is known as its weighted adjacency (see Bacry *et al.* [5]). The null hypothesis conveniently re-phrases in terms of the matrix parameter α as

$$(10) \quad \alpha_{kl} = 0, \text{ for every } (k, l) \in J,$$

and the alternative as

$$(11) \quad \alpha_{kl} > 0, \text{ for every } (k, l) \in J.$$

We will use formulations (2), (3) and (10),(11) interchangeably.

Remark 3. *The notion of "adjacency" is chosen to reflect a salient property of MHPs in terms of graphical models. Using the definition of Granger-causality for point processes introduced by Didelez [17], Xu *et al.* [51] show that an arrow exists from coordinate l to coordinate k in the causal graph of the MHP if and only if $\alpha_{kl} > 0$. In this regard, our test can be understood as a causality test.*

Additionally to the parametric specification above, the baseline and kernel are subject to the following assumptions.

Assumption 1.

- (i) For any $\theta \in \Theta$, $k \in \llbracket 1, K \rrbracket$ and $t \in [0, 1]$, $\mu_k(t, \theta) > 0$.
- (ii) For any $\theta \in \Theta$ and $k \in \llbracket 1, K \rrbracket$, $\mu_k(\cdot, \theta) \in \mathcal{C}^0([0, 1])$.

Assumption 2. *There exist two unique functions f, g on \mathbb{R}^+ with values in $\mathcal{M}_K(\mathbb{R}^+)$ such that, for any $s \in \mathbb{R}^+$, $x \in \mathbb{R}^m$ and $\theta \in \Theta$,*

$$(12) \quad \varphi(s, x, \theta) = g(x, \theta) \odot f(s, \theta),$$

where \odot denotes the Hadamard (or coefficient-wise) product over $\mathcal{M}_K(\mathbb{R}^+)$.

Remark 4. *The uniqueness of decomposition (12) is not a benign assumption, it de facto excludes non identifiable choices of the type $g(\gamma, x)\varphi(s, \alpha, \beta) = \alpha\gamma x\phi(s, \beta)$. Conversely, the Hadamard product is only meant to reproduce commonly found parameterisations (see for instance Bacry *et al.* [5]). Other choices would not alter our results, including the usual matrix product.*

Assumption 3.

- (i) For any $\theta \in \Theta$, $f(\cdot, \theta) \in \mathcal{C}^1(\mathbb{R}^+)$.
- (ii) The function f is uniformly equi-continuous in θ .

Assumption 4.

- (i) For any $\theta \in \Theta$, $g(\cdot, \theta) \in L^2(F(dx))$.
- (ii) The function g is uniformly equi-continuous in θ .

Remark 5. *Any kernel of the type*

$$(13) \quad \varphi(s, x, \theta) = g(x) \odot f(s, \theta)$$

where $g \in L^2(F(dx))$ and f is a \mathcal{C}^1 function over $\mathbb{R}^+ \times \Theta$ will satisfy assumptions 3 and 4. In particular, fixing $g \equiv 1$ retrieves Chen & Hall's [9] original setup, and our assumptions can be regarded as a natural extension of the univariate case's conditions in this respect. While theoretically restrictive, the splitting Assumption (2) encompasses most of the existing literature regarding applications of marked Hawkes process;

with use cases including seismology [36], information diffusion [10], epidemiology [12], and price dynamics on financial markets [34][16] [11]. The choice of i.i.d marks is a more limiting Assumption. Embrecht et al.'s model [18] features a split kernel but remains out of the scope of this article since its marks exhibit a dynamic distribution.

We also require some identifiability conditions on μ and φ .

Assumption 5. If $\vartheta \neq \theta$, then there exist a sub-interval $J \subset [0, T]$ and a subset $S \subset \mathbb{R}^m$ of non null-measures on which, for every $(t, x) \in J \times S$, either $\mu(t, \vartheta) > \mu(t, \theta)$ or $\mu(t, \vartheta) < \mu(t, \theta)$ or $g(x, \vartheta)f(t, \vartheta) < g(x, \theta)f(t, \theta)$ or $g(x, \vartheta)f(t, \vartheta) > g(x, \theta)f(t, \theta)$.

As we move to the weak convergence of the LRS, we will be concerned with derivatives of the likelihood, requiring some higher order regularity conditions on the kernel φ .

Assumption 6.

- (i) For any $t \in [0, T]$, $\theta \mapsto \mu(t, \theta)$ is twice continuously differentiable over Θ .
- (ii) For any $\theta \in \Theta$, and any $p \in \{1, 2\}$, $t \mapsto \partial_\theta^p \mu(t, \theta) \in \mathcal{C}^1(\mathbb{R}^+)$.

Assumption 7.

- (i) For any $t \in [0, T]$, $\theta \mapsto f(t, \theta)$ is twice continuously differentiable over Θ .
- (ii) For any $\theta \in \Theta$, and any $p \in \{1, 2\}$, $t \mapsto \partial_\theta^p f(t, \theta) \in \mathcal{C}^1(\mathbb{R}^+)$.

Remark 6. Take note that the differentiability of f is not restricted to the interior of Θ , and we therefore require its right-differentiability at boundaries where some α_{kl} are null.

Assumption 8.

- (i) for any $x \in \mathbb{R}^m$, $\theta \mapsto g(x, \theta)$ is continuously differentiable over Θ .
- (ii) For any $q \in \{1, 2\}$, $\partial_\theta^q g$ is uniformly p -Hölder in θ for some $p \in (0, 1]$: there is some $C > 0$ such that, for any $x \in \mathbb{R}^m$ and any $\theta, \nu \in \Theta$

$$\|\partial_\theta^q g(x, \theta) - \partial_\theta^q g(x, \nu)\|_1 \leq C \|\theta - \nu\|_2^p$$

Remark 7. Hypothesis 8 drastically restricts the theoretical class of functions for g , but it is inconsequential in practice. In other works, it is often the case that g does not depend on θ , and the conditions above are then immediately met (see Deschatre and Gruet [16] and Goda [20] for two examples). When g does depend on θ , recalling the instantaneous probability interpretation of the intensity, a fitting choice is the logistic function

$$g(x, \theta) = \frac{1}{1 + e^{-\theta x}},$$

which derivative of order p in θ is $|x|^p$ -Lipschitz. Assumption 8 is thus easily satisfied in practice by truncating the marks, and the remark extends to most reasonable choices for g .

The next hypothesis is standard in the study of linear Hawkes processes, and notably suffices to guarantee the non-explosion of N_t (see for instance Bacry et al. [6][Lemma 1]).

Assumption 9. The process is stable in that for any θ in Θ

$$\rho \left(\int_{(0, \infty] \times \mathbb{R}^m} \varphi(s, x, \theta) ds F(dx) \right) < 1.$$

where ρ denotes the spectral radius.

Remark 8. In Hawkes & Oakes' [26] cluster representation, the univariate Hawkes process is an immigration-birth process in which migrants arrive at rate μ and have descendants according to a Galton-Watson process with branching ratio $\int_0^\infty \varphi(s) ds$. Assumption 9 then has the intuitive interpretation that the ratio of non-immigrants in the accumulated population cannot exceed 1. A popular choice for φ is the exponential parameterisation $t \mapsto \alpha \exp(-\beta t)$. The condition then expresses straightforwardly as $\alpha/\beta < 1$, and is comparably simple to verify for most kernel choices.

In our context, Assumption 9 guarantees a semi-analytic form of the normalized mean intensity, as exposed in lemma 1, which proof is straightforward and delayed to section 5

Lemma 1. Under Assumption 9, for any $\theta_0 \in \Theta$ and any $t \in [0, T]$

$$\frac{1}{n} \mathbb{E}^{\theta_0} [\lambda^{(n)}(t, \theta_0)] = h(t, \theta_0) := \mu\left(\frac{t}{T}, \theta_0\right) + \int_0^t \langle F, \Psi \rangle(t-s, \theta_0) \mu(s, \theta_0) ds,$$

where \mathbb{E}^θ is the expectation under $\mathbb{P}(\theta)$ and $\langle F, \Psi \rangle = \sum_{k \geq 1} \langle F, \varphi \rangle^{\star k}$, with $\star k$ the k -fold convolution.

Throughout, we denote by $H(\cdot, \cdot)$ the function

$$(14) \quad H: t, \theta \mapsto \int_0^t h(s, \theta) ds.$$

The definition of h being independent from the value of n allows to express the asymptotic information $I(\theta)$ of the model in a convenient form, as will be made clear in section 3. In the following, for any vector $u = (u_i)$ in some \mathbb{R}^d , we use the tensor involution notation $u^{\otimes 2}$ to denote the $d \times d$ matrix $u^{\otimes 2} = (u_i u_j) = uu^\top$.

Assumption 10. (Non degeneracy of the observed information)

The matrix valued function

$$(15) \quad I(\theta) = \int_0^T \sum_{k=1}^K \frac{\{\partial_\theta \mu_k(s, \theta) + \int_0^s \sum_{l=1}^K \partial_\theta \langle F, \varphi \rangle_{kl}(s-u, \theta) h_l(u, \theta) du\}^{\otimes 2}}{\mu(s, \theta) + \int_0^s \sum_{l=1}^K \langle F, \varphi \rangle_{kl}(s-u, \theta) h_l(u, \theta) du} ds$$

takes non singular values, where $h(t, \theta) = (h_k(t, \theta))$ is defined in Lemma 1.

Remark 9. Verifying assumption 10 relates to the parameter identifiability problem, which is a non trivial question for MHPs. consider for instance the exponential kernel

$$\varphi_{kl}: t \mapsto \alpha_{kl} \exp(-\beta_{kl} t).$$

for $(k, l) \in \llbracket 1, K \rrbracket^2$. It suffices that $\alpha_{kl} = 0$ in the true parameter to prevent β_{kl} from being identified. Under the non degeneracy assumption 10, the size and structure of the decay parameter are hence constrained by those of the weighted adjacency. We refer to Bonnet et al. [7] for a simple identifiability criterion in the context of exponential MHPs. In general, we provide a consistent estimator for $I(\theta)$ in Lemma 11 on which assumption 10 can be numerically verified.

2.3. Likelihood-based inference. The log-likelihood of $\theta \in \Theta$ for the (marked) MHP is given by

$$(16) \quad \mathcal{L}(n, \theta) = \sum_{k=1}^K \int_0^T \ln \lambda_{k,s}^{(n)}(\theta) dN_{k,s}^{(n)} - \int_0^T \lambda_{k,s}^{(n)}(\theta) ds,$$

where $(N_t^{(n)})$ is the process defined by (6). We refer to Daley & Vere-Jones [14][Proposition 7.2.3 p. 232] for a proof of (16). The maximum likelihood estimator $\hat{\theta}_n$ (MLE) for θ is then defined as the global argmax of the likelihood $\mathcal{L}(n, \cdot)$ over Θ . For any sub-model Θ_0 , we define similarly $\hat{\theta}_n^0$ as the global argmax of $\mathcal{L}(n, \cdot)$ over Θ_0 . Note that $\hat{\theta}_n^0$ writes as a random vector of \mathbb{R}^d with p null

coordinates. Our core problem is to derive the asymptotic distribution of the (log) likelihood ratio statistic (LRS)

$$\Lambda_n = 2 \left(\sup_{\theta \in \Theta} \mathcal{L}(n, \theta) - \sup_{\theta \in \Theta_0} \mathcal{L}(n, \theta) \right) = 2(\mathcal{L}(n, \hat{\theta}_n) - \mathcal{L}(n, \hat{\theta}_n^0))$$

for sub-models of the type

$$(17) \quad \Theta_0 = \{ \theta = (\gamma, \alpha, \beta) \in \Theta \mid \alpha_{kl} = 0; (k, l) \in \mathcal{J} \}$$

corresponding to the set of hypotheses (10) and (11). As is usual for likelihood ratio tests the supremum over the whole parameter space is used as a proxy for the alternative. The distribution of the MLE is of course also of great interest and we cover the problem as a by-product of our general results on Λ_n . Finally, we stress that the class of test we introduce is quite flexible, and Θ_0 may be chosen to match a larger set of problems.

3. MAIN RESULTS

3.1. Consistency of the MLE. Before getting to the likelihood ratio, a preliminary consistency result for the MLE is stated.

Proposition 1. *Under Assumptions 1 to 5 and Assumption 9 the MLE $\hat{\theta}$ is consistent: for any $\theta_0 \in \Theta$,*

$$(18) \quad \hat{\theta}_n \xrightarrow[n \rightarrow \infty]{\mathbb{P}(\theta_0)} \theta_0.$$

Observe that we have not introduced any restriction locating θ_0 in the interior $\overset{\circ}{\Theta}$ of its parameter space Θ . When the true parameter belongs to $\overset{\circ}{\Theta}$, the consistency of the MLE guarantees it eventually positions itself away from the border with probability tending to 1. We may then expect the same asymptotic behaviour as in the univariate case, namely, the asymptotic normality of the MLE. In the general case where this restriction is lifted, the MLE may find itself pinned along the boundary with strictly positive probability, giving rise to singularities in the distribution of the LRS Λ_n . Note in particular that in the univariate case, the main result of Chen & Hall [9] is thus non applicable to our purpose of testing for self-excitation as it breaks down under the Poisson null hypothesis $\varphi = 0$. The general statement for the convergence of the LRS however remains similar to the standard case, as still expresses in terms of quadratic forms in a Gaussian variable, see Van den Vaart [50][section 16.3].

3.2. Asymptotic of the likelihood ratio test. Let Θ_0 be the sub-model (17) associated to a null hypothesis of the type (2). Without loss of generality we may assume that the null adjacencies correspond to the first p parameter coordinates $\theta_1, \dots, \theta_p$. Then, $\sqrt{n}(\Theta - \{\theta_0\})$ converges¹ to $H_0 = \{0\}^p \times \mathbb{R}^{d-p}$ and $\sqrt{n}(\Theta - \{\theta_0\})$ to $H = (0, \infty)^p \times \mathbb{R}^{d-p}$. The asymptotic distribution of the LRS Λ_n is then given as the difference between the squared distances of a certain Gaussian variable to H_0 and H .

Theorem 1. *Under Assumptions 1 to 10, for any $\theta_0 \in \Theta_0$*

$$(19) \quad \Lambda_n \xrightarrow[n \rightarrow \infty]{\mathcal{L}(\mathbb{P}(\theta_0))} \|I(\theta_0)^{1/2} X - I(\theta_0)^{1/2} H_0\|_2^2 - \|I(\theta_0)^{1/2} X - I(\theta_0)^{1/2} H\|_2^2$$

where $X \sim \mathcal{N}(0, I(\theta_0)^{-1})$

¹In the sense of Painlevé-Kuratowski, see the introduction of Geyer [19].

In the standard case, H contains the whole parameter space and the second term vanishes. Here the term subsists as the singularities in the distribution of the MLE echo into that of the LRS. This gives rise to a $\bar{\chi}^2$ distribution.

Proposition 2. *Under Assumptions 1 to 10, the likelihood ratio for the set of hypotheses (10) and (11) is asymptotically distributed under any $\mathbb{P}(\theta_0)$ with $\theta_0 \in \Theta_0$ as a chi-bar law:*

$$(20) \quad \Lambda_n \xrightarrow[n \rightarrow \infty]{\mathcal{L}(\mathbb{P}(\theta_0))} \bar{\chi}^2(p)$$

The chi-bar distribution can be conceived as the one of a χ^2 law with random degrees of freedom depending on the number of zeros in the projection on H of its underlying Gaussian variable X in Theorem 1. The $\bar{\chi}^2$ thus expresses as a mixture of standard χ^2

$$\bar{\chi}^2(p) = \sum_{k=0}^p \omega_k \chi^2(k).$$

We refer to Annex A.2 for a rigorous derivation of the existence and expressions of the weights in our case, and to the works of Shapiro [43], Kudo [33], Self & Liang [42], and Susko [47] for further details on the links between the $\bar{\chi}^2$ and the standard χ^2 distributions. The weights (ω_k) have the unfortunate property of generally depending non-explicitly on the value of θ_0 . This problematic dependency can however easily be circumvented. In some specific albeit useful configurations – including the test of an univariate Poisson against the self exciting alternative – the weights are in fact known and independent from the true parameter. In the general case, Susko [47] suggests exploiting the relation between the effective degree of the chi-bar and its underlying normal variable X by approximating the conditional distribution of Λ_n by a $\chi^2(p-k)$ law when k zeros are estimated in the MLE $\hat{\theta}_n$. This yields the *conditional* chi-bar test. The p-values for our test are therefore given by the following rule:

- When the $(\omega_k)_{k \in \llbracket 1, p \rrbracket}$ are known, by Proposition 2, for any $\theta_0 \in \Theta_0$

$$\mathbb{P}_{\theta_0}(\Lambda_n \leq z_a^\omega) \xrightarrow[n \rightarrow \infty]{} a,$$

where z_a^ω is the quantile at level a for the $\sum_k \omega_k \chi^2(k)$ law .

- When the $(\omega_k)_{k \in \llbracket 1, p \rrbracket}$ are unknown, by Theorem 1 and Susko's Theorem [47], denoting \hat{p} the estimated number of zeros in $\hat{\theta}_n$, for any $k \in \llbracket 1, p \rrbracket$,

$$\mathbb{P}_{\theta_0}(\Lambda_n \leq z_a^{p-k} \mid \hat{p} = k) \xrightarrow[n \rightarrow \infty]{} a,$$

where z_a^{p-k} is the quantile at level a for the $\chi^2(p-k)$ law .

As the test statistic goes to ∞ as $n \rightarrow \infty$ under any $\mathbb{P}(\theta)$ with θ in the alternative, the test is consistent. As a by-product of Theorem 1, we recover the asymptotic distribution of the MLE.

Proposition 3 (Central limit theorem). *Under Assumptions 1 to 10, for any $\theta_0 \in \Theta_0$, the distribution of $\sqrt{n}(\hat{\theta}_n - \theta_0)$ converges under $\mathbb{P}(\theta_0)$ to the distribution of*

$$\arg \min_{Y \in H} \|I(\theta_0)^{1/2} X - I(\theta_0)^{1/2} Y\|_2^2$$

where $X \sim \mathcal{N}(0, I(\theta_0)^{-1})$. Furthermore, denoting by $\tilde{\theta}_n^0$ the non-null coordinates of the sparse MLE $\hat{\theta}_n^0$ and by $\tilde{\theta}_0$ the non-null coordinates of the true parameter, the distribution of $\sqrt{n}(\tilde{\theta}_n^0 - \tilde{\theta}_0)$ converges to a centered normal distribution with variance given by the inverse of the principal submatrix of $I(\theta_0)$ corresponding to the null hypothesis parameters.

4. NUMERICAL STUDY

In this section, we illustrate the preceding theoretical results with examples from simulated and real data. While the present work mainly deals with model selection for MHPs, allowing for a null adjacency opens interesting perspective for the univariate Hawkes process. Our first example is concerned with testing for self-excitation in such setting. This allows for an application on Ebay auctions, on which we test for potential deviations from a naive equilibrium. The second example is concerned with the bivariate Hawkes, which is of frequent use in financial applications. We show how our test may be used to differentiate between different price feedback regimes on intraday power markets. In these first two examples, the asymptotic distribution of the LRS is explicit and independent from the value of the true parameter. Finally, in our third and last example, we show how the test may be conducted in the general case, and demonstrate how Susko's [47] methodology applies in our setting.

4.1. The Poisson versus Self-exciting test. Self-excitation is the prominent feature of the univariate Hawkes process. A natural problem is to determine whether the observed event sequence does feature such a property at all. Recall that in spite of the univariate setting of this test, the results of Chen & Hall are non applicable under a Poisson null and we must have recourse to Theorem 1 instead. The Poisson *versus* Hawkes problem was first raised by Dachian & Kutoyants [13], whom propose a test in the long time asymptotic for the Poisson null hypothesis against the one sided alternative. It has since arisen again in the context of applications. In Kramer & Kiesel [31] for instance, the authors identify the $\bar{\chi}^2$ family of distribution as a the correct asymptotic for the likelihood ratio, but they are unable to conclude.

4.1.1. Setting and model. We first work in the setting of Dachian & Kutoyants [13], where the kernel $\varphi: \mathbb{R}^+ \mapsto \mathbb{R}^+$ of the Hawkes process depends on a single parameter $\alpha \in [0, 1)$

$$(21) \quad \varphi: t, \alpha \mapsto \alpha \Phi(t),$$

and is otherwise completely specified by a known positive function $\Phi: \mathbb{R}^+ \mapsto \mathbb{R}^+$. We also let the baseline fluctuate in time according to $\mu: t, \mu_0, \kappa \mapsto \mu_0 \exp(\kappa \frac{t}{T})$. Our question amounts to testing

$$(22) \quad \alpha = 0 \quad (\text{the process is a Poisson process}),$$

against the one sided alternative

$$(23) \quad \alpha > 0 \quad (\text{the process is a true Hawkes process}).$$

The asymptotic distribution under the null of Λ_n (22) is given by Theorem 1 as the one of $\mathbb{1}_{X>0} X^2$ where $X \sim \mathcal{N}(0, 1)$ (see annex A.2) and simplifies as

$$(24) \quad \bar{\chi}^2 = \frac{1}{2} \chi_0^2 + \frac{1}{2} \chi_1^2.$$

The test provides a straightforward decision rule as it rejects (22) at the significance level $a \in (0, 1/2)$ when Λ_n is above the $(1 - 2a)$ -quantile of the standard $\chi^2(1)$ distribution. The framework of Dachian & Kutoyants [13] is however too restrictive in practice. The function Φ usually depends on some additional decay parameter β , which is not known. We then face the issue that two different values for β yield the same distribution for the process under the null (22): a non identifiable nuisance parameter must be dealt with in addition to the boundary problem. The exact same configuration can be encountered in Ritz [40][Remark 4] for instance, where the author remarks that the distribution of the LRS then differs from (24).

A naive answer to the issue consists in falling back on the simpler case (21) by setting β at some pre-determined value β^* to conduct the test. The distribution of Λ_n under the null (22) is still given by (24) and the level of the test is therefore unchanged. The resulting mis-specification of β will however decrease the value of the LRS under the alternative, as underlined by Hansen [22]. In our case, we find numerical evidence that the subsequent power loss is very modest even for gross mis-specifications of β . In light of these results, we leave potential improvements on the naive test to further study (see for instance Andrews [4] for sup-likelihood ratio tests). Note also that the question of the test's robustness to errors in β goes beyond identifiability questions as the estimation of the decay parameter is subject to notoriously pervasive numerical instability issues.

4.1.2. *Simulation study.* We report the results for 10000 simulations over $T = 10$ s of the point process $N^{(n)}$ with intensity $n\mu$, corresponding to the sum of $n = 2000$ non-homogeneous Poisson process with intensity μ . The parameters are set to $\mu_0 = 2$ and $\kappa = 2$, and the simulation performed using a standard thinning approach. At each step we compute the value of the LRS Λ_n for the full model against the Poissonian one by numerically maximising the likelihood over the full ($\alpha > 0$) and sparse ($\alpha = 0$) parameter set corresponding to each hypothesis. Three sets of such simulations are performed, with the kernel of the process in the full model successively taken as a Pareto kernel, an exponential kernel, and a gamma kernel, all with decay $\beta = 10$.

$$(25) \quad \Phi: t, \beta \mapsto 1/\beta(1+t)^{-(1+\beta)} \quad \Phi: t, \beta \mapsto \beta \exp(-\beta t) \quad \text{and} \quad \Phi: t, \beta \mapsto \beta^2 t \exp(-\beta t).$$

We provide in figure 1 the QQ-plot for the empirical distribution of Λ_n conditionally on its being positive, which is expected to follow a $\chi^2(1)$ distribution. In practice, we use a threshold of $\varepsilon = 10^{-4}$ above which the LRS is considered positive.




FIGURE 1. QQ-plots for Λ_n conditional on positivity against its theoretical asymptotic distribution for $n = 2000$ trajectories. From left to right: Power-law, exponential and gamma kernels with decay $\beta = 10$.

Note that under our chosen parameter values, the simulated process averages about 20 jumps per realisation, which would prevent using standard estimation procedure based on long-time realisations of the process. The likelihood ratio behaves as expected under the null and we turn our attention to the distribution of the LRS under the alternative. Since this supposes the simulation of a large number of Hawkes processes, we limit our numerical study to the case of an exponential Hawkes process to keep the computation time under a reasonable length.

A collection of $N = 2000$ sets of n trajectories of a Hawkes process with constant baseline $\mu = 4$, decay $\beta = 3$, and kernel $\varphi: t \mapsto \alpha\beta \exp(-\beta t)$ is simulated for 12 different values of the weighted adjacency (or endogeneity) parameter α , and n successively set at $n = 500$ and $n = 2000$, totalling 60 million simulated trajectories. We compute in figure 2 the empirical power of the test at the 5% level for two specifications of the decay, one at the true value $\beta_0 = 3$, the other at $\beta = 30$ corresponding to a 900% misspecification.




FIGURE 2. Empirical power of the test for $N = 2000$ sets of n trajectories. Left: $n = 500$ trajectories. Right $n = 2000$ trajectories.

The numerical results suggest that a tenfold error in the choice of the decay incurs only a limited power loss, with most of the shift in the power function being contained around small local alternatives. In particular, the test remains consistent.

4.1.3. *Strategic bidding in online auctions.* We turn our attention to the real world setting of bid arrivals on online auction. The Ebay.com website runs second best price open auctions, meaning that the bids are public and the transaction settled at the second highest recorded bid. Were the bids sealed and static instead of public and dynamic, this would give rise to a Vickrey auction (see Krishna [32][p. 181]), where a Nash equilibrium is reached with participants bidding their full capacity. If Ebay bidders are insensitive to each other's behaviour, the observed equilibrium should resemble the one of the Vickrey auction. One would then expect bid arrivals to occur independently consistently with an homogeneous Poisson model. This is of course not what is observed in practice. Shmueli *et al.* [44] remark bid arrivals depend on complex factors, and use a three-stages non-homogeneous Poisson process to account for the changing nature of the process. Participants engage for instance in last minute bidding, a behaviour known as *sniping*. We propose an explanation for the deviation from the static equilibrium based on the categorisation of bidders into two cohorts:

- *Price sensitive* bidders, whose estimation is dynamic and depends on other bidder's behaviour.
- *Price insensitive* bidders, who place a single bid at full capacity, either at an entirely random time, or at the last few moments in an effort to squeeze out price-sensitive bidders.

We aim to prove the existence of the first category of agents by fitting a Hawkes model on bid arrival times. Ebay auctions participation is mostly human and we use a gamma kernel (25) to reflect the additional reaction time this incurs. We also retain the exponential baseline of model (21) in an effort to match the increasing intensity pattern. If bidders were to all belong to the first category, the increasing Poisson-like exponential intensity should account entirely for the sniping phenomenon. If it results instead from agents competitively interacting, the Hawkes model is more appropriate. Our null and alternative hypotheses are therefore the same as for the preceding simulation study, that is a Poisson null $\alpha = 0$ against the one-sided positive alternative. Should the test reject the null hypothesis, Ebay auctions would fall in the framework of interdependent values. The dynamic nature of equilibrium bidding strategies is well documented in such models (Krishna [32][Proposition 6.2 p. 91]), consistently with our proposed interpretation.

The data consists in bid arrival times of 628 Ebay auctions opened and settled in 2020 for luxury watches, digital assistance devices and game consoles. Auctions may run over 7, 5 or 3 days from which we keep only the last 2.8 days. This allows for a simple synchronous model, and avoids the non stationarity due to novelty-induced bids at the start of 3-days auctions, which are of little relevance to our question.

We set the parameter space for the decay as the closed real interval from 1 to 10000 days⁻¹ and repeat the test with the two values $(\beta_1, \beta_2) = (5, 1000)$ to reduce the potential misspecification. Using the Bonferroni correction

$$(26) \quad \mathbb{P} \left[\bigcup_{i=1}^m \{ \Lambda_n(\beta_i) \geq q_{1-\frac{a}{m}} \} \right] \leq \sum_{i=1}^m \mathbb{P} [\Lambda_n(\beta_i) \geq q_{1-\frac{a}{m}}] \leq m \frac{a}{m} = a,$$

we obtain a level a test from the repeated procedure by rejecting the null as soon as one of the LRS is above the quantile of order $1 - a/m$ of the χ_1^2 distribution. When the value set for β is large, the repetition allows to recoup some of the lost power as long as m is kept reasonably small (here $m=2$). We report in table 1 the LRS and p-value for each β_i , with $\beta_0 = 0$ representing the Poisson null.

β	\hat{m}	$\hat{\kappa}$	$\hat{\alpha}$	LRS	p
0	4.95	3.43	N.A	N.A	N.A
5	0.24	3.7	0.96	2.26e2	7e - 51
1000	0.31	1.50	0.89	4.88e3	~ 0

TABLE 1. Estimated parameters and likelihood ratio statistic against the Poisson null of a gamma Hawkes process on bid arrival times of 627 ebay auctions.

Both decay choices yield p-values far below 0.005, the Poisson model is rejected at the level 0.01, matching our supposition of the price being formed through competitive price discovery. Though the second-price mechanism of Ebay auctions does not reward late bidding, their dynamic nature favors interdependent strategies, in which bidders await and react to the information flow from competing bids. Providing a complete model for the dynamic of the auction would require a more flexible baseline, mimicking the varying intensity of Shmueli *et al.* [44]. Such ambition is out of the scope of this exploratory example which is intended to showcase the general relevance of our inference procedure in the context of dynamic auctions. In doing so, we have uncovered a

feedback mechanism between bidders' valuations and bid values. Finally, we acknowledge our conclusions are perhaps dataset-specific. The auctions set we estimated the model upon suffers from a high prevalence of luxury goods, the demand for which features a characteristic positive price elasticity. Accessing additional data would here allow for more general conclusions.

4.2. Testing for cross-excitation in the bivariate Hawkes process. Suppose one wishes to test for the joint significance of two (weighted) adjacency coefficients α_{ij} and α_{kl} belonging to different coordinates k and l of a multivariate Hawkes process. From the expression of the Fisher information (15), that $k \neq l$ entails the asymptotic independence of the $\hat{\alpha}_{ij}$ and $\hat{\alpha}_{kl}$. One may then derive the limit distribution of the likelihood ratio as

$$(27) \quad \bar{\chi}^2 = \frac{1}{4}\chi_0^2 + \frac{1}{2}\chi_1^2 + \frac{1}{4}\chi_2^2$$

by remarking the weights of the mixture express either in terms of the probability of two independent Gaussian variables being positive or of a single Gaussian being positive. Alternatively, setting the antidiagonal term to 0 in example 7 of Self & Liang [42] yields the same result. While somewhat specific, the present example has a practical use in the context of financial statistics, and we shape our simulations and application around the microstructure model of Bacry *et al.* The price of an asset (P_t) over a timeframe $t \in [0, T]$ is modelled as an accumulation of positive and negative jumps across the discrete price grid, respectively corresponding to pure jump processes (N_t^+) and (N_t^-) with null initial conditions:

$$P_t - P_0 = N_t^+ - N_t^-.$$

For liquid enough assets, the processes are simple as price jumps rarely exceed the mesh size of the grid (known as the *tick size*). In the converse, the (N_t^\pm) are marked with price increment sizes. The two jump processes are chosen as the coordinates of a bivariate Hawkes process with baseline μ and kernel φ . This model, originally intended for high frequency prices at the level of the tick grid, has also become an instrument of choice for power prices, which retain at large time scales some features reminiscent of microstructure phenomena – see Deschatre & Gruet [16] for a price model and see Kramer & Kiesel [31] for order book dynamics. From now on we suppose the baseline and kernel take the shapes

$$(28) \quad \mu = \exp(\kappa t) [m \quad m]^T \quad \varphi: t, x \mapsto \beta x \begin{bmatrix} \gamma_1 & \alpha \\ \alpha & \gamma_2 \end{bmatrix} \exp(-\beta t).$$

In the classical setting of Bacry *et al.*, the price trajectories are guided purely by reversion effects, corresponding to the $\gamma_1 = \gamma_2 = 0$ regime. One may wonder whether this kernel shape, which was introduced to account for the negative auto-correlation of price increments at high frequency remains reasonable at larger scale. This corresponds to testing the following hypotheses:

$$(29) \quad \begin{array}{ll} \mathcal{H}_0 : \gamma_1 = 0, \gamma_2 = 0 & \text{(Prices are purely reverting),} \\ \mathcal{H}_1 : \gamma_1 > 0, \gamma_2 > 0 & \text{(Prices have a momentum component),} \end{array}$$

where the term *momentum* is borrowed from the financial vernacular to designate a self-sustained directional dynamic in prices.

4.2.1. Simulation study. The simulated process has the specification (28) with $m = 5$, $\alpha = 5$, $\kappa = 2\text{Hz}$, $\beta = 10\text{Hz}$ and lives under the null $\gamma_1 = \gamma_2 = 0$. The distribution of the marks is the one of $|X| + \delta$ where $X \sim \mathcal{N}(0, 1)$ and $\delta = 10^{-2}$ is an offset parameter meant to prevent numerical error in the simulation. We report the results of repeated estimations for a set of 10000

simulations over a timeframe of $T = 1s$ and $n = 800$. Figure 3 displays the QQ-plot for the values obtained for Λ_n , conditionally on its being strictly positive, with a non-nullity threshold of $\varepsilon = 10^{-5}$.




FIGURE 3. QQ-plots for Λ_n conditionally on its positivity for $N = 10000$ simulations of $n = 800$ trajectories of a marked MHP with characteristics (28) and absolute normal marks.

Conditional on positivity, a chi-bar variable with weights (27) has a $2/3\chi^2(1) + 1/3\chi^2(2)$ law, and one observes $\max(\Lambda_n, \varepsilon) \simeq \max(\Lambda_n, 0)$ to converge well enough to such a distribution .

4.2.2. Single asset price dynamics of late night delivery power futures. We proceed to our application to intraday power prices. The dataset consists of 4-hours long windows from 5 to 1 hour to delivery of hourly power future mid-prices² for the $n = 364$ trading sessions occurring from deliveries from January 1, 2019 to December 30th, 2019. The exclusion of the last hour of trading serves so as to avoid taking into account effects associated with the delivery zone reducing in size after that point. In Deschatre & Gruet [16], the authors focus on the most liquid of the hourly products of the German intraday power market. Thanks to the aggregation procedure (6), we are also able to estimate the Hawkes-based power prices model on late night and early morning deliveries future contracts, which are less intensively traded. The retained futures cover the 12 consecutive hourly delivery periods beginning from 18h to 5h.

The null hypothesis is rejected with high certainty for each future contract, indicating the existence of a statistically significant momentum component in the dynamics of German intraday power prices. We hypothesise this stems from market participants competing to capture scarce liquidity. The cascading patterns observed in mid-prices hint at liquidity reconstituting slower than it is consumed, which would be consistent with our interpretation. Using a stochastic baseline Hawkes process, Kramer and Kiesel [31] find an influence of wind and solar forecast errors on exogenous order arrivals. Marking the kernel for similar exogenous factors could allow for testing potential relations between common stresses on power traders and episodes of directional variations in price.

Remarkably, our estimations situate the price endogeneity η around the low value of 0.45 which is consistent with the findings of Deschatre & Gruet [16], meaning that price moves do not generate as many reactions as one would expect. This would put the intraday power markets at

²Arithmetic average of best bid and best ask prices.

Delivery start	18h	19h	20h	21h	22h	23h	0h	1h	2h	3h	4h	5h
$\alpha^\#$	2.30	2.54	2.11	2.21	2.40	2.11	2.30	3.04	2.90	2.44	2.20	2.35
γ_1	2.03	2.26	2.41	2.25	2.77	1.37	2.03	2.41	2.26	2.71	2.33	2.35
γ_2	1.62	2.49	2.30	2.01	2.36	1.34	1.62	2.27	2.20	2.35	1.69	1.52
$\beta^\#$	930	516	559	510	403	369	930	460	350	556	488	626
$\mu_0^\#$	68.2	95.0	104	99.3	98.1	67.6	68.2	60.9	66.6	51.1	46.7	93.0
$\kappa^\#$	1.14	0.74	0.76	0.74	0.60	0.54	1.14	0.89	0.91	1.11	1.21	0.56
α^b	3.86	4.40	4.44	4.19	4.89	3.27	3.86	5.38	4.99	4.70	4.02	3.99
β^b	692	385	371	467	302	327	692	296	253	444	470	515
μ_0^b	70.2	61.9	105	101	98.9	67.6	70.2	47.6	62.3	69.3	77.1	96.8
κ^b	1.16	1.22	0.77	0.75	0.61	0.58	1.16	0.88	0.55	0.93	0.89	0.58
$\mathbb{E}[X]$	0.10	0.10	0.10	0.1	0.11	0.11	0.10	0.09	0.10	0.1	0.11	0.11
$\eta^\#$	0.41	0.47	0.44	0.46	0.55	0.46	0.41	0.54	0.51	0.50	0.42	0.34
η^b	0.39	0.48	0.45	0.41	0.54	0.48	0.39	0.54	0.50	0.47	0.43	0.43
LRS	647	536	371	510	644	280	647	344	249	718	954	1379
p	~ 0											

TABLE 2. Estimates for model (28) as applied to German intraday hourly power futures mid-prices, 5 to 1h to delivery on the first 364 trading sessions of 2019. The $\#$ and b superscripts respectively denote the full and sparse models.

odds with more conventional asset classes, were values close to criticality $\eta = 1$ are repeatedly observed (see for instance Hardiman *et al.* [24] [25]), reflecting a no arbitrage condition (see Juselin & Rosenbaum [30]). One could assess the robustness of this conclusion and its potential reasons by repeating the preceding analysis with different choices for the marks, as precedently suggested. The low endogeneity would be compatible with the conjecture of liquidity constraints being an important price driver, since it suggests some sense of "missing participants". The particularities of power trading (which is typically a capital-intensive activity and incurs specific risks) may indeed be repulsive to some of the conventional liquidity providers. Depending on the role of such participants, the momentum component could vanish on more accessible markets. In all generality, one would expect the additional barriers to entry to limit the diversity of market participants and favour the appearance of significant non-reverting patterns in price dynamics as a result of the alignment of traders' incentives and constraints.

4.3. Testing for an arbitrary set of adjacency coefficients.

4.3.1. *Setting and procedure.* Apart from the simple configurations above, the weights are usually intractably tied to the value of the true parameter. As exposed with Proposition 2, a conditional test can then be conducted. Recall that, from Susko's theorem [47], conditionally on k zeros being observed in the MLE, the LRS Λ_n closely follows a $\chi^2(p - k)$ distribution. The resulting procedure is the following: if Λ_n is null, the test never rejects \mathcal{H}_0 . If Λ_n is positive, the full and sparse MLE should differ and a number $k < p$ of null adjacency coefficients are observed in $\hat{\theta}_n$. The test then rejects \mathcal{H}_0 when Λ_n is above the quantile of chosen level for the $\chi^2(p - k)$ distribution.

4.3.2. *Simulation study.* We apply the general procedure to a 10-dimensional Hawkes processes. The exponential kernel was again retained to keep an acceptable computation time. Under the null, the process is a juxtaposition of two 5-dimensional Hawkes process with cyclic interaction

patterns, each parameterised by a single adjacency coefficient α_i . Under the alternative, four coordinate pairs interact, binding the two cycles through four weights $\gamma_1 \cdots \gamma_4$. The precise graph of interactions is encoded into Figure 4 below.




FIGURE 4. Interaction graph of the example MHP model. Full edges: interactions under the null. Dotted edges: Additional interactions under the alternative.

The baseline and decay are constant and set to the same values $\mu = 8, \beta = 10$ for all 10 coordinates. In figure 5 are the QQ-plots for the distribution of Λ_n conditionally on the number of zeros being observed among the MLEs, with a threshold at $\varepsilon = 10^{-5}$. We observe the conditional distributions of the LRS converge well to the respective asymptotic counterparts they are given by Susko's theorem. An occasional degree mis-attribution seemingly persists, resulting in a more pronounced deviation of the distribution conditional on the effective degree of the LRS being 1.




FIGURE 5. Conditional QQ-plots for the empirical distribution of Λ_n against their theoretical $\chi^2(p - k)$ asymptotic distribution. 60000 simulations, $n = 1000$.

4.3.3. *Multi-asset price dynamics of late night delivery power futures.* We continue with German intraday power futures mid prices. Our aim here is to retrieve the temporal dependency structure between multiple hourly futures. The dataset is the one of example 4.2.2. We consider the four hourly futures with deliveries starting at each hour from 22 to 1h on the 364 trading sessions of 2019. From each session is kept the 4 hour time window from 21h to midnight. The model of Deschatre & Gruet [16] is extended to incorporate inter-asset excitation in addition to reversion and momentum factors. This results in the kernel shape

$$\alpha = (\alpha_{kl}) = \begin{bmatrix} a_1 & a_2 & 0 & 0 & 0 & 0 & 0 & 0 \\ a_2 & a_1 & 0 & 0 & 0 & 0 & 0 & 0 \\ \gamma_1 & 0 & a_1 & a_2 & 0 & 0 & 0 & 0 \\ 0 & \gamma_1 & a_2 & a_1 & 0 & 0 & 0 & 0 \\ \gamma_4 & 0 & \gamma_2 & 0 & a_1 & a_2 & 0 & 0 \\ 0 & \gamma_4 & 0 & \gamma_2 & a_2 & a_1 & 0 & 0 \\ \gamma_6 & 0 & \gamma_5 & 0 & \gamma_3 & 0 & a_1 & a_2 \\ 0 & \gamma_6 & 0 & \gamma_5 & 0 & \gamma_3 & a_2 & a_1 \end{bmatrix}$$

where the simple cross-dependence structure is adapted from Bacry *et al* [6][section 5.2]. The model forces a lead-lag relation wherein the less liquid late night futures react to price moves in the more liquid evening deliveries. This rules out temporal inconsistencies in which market factors materialising past the delivery period of a future may move its price by a ricocheting along the edges of the causal graph of (N_t) . We have also taken advantage of the closeness of momentum and reversion factors estimates of example 4.2.2, keeping only two parameters for the a_i s. Likewise, we take only one decay parameter β for the whole kernel. More precisely, our kernel has the shape

$$\varphi_{kl} : t, x \mapsto \alpha_{kl} x \exp(-\beta t).$$

A specificity of our setting is the asynchronicity of trading windows for each future. Unrestricted trading of the asset expiring at hour $n - 1$ halts one hour before it does for the future expiring at hour n , which is a substantial fraction of the estimation window. Our marked Hawkes process lives on a multidimensional triangular time-frame $[0, T_1] \times \dots \times [0, T_4]$, which only incurs an innocuous modification of the log-likelihood and does not alter our results. Accordingly, the baseline is parameterised by two 4-dimensional parameters (m_i) and (κ_i) and by the known end times (T_i) :

$$\mu : t \mapsto \left[m_1 e^{\kappa_1 \frac{t}{T_1}} \quad m_1 e^{\kappa_1 \frac{t}{T_1}} \quad m_2 e^{\kappa_2 \frac{t}{T_2}} \quad m_2 e^{\kappa_2 \frac{t}{T_2}} \quad m_3 e^{\kappa_3 \frac{t}{T_3}} \quad m_3 e^{\kappa_3 \frac{t}{T_3}} \quad m_4 e^{\kappa_4 \frac{t}{T_4}} \quad m_4 e^{\kappa_4 \frac{t}{T_4}} \right]^T.$$

Transaction costs may incentivize a reduction in the number of simultaneously traded assets. Such efficiency gains should come naturally in a MHP-based approach as the model isolates direct interactions. We record in table 3 the results of our estimations for adjacency parameters.

Model	\hat{a}_1	\hat{a}_2	$\hat{\gamma}_1$	$\hat{\gamma}_2$	$\hat{\gamma}_3$	$\hat{\gamma}_4$	$\hat{\gamma}_5$	$\hat{\gamma}_6$	LRS
every $\gamma_j > 0$	4.26	4.75	1.34	0.93	0.95	0.29	0.26	0.13	N.A
$\alpha_{ij} = 0$ when $i > j + 1$	3.9	5.12	1.24	0.95	1.54	N.A	N.A	N.A	2361
$\alpha_{ij} = 0$ when $i > j + 2$	4.2	4.89	1.19	1.22	1.47	0.29	0.9	N.A	1569

TABLE 3. Inferred adjacency parameters for the first 364 trading sessions of 2019 on the German intraday power market.

A first test is run with null hypothesis that direct dependence between futures vanishes beyond a 2 hours delay, that is $\gamma_4 = \gamma_5 = \gamma_6 = 0$ against the alternative that they are all positive. We use a threshold of $\varepsilon = 10^{-5}$ under which estimates are considered null. Since no estimates fall below ε , the p-value for the test is obtained by comparing the LRS with the quantile of appropriate level of a $\chi^2(3)$ distribution. As the null is rejected with high certainty, a second test is run with the null hypothesis that $\gamma_6 = 0$ against the one-sided alternative. The null is again rejected at the 0.05 confidence level. Despite these unexpected results it is compelling to note the amplitude of interactions between two futures i, j decreases fast as $T_i - T_j$ increases. In practice, the transactions costs of the intraday power market will often prevent executing on such weak signal.

4.4. Comparison to other methods. Our method is natural in that it leverages the specific property of linear Hawkes processes of remark 2 into a useful Poisson approximation of the Hawkes process. When T is large enough to allow for credible trajectory-by-trajectory estimations, different procedures are usually preferred. In Bonnet *et al.* [7], the authors estimate the MLE on each trajectory and average the resulting sequence. In Deschatre & Gruet [16], the likelihood for the full dataset (or the sum of each log-likelihood) is maximised. One may wonder whether the two more conventional approaches keep producing acceptable result as T decreases and how they compare to our method. We apply the three options to an univariate Hawkes model with constant baseline and an exponential kernel $t \mapsto \alpha\beta \exp -\beta t$. In figure 6 are the empirical distribution of the estimates for α according to each method, the ground truth being an homogeneous Poisson process $\alpha = 0$ with intensity $\mu = 5\text{Hz}$. The simulations are performed on a horizon of $T = 10\text{s}$. Note that this creates about the same number as in a trajectory of the real world dataset of example 4.1.




FIGURE 6. Histograms for the estimators of α for each method, obtained for $N = 4000$ simulations of $n = 500$ trajectories of a Poisson process with intensity 5Hz over a timeframe of $T = 10\text{s}$. Left: our method (summing the n trajectories). Top-right: maximising the sum of the n log-likelihoods. Bottom-right: averaging the MLEs over the n trajectories.

When applied to our setting, the two conventional approaches produce positive biases, which is especially detrimental to our main objective of testing for self-excitation. Consequently, we do not investigate further their properties at small time scales. Though the present example tells nothing of large scale properties, it is however interesting to observe our method produces the largest variance and might become a subpar solution as T increases and standard estimators converge again. Note also that in the case of the average of MLES strategy, the bias is at least partly due to the positivity restrictions imposed upon the weighted adjacency α , and could therefore disappear entirely in a more flexible model *à la* Bonnet *et al.* [7] where α can reach negative values.

5. PREPARATION FOR THE PROOFS

Throughout the proofs, θ_0 is a parameter in Θ_0 , and $(N_t^{(n)})$ a point process with intensity $(\lambda_t^{(n)}(\theta_0))$. Unless specified otherwise, any expectation \mathbb{E} is taken under $\mathbb{P}(\theta_0)$. We will be concerned with the asymptotic behaviour of a variety of derivatives of the log-likelihood, which we introduce hereafter. Define the score $\mathcal{S}(n, \cdot) = (\mathcal{S}(n, \cdot)_i) = (\partial_{\theta_i} \mathcal{L}(n, \theta))$ where for any $\theta \in \Theta$ and $i \in \llbracket 1, d \rrbracket$,

$$(30) \quad \mathcal{S}(n, \theta)_i = \sum_{k=1}^K \int_0^T \frac{\partial_{\theta_i} \lambda_{k,s}^{(n)}(\theta)}{\lambda_{k,s}^{(n)}(\theta)} dN_k^{(n)} - \sum_{k=1}^K \int_0^T \partial_{\theta_i} \lambda_{k,s}^{(n)}(\theta) ds.$$

Similarly, $\partial_{\theta} \mathcal{S}(n, \cdot) = (\partial_{\theta} \mathcal{S}(n, \cdot)_{ij})$, where for any $i, j \in \llbracket 1, d \rrbracket$

$$(31) \quad \begin{aligned} \partial_{\theta} \mathcal{S}(n, \theta)_{ij} &= \sum_{k=1}^K \int_0^T \frac{\partial_{\theta_i} \partial_{\theta_j} \lambda_{k,s}^{(n)}(\theta)}{\lambda_{k,s}^{(n)}(\theta)} - \frac{\partial_{\theta_i} \lambda_{k,s}^{(n)}(\theta) \partial_{\theta_j} \lambda_{k,s}^{(n)}(\theta)}{(\lambda_{k,s}^{(n)}(\theta))^2} dN_k^{(n)} \\ &\quad - \sum_{k=1}^K \int_0^T \partial_{\theta_i} \partial_{\theta_j} \lambda_{k,s}^{(n)}(\theta) ds. \end{aligned}$$

Finally, define the empirical information $\mathcal{J}(n, \cdot) = (\mathcal{J}_{ij}(n, \cdot))_{ij}$ as

$$(32) \quad \mathcal{J}_{ij}(n, \theta) = \sum_{k=1}^K \int_0^T \frac{\partial_{\theta_i} \lambda_{k,s}^{(n)}(\theta)}{\lambda_{k,s}^{(n)}(\theta)} \frac{\partial_{\theta_j} \lambda_{k,s}^{(n)}(\theta)}{\lambda_{k,s}^{(n)}(\theta)} dN_{k,s}^{(n)}.$$

We first recall some well-known results, which make a recurrent appearance in our proofs.

Lemma 2. *Let $(X_n(\nu))_{n,\nu}$ be a family of random variables with values in some normed space $(E, \|\cdot\|)$, indexed by $n \in \mathbb{N}$ and ν in some compact $A \subset \mathbb{R}^d$. If for any $\nu \in A$,*

$$(33) \quad X_n(\nu) = O_{\mathbb{P}}(1) \quad (\text{respectively: } o_{\mathbb{P}}(1)),$$

and $(X_n(\cdot))$ is stochastically equicontinuous: for any $\eta > 0$ and $\varepsilon > 0$, there is $\delta > 0$ such that

$$(34) \quad \limsup_{n \rightarrow \infty} \mathbb{P} \left[\sup_{\|\nu_1 - \nu_2\| < \delta} \|X(n, \nu_1) - X(n, \nu_2)\| > \eta \right] \leq \varepsilon,$$

then (33) holds uniformly over any compact $\Gamma \subset E$, that is

$$\sup_{\nu \in \Gamma} \|X_n(\nu)\| = O_{\mathbb{P}}(1) \quad (\text{respectively: } o_{\mathbb{P}}(1)).$$

In the sequel, the conditions for the supremum of random variables to remain measurable will always be met, see appendix C of Pollard [37] for more details. Lemma 2 is commonly found in the context of parametric inference, and a similar argument appears in concise form in Chen & Hall [9]. We recall its proof for the sake of completeness.

Proof of Lemma 2. Let $(X_n(\nu))$ satisfy the conditions of Lemma 2, Γ be a compact set, and let $\varepsilon > 0$. For any $\delta > 0$, we have a finite covering of Γ with open sets included in balls with centres (c_i) , $i \in \llbracket 1, P_\delta \rrbracket$, and radius strictly below δ . For every $\nu \in \Gamma$, write $c(\nu)$ for the ball center closest to ν .

$$(35) \quad \begin{aligned} & \mathbb{P} \left[\sup_{\nu \in \Gamma} \|X_n(\nu)\| > M \right] \\ & \leq \mathbb{P} \left[\sup_{\nu \in \Gamma} \|X_n(\nu) - X_n(c(\nu))\| > M/2 \right] + \mathbb{P} \left[\sup_{\nu \in \Gamma} \|X_n(c(\nu))\| > M/2 \right]. \end{aligned}$$

Now, under condition (34), for any sufficiently small δ , the first term on the right hand side of (35) vanishes as $n \rightarrow \infty$. The second term is bounded by

$$\sum_{i=1}^{P_\delta} \mathbb{P}(\|X_n(c_i)\| > M).$$

When $X_n(\cdot)$ is pointwise $O_{\mathbb{P}}(1)$ over Γ (respectively: pointwise $o_{\mathbb{P}}(1)$), this bound may be made arbitrarily small by choosing M sufficiently large (respectively: uniformly in M), yielding the desired results. \square

Stochastic equi-continuity results will most often proceed directly from the regularity of the kernel φ . In some other instances, we have recourse to a chaining Lemma.

Lemma 3 (Chaining, Pollard [37], Section VII.2, p. 142-146). *Let (Z_x) be a stochastic process indexed by x in a compact metric space (E, d) with values in some normed space $(F, |\cdot|)$. If there exists a strictly positive constant D such that for any $x, y \in E$, and any $\eta > 0$,*

$$(36) \quad \mathbb{P} \left[|Z(x) - Z(y)| \geq \eta d(x, y) \right] \leq 2 \exp \left[- \frac{1}{2} \frac{\eta^2}{D^2} \right],$$

then for any $0 < \varepsilon < 1$,

$$(37) \quad \mathbb{P} \left[\text{There is some } x, y \in E \text{ such that } |Z(x) - Z(y)| \geq 26DJ(d(x, y)) \text{ and } d(x, y) < \varepsilon \right] \leq 2\varepsilon,$$

where J is the covering integral

$$J(\delta) = \int_0^\delta \left(2 \ln \left[\frac{N(u)^2}{u} \right] \right)^{\frac{1}{2}} du,$$

in which $N(u)$ is the covering number, that is the size of the smallest u -net on E for the metric d .

Observe as in Pollard [37] that, in practice, one may replace $J(\cdot)$ in (37) by any finite upper bound on the covering integral, and the exponential bound in (36) by any bound decreasing in η at the same speed. The chaining lemma requires concentration inequalities, which will arise from Lemma 4 below.

Lemma 4 (Lemma 2.1 of Van de Geer [49]). *Let (M_t) be a martingale over $[0, T]$ with predictable variation $V_t = \langle M, M \rangle_t$ and such that $|\Delta M_t| = |M_t - M_{t-}| < \kappa$ for all $t \in [0, T]$. Then, for any $a > 0$ and $b \in \mathbb{R}$,*

$$\mathbb{P} \left[\text{There is some } t \in [0, T] \text{ such that } |M_t| > a \text{ and } V_t \leq b^2 \right] \leq 2 \exp \left[- \frac{1}{2} \frac{a^2}{a\kappa + b^2} \right].$$

Some more specific devices now come into play. The definition of the intensity 1 gives rise to a variety of Volterra integral equations. We recall some elementary results regarding their resolvents.

Lemma 5 (On Volterra equations). *Let $\kappa: \mathbb{R}^+ \mapsto \mathcal{M}_K(\mathbb{R}^+)$ be a locally bounded function such that*

$$\rho\left(\int_0^\infty \kappa(s) ds\right) < 1,$$

and m a continuous function from \mathbb{R}^+ to \mathbb{R}^K . Define $\mathcal{K} = \sum_{k=1}^\infty \kappa^{\star k}$, wherein the $\star k$ -superscript denotes the k -fold convolution. Then, there exists a unique solution in locally bounded functions from \mathbb{R}^+ to \mathbb{R}^K to the integral equation in g

$$(38) \quad g(t) = m(t) + \int_0^t \kappa(t-s)g(s) ds, \quad t \in \mathbb{R}^+,$$

given by

$$g: t \mapsto m(t) + \int_0^t \mathcal{K}(t-s)m(s) ds.$$

See for instance Bacry *et al.* [6][Lemma 3 p. 16] or Gripenberg [21][Theorem 3.5 p. 44] for a proof. Lemma 1 is then an immediate consequence of Lemma 5. Before getting to its short proof, we extend its assertion into the more general form of Corollary 1 which allows for values of θ different to the true parameter.

Corollary 1 (An extension of Lemma 1). *Under assumption 9, for any $t \in [0, T]$, $n \in \mathbb{N}^*$ and $\theta \in \Theta$,*

$$\mathbb{E}\left[\frac{1}{n}\lambda_t^{(n)}(\theta)\right] = \mu(t, \theta) + \int_0^t \langle F, \varphi \rangle(t-s, \theta)h(s, \theta_0) ds,$$

where $s \mapsto h(s, \theta_0)$ is the unique solution in χ to the Volterra equation $\chi = \mu(\cdot, \theta_0) + \langle F, \varphi \rangle(\cdot, \theta_0) \star \chi$.

Proof. Let $t \in [0, T]$ and $\theta \in \Theta$. Under assumptions 4 and 9, The process

$$\left(\int_{[0, s] \times \mathbb{R}^m} \mathbb{1}_{u < t} \varphi(t-u, x, \theta_0) \mathbf{M}^{(n)}(du, dx)\right), \quad s \in [0, T],$$

is a martingale. In particular, at $s = t$,

$$(39) \quad \mathbb{E}\left[\int_{[0, t] \times \mathbb{R}^m} \varphi(t-s, x, \theta) \mathbf{N}^{(n)}(ds, dx)\right] = \int_0^t \langle F, \varphi \rangle(t-s, \theta) \mathbb{E}\left[\frac{1}{n}\lambda_s^{(n)}(\theta_0)\right] ds.$$

Applying (39) to the intensity (1), one gets

$$(40) \quad \mathbb{E}\left[\frac{1}{n}\lambda_t^{(n)}(\theta)\right] = \mu(t, \theta_0) + \int_0^t \langle F, \varphi \rangle(t-s, \theta) \mathbb{E}\left[\frac{1}{n}\lambda_s^{(n)}(\theta_0)\right] ds.$$

Setting $\theta = \theta_0$ shows the normalised intensity at the true parameter satisfies a Volterra integral equation with baseline $\mu(\cdot, \theta_0)$ and kernel $\langle F, \varphi \rangle(\cdot, \theta_0)$. By virtue of Lemma 5, we have Lemma 1. Inserting the resulting expression for $\lambda_s^{(n)}(\theta_0)$ into the last term of (40) yields corollary 1. \square

Following corollary 1, define

$$\bar{\lambda}(t, \theta, \theta_0) = (\bar{\lambda}_k(t, \theta, \theta_0))_k = \left(\mu_k(t, \theta) + \int_0^t \sum_{l=1}^K \langle F, \varphi \rangle_{kl}(t-s, \theta) h_l(s, \theta_0) ds\right)_k.$$

With the same differential notation $\partial_\theta^{\otimes 1} u = (\partial_{\theta_i} u)_i$ and $\partial_\theta^{\otimes 2} u = (\partial_{\theta_i} \partial_{\theta_j} u)_{ij}$, as Chen & Hall [9], define for any $p \in \{1, 2\}$

$$\partial_\theta^{\otimes p} \bar{\lambda}(t, \theta, \theta_0) = (\partial_\theta^{\otimes p} \bar{\lambda}_k(t, \theta, \theta_0)) = \left(\partial_\theta^{\otimes p} \mu_k(t, \theta) + \int_0^t \sum_{l=1}^K \partial_\theta^{\otimes p} \langle F, \varphi \rangle_{kl}(t-s, \theta) h_l(s, \theta_0) ds\right)_k.$$

Note that $\bar{\lambda}$ does not depend on n and that $\bar{\lambda}(\cdot, \theta_0, \theta_0)$ is simply $h(\cdot, \theta_0)$. As mentioned in the introduction, we will be concerned with obtaining a law of large number for the intensity. The deviation of $n^{-1}\lambda^{(n)}$ from $\bar{\lambda}$ need be further precised. To this end, we extend a result of Jaisson & Rosenbaum [29][proposition 2.1 p. 7] (see also Takeuchi [48][Proposition 1 p. 230]).

Lemma 6. *Under Assumption 9, for any $\theta \in \Theta$ and $t \in [0, T]$,*

$$(41) \quad \frac{1}{n}\lambda^{(n)}(t, \theta) = \bar{\lambda}(t, \theta, \theta_0) + \int_{[0, t] \times \mathbb{R}^m} \Psi(t-s, x, \theta_0, \theta) \frac{M^{(n)}}{n}(ds, dx)$$

where Ψ is defined by

$$\Psi = \sum_{k \geq 1} \tilde{\varphi}_k \quad \text{with} \quad \tilde{\varphi}_k(s, x, \theta_0, \theta) = \langle F, \varphi \rangle^{*(k-1)}(\cdot, \theta_0) \star \varphi(\cdot, x, \theta)(s).$$

Remark 10. *The function Ψ defined in Lemma 6 verifies for any $\theta \in \Theta$*

$$\langle F, \Psi \rangle(\cdot, \theta_0, \theta) = \int_{\mathbb{R}^m} \Psi(\cdot, x, \theta_0, \theta) F(dx) = \langle F, \varphi \rangle(\cdot, \theta) + \langle F, \varphi \rangle(\cdot, \theta) \star \sum_{k=1}^{\infty} \langle F, \varphi \rangle^{*k}(\cdot, \theta_0).$$

At $\theta = \theta_0$, this yields $\langle F, \Psi \rangle(\cdot, \theta_0) = \sum_{k=1}^{\infty} \langle F, \varphi \rangle^{*k}(\cdot, \theta_0)$. The notation of Lemma 6 is thus consistent with that of Lemma 1 and Corollary 1, which it implies.

Proof of Lemma 6. Let $\theta \in \Theta$ and $n \in \mathbb{N}^*$. One has for any $t \in [0, T]$

$$(42) \quad \frac{1}{n}\lambda_t^{(n)}(\theta) = \mu(t, \theta_0) + \int_{[0, t] \times \mathbb{R}^m} \varphi(t-s, x, \theta) \frac{1}{n} M^{(n)}(ds, dx) + \int_0^t \langle F, \varphi \rangle(t-s, \theta) \frac{1}{n} \lambda_s^{(n)}(\theta_0) ds.$$

At $\theta = \theta_0$ in particular, $n^{-1}\lambda_t^{(n)}(\theta_0)$ verifies a Volterra equation with kernel $\langle F, \varphi \rangle(\cdot, \theta_0)$. Applying Lemma 5, using Fubini's theorem, and re-arranging the terms, for any $t \in [0, T]$,

$$(43) \quad \begin{aligned} \frac{1}{n}\lambda_t^{(n)}(\theta_0) &= \mu(t, \theta_0) + \int_0^t \langle F, \Psi \rangle(t-s, \theta_0, \theta) \mu(s, \theta_0) ds + \int_{[0, t] \times \mathbb{R}^m} \varphi(t-s, x, \theta_0) \frac{1}{n} M^{(n)}(ds, dx) \\ &+ \int_{[0, t] \times \mathbb{R}^m} \int_0^{t-u} \langle F, \Psi \rangle(t-u-s, \theta_0, \theta) \varphi(s, x, \theta_0) ds \frac{1}{n} M^{(n)}(du, dx). \end{aligned}$$

Recalling that $\bar{\lambda}(\cdot, \theta_0, \theta_0) = \mu(\cdot, \theta_0) + \langle F, \Psi \rangle(\cdot, \theta_0) \star \mu(\cdot, \theta_0)$ this yields expression (41) at $\theta = \theta_0$. Reinserting the obtained decomposition for $\lambda_s^{(n)}(\theta_0)$ into (42) and using Fubini's theorem again ends the proof. \square

It is useful to observe the properties of the kernel φ extend readily to the resolvent Ψ . When φ is split in the sense of Assumption 2, Ψ is split too with, for any $t, x \in \mathbb{R}^+ \times \mathbb{R}^m$ and $\theta \in \Theta$,

$$(44) \quad \Psi: (t, x, \theta_0, \theta) = g(x, \theta) \odot \left(f(\cdot, \theta) + f(\cdot, \theta) \star \langle F, \Psi \rangle(\cdot, \theta_0, \theta_0) \right)(t).$$

Furthermore, for any $t, s \in [0, T]$ and $k \in \mathbb{N}^*$

$$\|\langle F, \varphi \rangle'(t-s, \theta_0) \langle F, \varphi \rangle^{*k}(s, \theta_0)\|_2 \leq \sup_{u \in [0, T]} \|\langle F, \varphi \rangle'(u, \theta_0)\|_2 \|\langle F, \varphi \rangle^{*k}(s, \theta_0)\|_2,$$

where $\langle F, \varphi \rangle^{*k}$ is integrable over \mathbb{R}^+ under Assumption 9 and *a fortiori* integrable over $[0, T]$, hence $\langle F, \varphi \rangle^{*k+1} \in \mathcal{C}^1([0, T])$ for any $k \in \mathbb{N}$. From similar elementary bounds and the dominated convergence theorem, one finds for any $\theta \in \Theta$ that $f(\cdot, \theta) + f(\cdot, \theta) \star \langle F, \Psi \rangle(\cdot, \theta_0, \theta_0) \in \mathcal{C}^1([0, T])$ and Ψ thus writes as the Hadamard product of g with a continuously differentiable function of $t \in [0, T]$.

6. PROOF OF THE CONSISTENCY OF THE MLE

6.1. Technical results. Our proof differs somewhat from the univariate case of Chen & Hall [9] in which the MLE is seen as a critical point of $\mathcal{L}(n, \cdot)$. Such characterisation falls short in the multivariate case, where θ_0 is not necessarily an inner point of Θ . As mentioned in the introduction, an M-estimator approach is preferred instead. We are then compelled to prove the convergence in probability of the local likelihood ratio

$$(45) \quad \frac{1}{n} \Lambda_n(\theta, \theta_0) = \frac{2}{n} (\mathcal{L}(n, \theta) - \mathcal{L}(n, \theta_0))$$

to some limit constrast function

$$(46) \quad \bar{\Lambda}(\theta, \theta_0) = 2 \sum_{k=1}^K \int_0^T \ln \left\{ \frac{\bar{\lambda}_k(s, \theta, \theta_0)}{\bar{\lambda}_k(s, \theta_0, \theta_0)} \right\} \bar{\lambda}_k(s, \theta_0, \theta_0) - (\bar{\lambda}_k(s, \theta, \theta_0) - \bar{\lambda}_k(s, \theta_0, \theta_0)) ds,$$

as will be precised in Lemma 10 (the multiplicative factor 2 serves so as to homogenise (45) with future notation). From (16), the random function Λ_n (45) writes as a functional of the intensity process $(\lambda_s^{(n)}(\theta))$. From the definition (6) of $(N_t^{(n)})$ as a sum process and the linearity of the intensity, for any $\theta \in \Theta$, we may write $\int_0^T \lambda_t^{(n)}(\theta) dt = \sum_{k=1}^n Y_k(\theta)$ where the $Y_k(\theta)$ are i.i.d and have the distribution of $\int_0^T \lambda_t^{(1)}(\theta) dt$. One may therefore already observe that, from the classical law of large numbers,

$$(47) \quad \frac{1}{n} \int_0^T \lambda_s^{(n)}(\theta) ds \xrightarrow[n \rightarrow \infty]{\mathbb{P}(\theta_0)\text{-a.s.}} \int_0^T \bar{\lambda}(s, \theta, \theta_0) ds.$$

Corollary 2 below will serve to extend such results to a broad class of functionals of the intensity, encompassing Λ_n . We first prove a uniform convergence for the type of processes appearing in decomposition (41).

Lemma 7. *Let $v \in L^2(F(dx))$. For any $\theta \in \Theta$,*

$$\begin{aligned} \sup_{t \in [0, T]} \left\| \int_{[0, t] \times \mathbb{R}^m} v(x) \frac{1}{n} \mathbf{M}^{(n)}(ds, dx) \right\|_2 &= o_{\mathbb{P}(\theta)}(1) \\ \sup_{t \in [0, T]} \left\| \int_{[0, t] \times \mathbb{R}^m} v(x) \frac{1}{\sqrt{n}} \mathbf{M}^{(n)}(ds, dx) \right\|_2 &= O_{\mathbb{P}(\theta)}(1) \end{aligned}$$

Proof of Lemma 7. Let $v = (v_{kl}) \in L^2(F(dx))$. For any $n \in \mathbb{N}^*$. We define the process

$$(48) \quad \tilde{\mathbf{M}}_t^{(n)} = \int_{[0, t] \times \mathbb{R}^m} v(x) \mathbf{M}^{(n)}(ds, dx) = \left(\sum_{l=1}^K \int_{[0, t] \times \mathbb{R}^m} v_{kl}(x) M_l^{(n)}(ds, dx) \right)_k.$$

Since $v \in L^2(F(dx))$, $(\tilde{\mathbf{M}}_t)$ is a square integrable martingale (see Brémaud [8][Theorem 5.1.33 p. 170]). Using Remark 1, the covariation of $(\tilde{\mathbf{M}}_t)$ simplifies as

$$\langle \tilde{M}_{k,\cdot}^{(n)}, \tilde{M}_{l,\cdot}^{(n)} \rangle_t = \sum_{p=1}^K \int_0^t \int_{\mathbb{R}^m} v_{kp}(x) v_{lp}(x) \lambda_{p,s}^{(n)}(\theta_0) ds F_p(dx).$$

Hence the predictable compensator of the sub-martingale $\|\tilde{\mathbf{M}}\|_2^2$ simplifies too, so that

$$\begin{aligned} \left\| \frac{1}{n} \tilde{\mathbf{M}}^{(n)} \right\|_2^2 &= \sum_{k=1}^K \left\langle \frac{1}{n} \tilde{\mathbf{M}}_{k,\cdot}^{(n)}, \frac{1}{n} \tilde{\mathbf{M}}_{k,\cdot}^{(n)} \right\rangle \\ &= \left(\left\| \frac{1}{n} \tilde{\mathbf{M}}_t^{(n)} \right\|_2^2 - \frac{1}{n} \int_0^t \left\{ \sum_{k=1}^K \sum_{p=1}^K \int_{\mathbb{R}^m} v_{kp}^2(x) F_k(dx) \right\} \frac{1}{n} \lambda_{p,s}^{(n)}(\theta_0) ds \right)_{k,t} \end{aligned}$$

is also a martingale. From this it follows, together with Lemma 1 and the positivity of the $\lambda_{k,\cdot}^{(n)}$,

$$\mathbb{E} \left[\left\| \frac{1}{n} \tilde{\mathbf{M}}_T^{(n)} \right\|_2^2 \right] \leq \frac{1}{n} \left(\sum_{k=1}^K \sum_{p=1}^K \int_{\mathbb{R}^m} v_{kp}^2(x) F_k(dx) \right) \|H(T, \theta)\|_1,$$

where $H(\cdot, \cdot)$ is defined in (14), and the same bound with n^{-1} omitted holds for $n^{-1/2} \tilde{\mathbf{M}}$. Applying Doob's maximal inequality to the sub-martingale $\|\tilde{\mathbf{M}}\|_2^2$ with the bound above yields Lemma 7. \square

Corollary 2. *Under Assumptions 1 to 4 and Assumption 9, for any $\theta \in \Theta$,*

$$\sup_{t \in [0, T]} \left\| \frac{1}{n} \lambda_t^{(n)}(\theta) - \bar{\lambda}(t, \theta, \theta_0) \right\|_2 = o_{\mathbb{P}(\theta_0)}(1).$$

Remark 11. *The splitting Assumption 2 proves instrumental at this point. It allows for the integral of $u(\cdot) \odot v(\cdot)$ against the counting measure $N^{(n)}(ds, dx)$ to be interpreted as an integral of $u(\cdot)$ against the Riemann-Stieltjes measure $\int_{\mathbb{R}} v(x) N^{(n)}(ds, dx)$. Thus, usual integration by parts applies (see: Daley & Vere-Jones [14][lemma 4.6.1]).*

Proof of Corollary 2. Let $\theta \in \Theta$. For any $n \in \mathbb{N}^*$ and $t \in [0, T]$,

$$\frac{1}{n} \lambda_t^{(n)}(\theta) - \bar{\lambda}(t, \theta, \theta_0) = \int_{[0, t] \times \mathbb{R}^m} \Psi(t-s, x, \theta_0, \theta) \frac{1}{n} \mathbf{M}^{(n)}(ds, dx).$$

From (44), we have some $u: t, \vartheta, \theta \mapsto u(t, \vartheta, \theta) \in \mathcal{M}_K(\mathbb{R})$ such that $u(\cdot, \theta, \theta_0) \in \mathcal{C}^1(\mathbb{R}^+)$ for any $\theta \in \Theta$, and $\Psi(t, x, \theta_0, \theta) = u(t, \theta, \theta_0) \odot g(x, \theta)$. Integrating by parts, for any $n \in \mathbb{N}^*$ and $t \in [0, T]$,

$$\begin{aligned} \frac{1}{n} \lambda_t^{(n)}(\theta) - \bar{\lambda}(t, \theta, \theta_0) &= u(0, \theta, \theta_0) \odot \int_{[0, t] \times \mathbb{R}^m} g(x, \theta) \frac{1}{n} \mathbf{M}(du, dx) \\ &\quad + \int_0^t u'(t-s, \theta, \theta_0) \odot \int_{[0, s] \times \mathbb{R}^m} g(x, \theta) \frac{1}{n} \mathbf{M}(du, dx) ds, \end{aligned}$$

hence for any $\theta \in \Theta$ and $n \in \mathbb{N}^*$,

$$(49) \quad \sup_{t \in [0, T]} \left\| \frac{1}{n} \lambda_t^{(n)}(\theta) - \bar{\lambda}(t, \theta, \theta_0) \right\|_2 \leq C(u(\cdot, \theta, \theta_0)) \sup_{t \in [0, T]} \left\| \int_{[0, t] \times \mathbb{R}^m} g(x, \theta) \frac{1}{n} \mathbf{M}^{(n)}(du, dx) \right\|_2,$$

where

$$C(u(\cdot, \theta, \theta_0)) = \|u(0, \theta, \theta_0)\|_2 + \int_0^T \|u'(s, \theta, \theta_0)\|_2 ds.$$

Corollary 2 then proceeds directly from Lemma 7. \square

We now extend the previous convergence to regular functions of the intensity.

Lemma 8. Let $\Phi \in \mathcal{C}^0(\mathbb{R}^{+*}, \mathbb{R})$. Under Assumptions 1 to 4 and Assumption 9, for any $\theta \in \Theta$, $k \in \llbracket 1, K \rrbracket$,

$$\sup_{t \in [0, T]} \left| \Phi\left(\frac{1}{n} \lambda_{k,t}^{(n)}(\theta)\right) - \Phi(\bar{\lambda}_k(t, \theta, \theta_0)) \right| = o_{\mathbb{P}(\theta_0)}(1).$$

Proof of Lemma 8. Let $\theta \in \Theta$. Under Assumption 1, for any $n \in \mathbb{N}^*$, $\lambda_k^{(n)}$ is strictly positive and $\Phi(n^{-1} \lambda_{k,s}^{(n)}(\theta))$ is indeed well defined for any $k \in \llbracket 1, K \rrbracket$. By the inverse triangle inequality, for any $t \in [0, T]$, and any $k \in \llbracket 1, K \rrbracket$,

$$(50) \quad \inf_{s \in [0, T]} \bar{\lambda}_k(s, \theta, \theta_0) + o_{\mathbb{P}(\theta_0)}(1) \leq \frac{1}{n} \lambda_{k,t}^{(n)}(\theta) \leq \sup_{s \in [0, T]} \bar{\lambda}_k(s, \theta, \theta_0) + o_{\mathbb{P}(\theta_0)}(1),$$

where the negligible terms are uniform in time. Hence, with probability tending to 1 under $\mathbb{P}(\theta_0)$ as n grows, $s \mapsto n^{-1} \lambda_s^{(n)}(\theta)$ eventually lies in some compact interval $J \subset \mathbb{R}$ in which $s \mapsto \bar{\lambda}(s, \theta, \theta_0)$ takes values. Let $\varepsilon > 0$. On account of Heine's Lemma, Φ is uniformly continuous over any compact subset of \mathbb{R} , and we have some $\eta > 0$ such that, for any two $x, y \in J$ verifying $|x - y| < \eta$, one gets $|\Phi(x) - \Phi(y)| \leq \varepsilon$. From Corollary 2, for any sufficiently large n , one has for any $t \in [0, T]$ that

$$\left| \frac{1}{n} \lambda_{k,t}^{(n)}(\theta) - \bar{\lambda}_k(t, \theta, \theta_0) \right| < \eta,$$

with probability tending to 1, thus proving Lemma 8. \square

The results above were written in the context of separable functions in the sense of Assumption 2. They apply immediately to finite sum of such functions, provided they satisfy the required regularity conditions. Together with Assumptions 2, 7 and 8, this remark guarantees the convergence of corollary 2 still holds for the class of functions on $\mathbb{R}^+ \times \mathbb{R}^m$ spanned by $\partial_\theta^p \varphi(\cdot, \cdot, \theta)$, $\theta \in \Theta$, $p \in \{0, 1, 2\}$. Consequently, for any $\theta \in \Theta$ and $k \in \llbracket 1, K \rrbracket$,

$$(51) \quad \sup_{s \in [0, T]} \left\| \frac{1}{n} \partial_\theta^{\otimes p} \lambda_{k,s}^{(n)}(\theta) - \partial_\theta^{\otimes p} \bar{\lambda}_k(s, \theta, \theta_0) \right\|_1 \xrightarrow[n \rightarrow \infty]{\mathbb{P}(\theta_0)} 0.$$

Such results will prove useful as we tackle derivatives of $(\lambda_t^{(n)})$ in the following sections. Applying the arguments of the proof of Lemma 8 to (51), we may already state the results below before getting back to the consistency of $\hat{\theta}_n$.

Lemma 9. Under assumptions 1 to 9, for any $k \in \llbracket 1, K \rrbracket$ and $i, j \in \llbracket 1, d \rrbracket$, and $t \in [0, T]$, denote

$$Y_{k,ij}^{(n)}(t, \theta) = (\lambda_{k,t}^{(n)}(\theta), \partial_{\theta_i} \lambda_{k,t}^{(n)}(\theta), \partial_{\theta_j} \lambda_{k,t}^{(n)}(\theta), \partial_{\theta_i} \partial_{\theta_j} \lambda_{k,t}^{(n)}(\theta)),$$

and

$$\bar{Y}_{k,ij}(t, \theta, \theta_0) = (\bar{\lambda}_k(t, \theta, \theta_0), \partial_{\theta_i} \bar{\lambda}_k(t, \theta, \theta_0), \partial_{\theta_j} \bar{\lambda}_k(t, \theta, \theta_0), \partial_{\theta_i} \partial_{\theta_j} \bar{\lambda}_k(t, \theta, \theta_0)).$$

Then, for any $\theta \in \Theta$ and any continuous function Φ from $(0, \infty) \times \mathbb{R}^3$ to \mathbb{R} ,

$$\sup_{t \in [0, T]} \left| \Phi\left(\frac{1}{n} Y_{k,ij}^{(n)}(t, \theta)\right) - \Phi(\bar{Y}_{k,ij}(t, \theta, \theta_0)) \right| = o_{\mathbb{P}(\theta_0)}(1).$$

6.2. Proof of proposition 1. The proof relies essentially on Lemma 10 which states the convergence of the contrast function Λ_n towards the limit function $\bar{\Lambda}$ defined in (46). Before getting to its proof, remark as in Ogata [35][proof of Lemma 3 pp. 252–253] that $\bar{\Lambda}$ re-expresses as

$$\theta \mapsto \bar{\Lambda}(\theta, \theta_0) = \sum_{k=1}^K \int_0^t L\left(\frac{\bar{\lambda}_k(s, \theta, \theta_0)}{\bar{\lambda}_k(s, \theta_0, \theta_0)}\right) \bar{\lambda}_k(s, \theta_0, \theta_0) ds,$$

where

$$L: y \in (0, \infty) \mapsto \ln(y) + \frac{1}{y} - 1$$

verifies $L(y) \leq 0$ for any $y > 0$ with equality if and only if $y = 1$. The function $\theta \mapsto \bar{\Lambda}(\theta, \theta_0)$ thus attains a separable maximum at θ_0 as long as $\lambda_k(\cdot, \theta, \theta_0)$ and $\bar{\lambda}_k(\cdot, \theta_0, \theta_0)$ differ on an interval of non-null measure when $\theta \neq \theta_0$, which always holds true under assumptions 9 and 5. When Lemma 10 holds, the M-estimator master theorem (see Van der Vaart [50][Theorem 5.7]) therefore applies, ensuring the consistency of the MLE as $n \rightarrow \infty$.

Lemma 10. *Under Assumptions 1 to 5 and Assumption 9*

$$(52) \quad \sup_{\theta \in \Theta} \left\| \frac{1}{n} \Lambda_n(\theta_0, \theta) - \bar{\Lambda}(\theta, \theta_0) \right\|_1 \xrightarrow[n \rightarrow \infty]{\mathbb{P}(\theta_0)} 0.$$

Proof of Lemma 10. We begin by showing convergence (52) holds pointwise at any $\theta \in \Theta$. Let $\theta \in \Theta$. Firstly,

$$\sum_{k=1}^K \int_0^T \frac{1}{n} \{ \lambda_{k,s}^{(n)}(\theta_0) - \lambda_{k,s}^{(n)}(\theta) \} - \{ \bar{\lambda}_k(s, \theta_0, \theta_0) - \bar{\lambda}_k(s, \theta, \theta_0) \} ds$$

vanishes in probability under $\mathbb{P}(\theta_0)$ as $n \rightarrow \infty$ as a direct application of the uniform convergence from Lemma 8. Secondly, recalling that $h(\cdot, \theta_0) = \bar{\lambda}(\cdot, \theta_0, \theta_0)$,

$$(53) \quad \begin{aligned} & \sum_{k=1}^K \int_0^T \ln\left(\frac{\lambda_{k,s}^{(n)}(\theta_0)}{\lambda_{k,s}^{(n)}(\theta)}\right) \frac{1}{n} dN_{k,s}^{(n)} - \ln\left(\frac{\bar{\lambda}_k(s, \theta_0, \theta_0)}{\bar{\lambda}_k(s, \theta, \theta_0)}\right) h_k(s, \theta_0) ds \\ &= \sum_{k=1}^K \int_0^T \ln\left(\frac{\bar{\lambda}_k(s, \theta_0, \theta_0)}{\bar{\lambda}_k(s, \theta, \theta_0)}\right) \frac{1}{n} dM_{k,s}^{(n)} \\ &+ \sum_{k=1}^K \int_0^T \ln\left(\frac{\bar{\lambda}_k(s, \theta_0, \theta_0)}{\bar{\lambda}_k(s, \theta, \theta_0)}\right) \left\{ \frac{1}{n} \lambda_{k,s}^{(n)}(\theta_0) - \bar{\lambda}_k(s, \theta_0, \theta_0) \right\} ds \\ &- \sum_{k=1}^K \int_0^T \left\{ \ln\left(\frac{\frac{1}{n} \lambda_{k,s}^{(n)}(\theta_0)}{\frac{1}{n} \lambda_{k,s}^{(n)}(\theta)}\right) - \ln\left(\frac{\bar{\lambda}_k(s, \theta_0, \theta_0)}{\bar{\lambda}_k(s, \theta, \theta_0)}\right) \right\} \frac{1}{n} dN_{k,s}^{(n)}. \end{aligned}$$

Now, the third and last term on the right-hand side of (53) is bounded in absolute value by

$$(54) \quad \sup_{t \in [0, T]} \sum_{k=1}^K \left| \ln\left(\frac{\frac{1}{n} \lambda_{k,s}^{(n)}(\theta_0)}{\frac{1}{n} \lambda_{k,s}^{(n)}(\theta)}\right) - \ln\left(\frac{\bar{\lambda}_k(s, \theta_0, \theta_0)}{\bar{\lambda}_k(s, \theta, \theta_0)}\right) \right| \left\| \frac{1}{n} \mathbf{N}_T^{(n)} \right\|_1,$$

where the two multiplicative factors in (54) are respectively $o_{\mathbb{P}(\theta_0)}(1)$ and $O_{\mathbb{P}(\theta_0)}(1)$ from Lemma 8 and the classical law of large numbers. As for the second term of (53), we have the bound

$$\sum_{k=1}^K \int_0^T \left| \ln\left(\frac{\bar{\lambda}_k(s, \theta_0, \theta_0)}{\bar{\lambda}_k(s, \theta, \theta_0)}\right) \right| ds \sup_{t \in [0, T]} \left\| \frac{1}{n} \lambda_{k,s}^{(n)}(\theta_0) - \bar{\lambda}_k(s, \theta, \theta_0) \right\|,$$

where $s \mapsto \bar{\lambda}(s, \theta, \theta_0)$ is locally bounded for any $\theta \in \Theta$ and Corollary 2 guarantees the second factor of the preceding decomposition vanishes as $n \rightarrow \infty$. Finally, a straightforward adaptation of the proof for Lemma 7 shows the first and remaining term on the right-hand side of (53) is $o_{\mathbb{P}(\theta_0)}(1)$ too as a result of Doob's maximal inequality. Consequently, for every $\theta \in \Theta$

$$(55) \quad \frac{1}{n} \Lambda_n(\theta_0, \theta) \xrightarrow[n \rightarrow \infty]{\mathbb{P}(\theta_0)} \bar{\Lambda}(\theta, \theta_0).$$

There remains to extend (55) uniformly over Θ . By virtue of Lemma 2 it suffices to show the stochastic equicontinuity of $n^{-1} \Lambda_n - \bar{\Lambda}$, which reduces to that of Λ_n since $\bar{\Lambda}$ is deterministic and continuous over the compact Θ . Let $\varepsilon > 0$. The baseline μ and the kernel φ are uniformly equicontinuous in θ . For any sufficiently close ϑ_1, ϑ_2 in Θ thus, one has for any $t \in [0, T]$ and $n \in \mathbb{N}^*$

$$\begin{aligned} & \left\| \frac{1}{n} \lambda_t^{(n)}(\vartheta_1) - \frac{1}{n} \lambda_t^{(n)}(\vartheta_2) \right\|_1 \\ & \leq \|\mu(t, \vartheta_1) - \mu(t, \vartheta_2)\|_1 + \int_0^T \|\varphi(t-s, x, \vartheta_1) - \varphi(t-s, x, \vartheta_2)\|_1 \frac{1}{n} N_{k,s}^{(n)}(ds, dx) \\ & \leq \varepsilon + \varepsilon \left\| \frac{1}{n} \mathbf{N}_T^{(n)} \right\|_1 = \varepsilon(1 + \|H(T, \theta_0)\|_1 + o_{\mathbb{P}(\theta_0)}(1)) = \varepsilon O_{\mathbb{P}(\theta_0)}(1), \end{aligned}$$

where $H(\cdot, \cdot)$ is defined in (14). Hence $n^{-1} \lambda^{(n)}$ is equi-continuous in the sense of Lemma 2. The equicontinuity of $n^{-1} \Lambda(n, \cdot)$, deduces directly from that of the normalized intensity by applying Heine's Lemma in the same fashion as in the proof of Lemma 8 and remarking again that $\frac{1}{n} N^{(n)} = O_{\mathbb{P}(\theta_0)}(1)$. The proof of Lemma 10 is then complete, yielding the consistency of the MLE as a consequence. \square

7. PROOF OF THE WEAK CONVERGENCE OF THE LIKELIHOOD RATIO

This section contains the proofs for Theorem 1 and Proposition 3. A classical argument for the weak convergence of the MLE relies on the second order expansion

$$(56) \quad n^{-\frac{1}{2}} \mathcal{S}(n, \theta_0) = n^{-\frac{1}{2}} \mathcal{S}(n, \hat{\theta}_n) + n^{-1} \partial_{\theta} \mathcal{S}(n, \theta_0) \sqrt{n}(\hat{\theta} - \theta_0) + o_{\mathbb{P}}(\sqrt{n}(\hat{\theta} - \theta_0)).$$

Under standard conditions, after cancelling the score at $\hat{\theta}_n$, the asymptotic normality of $n^{-\frac{1}{2}} \mathcal{S}(n, \theta_0)$ together with Slutsky's Lemma yields a central limit theorem for the MLE. The present setting differs in that $\mathcal{S}(n, \hat{\theta}_n)$ may now take non-null values, pulling apart $\sqrt{n}(\hat{\theta}_n - \theta_0)$ and the score at θ_0 . This leads to $\sqrt{n}(\hat{\theta}_n - \theta_0)$ behaving as the minimizer in h over $\Theta - \{\theta_0\}$ of the quadratic form

$$(57) \quad q_Z: h \mapsto (Z - h)^T I(\theta_0)(Z - h)$$

with $Z \sim \mathcal{N}(0, I(\theta_0)^{-1})$. Approximation (57) was introduced by Self & Liang [42][Theorem 3 and equ. (3.1) p. 607]. The precise conditions for its validity are detailed in the master theorem for constrained M-estimators of Geyer [19][Theorem 4.4 and Remark p. 2004], the verification of which assumptions' constitutes the main objective of the next few results. Notably, we prove in proposition 4 the asymptotic normality of the score at the true parameter, extending a result of Chen & Hall [9][equ. (12) p. 1021] to the multivariate case.

7.1. Technical results. The limit function $\bar{\Lambda}$ plays the role of $\mathbb{E}[n^{-1}\mathcal{L}(n, \cdot)]$ in Geyer's [19] setting. Lemma 11 shows the derivatives of $\bar{\Lambda}$ express as uniform limits of derivatives of $n^{-1}\mathcal{L}(n, \cdot)$. This will serve in the sequel so as to clarify some properties of $\bar{\Lambda}$. Additionally, we recover the consistency of $\mathcal{J}(n, \hat{\theta}_n)$ as an estimator of $I(\theta_0)$.

Lemma 11. *Under Assumptions 1 to 9,*

$$(58) \quad \sup_{\theta \in \Theta} \left\| \frac{1}{n} \mathbf{S}(n, \theta) - \partial_{\theta} \bar{\Lambda}(\theta, \theta_0) \right\|_2 = o_{\mathbb{P}(\theta_0)}(1),$$

$$(59) \quad \sup_{\theta \in \Theta} \left\| \frac{1}{n} \partial_{\theta} \mathbf{S}(n, \theta) - \partial_{\theta}^{\otimes 2} \bar{\Lambda}(\theta, \theta_0) \right\|_2 = o_{\mathbb{P}(\theta_0)}(1).$$

Moreover, using notation (32) for the empirical information,

$$(60) \quad \mathcal{J}(n, \hat{\theta}_n) \xrightarrow[n \rightarrow \infty]{\mathbb{P}(\theta_0)} I(\theta_0)$$

Proof. The normalised score $n^{-1}\mathbf{S}(n, \cdot)$ and its derivative write coordinate-wise as functionals of the type

$$\sum_{k=1}^K \int_0^T U\left(\frac{1}{n} Y_{k,ij}^{(n)}(s, \theta)\right) \frac{1}{n} dN_{k,s}^{(n)} + \int_0^T V\left(\frac{1}{n} Y_{k,ij}^{(n)}(s, \theta)\right) ds,$$

where U, V are continuous functions and $Y^{(n)}$ is defined in Lemma 9. Convergences (58) and (59) then proceed from Lemma 9 in the exact same fashion Lemma 10 proceeds from Lemma 8 and we do not detail their proof. From the same arguments, $\mathcal{J}(n, \cdot)$ converges uniformly in probability towards the asymptotic information function $I(\cdot)$. Together with the consistency of the MLE this yields (60) \square

Remark 12. *Other possibilities exist for an estimator of $I(\theta_0)$. The one we retained does not require the computation of the limit function h which is often non-explicit.*

Proposition 4. *Under Assumptions 1 to 9,*

$$(61) \quad \frac{1}{\sqrt{n}} \mathbf{S}(n, \theta_0) \xrightarrow[n \rightarrow \infty]{\mathcal{L}(\mathbb{P}(\theta_0))} \mathcal{N}(0, I(\theta_0))$$

Proof. Under $\mathbb{P}(\theta_0)$, the score $\mathcal{S}(n, \theta_0)$ exhibits a martingale structure :

$$(62) \quad \begin{aligned} \frac{1}{\sqrt{n}} \mathbf{S}(n, \theta_0) &= \sum_{k=1}^K \int_0^T \frac{1}{\sqrt{n}} \frac{\partial_{\theta} \mu_k(s, \theta_0) + \int_0^s \int_{\mathbb{R}^m} \sum_l \partial_{\theta} \varphi_{kl}(s-u, x, \theta_0) \frac{1}{n} N_l^{(n)}(du, dx)}{\mu_k(s, \theta_0) + \int_0^s \int_{\mathbb{R}^m} \sum_l \varphi_{kl}(s-u, x, \theta_0) \frac{1}{n} N_l^{(n)}(du, dx)} dM_{k,s}^{(n)} \\ &= \sum_{k=1}^K \int_0^T U_k^{(n)}(s) dM_{k,s}^{(n)}, \end{aligned}$$

from which we define the process

$$(\tilde{M}_t^{(n)}) = \left(\sum_{k=1}^K \int_0^t U_k^{(n)}(s) dM_{k,s}^{(n)} \right).$$

Using remark 1, the predictable co-variation of \tilde{M} expresses as

$$(63) \quad \left\langle \tilde{M}^{(n)}, \tilde{M}^{(n)} \right\rangle (t) = \int_0^t \sum_{k=1}^K \frac{\left(\partial_{\theta} \mu_k(s, \theta_0) + \int_0^s \int_{\mathbb{R}^m} \sum_l \partial_{\theta} \varphi_{kl}(s-u, \theta_0) \frac{1}{n} N_l^{(n)}(du, dx) \right)^{\otimes 2}}{\mu_k(s, \theta_0) + \int_0^s \int_{\mathbb{R}^m} \sum_l \varphi_{kl}(s-u, \theta_0) \frac{1}{n} N_l^{(n)}(du, dx)} ds.$$

Thanks to this simplification, expression (63) is quite similar to the one Chen & Hall [9] [equ. 12 p. 1021] or Ogata [35][Theorem 4 p. 255] obtain for the covariation of $\mathcal{S}(n, \theta_0)$ in the univariate case, hence we may use the same arguments. Namely, it suffices to show that \tilde{M} verifies a martingale central limit theorem, as introduced by Rebolledo [38] (see also Jacod & Shiryaev [28][Theorem VIII-3.11]). Define

$$\tilde{M}_\varepsilon^{(n)}(t) = \sum_{k=1}^K \int_0^t U_k^{(n)}(s) \mathbb{1}_{|U_k^{(n)}| > \varepsilon} dM_{k,s}^{(n)},$$

where $|\cdot|$ and $>$ are to be understood component-wise. The necessary conditions of the martingale central limit theorem can be re-expressed as in Andersen *et al.* [1][Th. II.5.1 and eq. (2.5.8) p. 83-84]) as

$$(64) \quad \forall t \in [0, T], \quad \langle \tilde{M}^{(n)}, \tilde{M}^{(n)} \rangle(t) \xrightarrow[n \rightarrow \infty]{\mathbb{P}(\theta_0)} V(t),$$

$$(65) \quad \forall \varepsilon > 0, \forall t \in [0, T], \quad \langle \tilde{M}_\varepsilon^{(n)}, \tilde{M}_\varepsilon^{(n)} \rangle(t) \xrightarrow[n \rightarrow \infty]{\mathbb{P}(\theta_0)} 0.$$

Firstly, using Lemma 8 with equation (63), condition (64) is immediately satisfied with

$$V(t) = \int_0^t \frac{(\partial_\theta \mu(s, \theta_0) + \int_0^s \partial_\theta \langle F, \varphi \rangle(s-u, \theta_0) h(u, \theta_0) du)^{\otimes 2}}{\mu(s, \theta_0) + \int_0^s \langle F, \varphi \rangle(s-u, \theta_0) h(u, \theta_0) du} ds.$$

Secondly, using remark 1 again, the predictable co-variation of $\tilde{M}_\varepsilon^{(n)}$ simplifies as in (63),

$$\langle \tilde{M}_\varepsilon^{(n)}, \tilde{M}_\varepsilon^{(n)} \rangle(t) = \sum_{k=1}^K \int_0^t \frac{(\partial_\theta \mu_k(s, \theta_0) + \int_0^s \int_{\mathbb{R}^m} \sum_l \partial_\theta \varphi_{kl}(s-u, x, \theta_0) \frac{1}{n} N_l^{(n)}(du, dx))^{\otimes 2}}{\mu_k(s, \theta_0) + \int_0^s \int_{\mathbb{R}^m} \sum_l \varphi_{kl}(s-u, x, \theta_0) \frac{1}{n} N_l^{(n)}(du, dx)} \mathbb{1}_{|U_k^{(n)}| > \varepsilon} ds.$$

where the integrand converges to 0 uniformly in time since for every $k \in \llbracket 1, K \rrbracket$,

$$\sup_{t \in [0, T]} |U_k^{(n)}(t)| = O_{\mathbb{P}(\theta_0)}\left(\frac{1}{\sqrt{n}}\right).$$

as a result of Lemma 9. The Lindeberg condition (65) is thus satisfied too, and $n^{-\frac{1}{2}} \tilde{M}^{(n)}$ converges in law for the Skorokhod topology to a Gaussian process with covariance V . Since convergence in Skorokhod topology implies convergence of finite-dimensional distributions, setting $t = T$, (61) holds as a consequence. \square

We now introduce an extension of Lemma 7 so as to ensure later the \sqrt{n} -consistency of certain functionals of the intensity.

Lemma 12 (A uniform extension of Lemma 7). *Under assumptions 1 to 9,,*

$$(66) \quad \sup_{\theta \in \Theta} \sup_{t \in [0, T]} \left\| \int_{[0, t] \times \mathbb{R}^m} g(x, \theta) \frac{1}{\sqrt{n}} \mathbf{M}^{(n)}(du, dx) \right\|_1 = O_{\mathbb{P}(\theta_0)}(1)$$

Proof of lemma 12. Recall from Lemma 7 that (66) already holds pointwise in θ . Consider the family of processes $\{Z^{(n)}(\theta)\}$ defined for any $t \in [0, T]$ and $\theta \in \Theta$ by

$$(67) \quad Z_t^{(n)}(\theta) = \int_{[0, t] \times \mathbb{R}^m} g(x, \theta) \frac{1}{\sqrt{n}} \mathbf{M}^{(n)}(ds, dx).$$

In view of Lemma 2, it thus suffices to show $\{\sup_{t \in [0, T]} \|Z_t^{(n)}(\theta)\|_1\}$ is stochastically equi-continuous in θ for Lemma 12 to hold. This will derive from the chaining argument detailed in Lemma 3, for which we require some concentration inequality on increments along θ of the family (67). We will work with the metrics $d_p : \Theta^2 \rightarrow \mathbb{R}^+$ defined for any $p \in (0, 1]$ by

$$d_p(\theta, \nu) = \|\theta - \nu\|_2^p.$$

Now, from the reverse triangle inequality, for any $\theta, \nu \in \Theta$,

$$(68) \quad \left| \sup_{t \in [0, T]} \|Z_t^{(n)}(\theta)\|_1 - \sup_{t \in [0, T]} \|Z_t^{(n)}(\nu)\|_1 \right| \leq \sup_{t \in [0, T]} \|Z_t^{(n)}(\theta) - Z_t^{(n)}(\nu)\|_1,$$

and, as far as tail bounds are concerned, we can look instead at the family of processes

$$Z_t^{(n)}(\theta, \nu) = \int_{[0, t] \times \mathbb{R}^m} (g(x, \theta) - g(x, \nu)) \frac{1}{\sqrt{n}} \mathbf{M}^{(n)}(ds, dx).$$

Note also that for any $n \in \mathbb{N}$, $\theta, \nu \in \Theta$ and $\eta > 0$,

$$(69) \quad \mathbb{P} \left[\sup_{t \in [0, T]} \|Z_t^{(n)}(\theta, \nu)\|_1 > \eta d_p(\theta, \nu) \right] \leq \sum_{k=1}^K \mathbb{P} \left[\sup_{t \in [0, T]} |Z_{k,t}^{(n)}(\theta, \nu)| > \frac{\eta}{K} d_p(\theta, \nu) \right]$$

and we can study each summand in the right-hand side of (69) separately to alleviate the notation. For any $n \in \mathbb{N}^*$, any $\theta, \nu \in \Theta$ and any $k \in \llbracket 1, K \rrbracket$, from Remark 1, the predictable variation of $(Z_{k,t}^{(n)}(\theta, \nu))$ writes

$$\sum_{l=1}^K \int_{\mathbb{R}^m} (g_{kl}(x, \theta) - g_{kl}(x, \nu))^2 F_l(dx) \frac{1}{n} \int_0^t \lambda_{l,s}^{(n)}(\theta_0) ds,$$

and is hence bounded uniformly in t by

$$\sum_{i,j} \int_{\mathbb{R}^m} |g_{ij}(x, \theta) - g_{ij}(x, \nu)|^2 F_j(dx) \frac{1}{n} \sum_{k=1}^K \int_0^T \lambda_{k,s}^{(n)}(\theta_0) ds.$$

Firstly, under Assumption 8, $x, \theta \mapsto g(x, \theta)$ is uniformly p^* -Hölder in θ for some $p^* \in (0, 1]$, and there is some $C > 0$ such that

$$(70) \quad \sum_{i,j} \int_{\mathbb{R}^m} |g_{ij}(x, \theta) - g_{ij}(x, \nu)|^2 F_j(dx) \leq C^2 \|\theta - \nu\|_2^{2p^*} = C^2 d_{p^*}(\theta, \nu)^2.$$

Secondly, for any $\zeta > 0$, from the continuity of the ℓ^1 norm and the classical law of large numbers (47), we have for any sufficiently large n that, $\mathbb{P}(\theta_0)$ -a.s.,

$$(71) \quad \frac{1}{n} \sum_{k=1}^K \int_0^T \lambda_{k,s}^{(n)}(\theta_0) ds \leq (1 + \zeta) \|H(T, \theta_0)\|_1.$$

Setting ζ to some fixed value, say, $\zeta = 1$, and then using (70) and (71) together, one has some $N \in \mathbb{N}^*$ such that, for any $n \geq N$, any $t \in [0, T]$ and any $\theta, \nu \in \Theta$,

$$(72) \quad \langle Z_{k,\cdot}^{(n)}(\theta, \nu), Z_{k,\cdot}^{(n)}(\theta, \nu) \rangle_t \leq 2C^2 \|H(T, \theta_0)\|_1 d_{p^*}(\theta, \nu)^2 = \mathcal{K} d_{p^*}(\theta, \nu)^2$$

$\mathbb{P}(\theta_0)$ -a.s. Furthermore, the jumps of $(Z_{k,t}^{(n)}(\theta, \nu))$ are bounded by

$$(73) \quad \|g(x, \theta) - g(x, \nu)\|_1 \leq C \|\theta - \nu\|_2^p = C d_{p^*}(\theta, \nu).$$

Applying the concentration inequality of Lemma 4 in light of (72) and (73), for any sufficiently large n , one has for any $\eta > 0$ and any $\theta, \nu \in \Theta$,

$$\begin{aligned} \mathbb{P}\left[\sup_{t \in [0, T]} |Z_{k,t}^{(n)}(\theta, \nu)| > \frac{1}{K} \eta d_{p^*}(\theta, \nu)\right] &\leq 2 \exp\left[-\frac{1}{2} \frac{\eta^2 d_{p^*}(\theta, \nu)^2}{K^2 (C d_{p^*}(\theta, \nu)^2 + \mathcal{K} d_{p^*}(\theta, \nu)^2)}\right] \\ &\leq 2 \exp\left[-\frac{1}{2} \frac{\eta^2}{K^2 (C + \mathcal{K})}\right]. \end{aligned}$$

Taking $D^2 = K^2 \max(C, \mathcal{K})$, summing over k in (69), and recalling (68) then, for any large enough n , for any $\eta > 0$ one has for any $\theta, \nu \in \Theta$,

$$\mathbb{P}\left[\left|\sup_{t \in [0, T]} \|Z_t^{(n)}(\theta)\|_1 - \sup_{t \in [0, T]} \|Z_t^{(n)}(\nu)\|_1\right| > \eta d_{p^*}(\theta, \nu)\right] \leq 2K \exp\left[-\frac{1}{2} \frac{\eta^2}{D^2}\right].$$

Apart from the unimportant factor K , this is the sub-exponential bound we needed to apply the chaining Lemma 3. Since we work with parametric families indexed on the compact Θ , the finiteness of the covering integral is immediate in our case. Together with Lemma 2, the chaining Lemma 3 therefore yields Lemma 12. \square

Corollary 3 (A consequence of Lemma 12). *For any $p, m \in \{0, 1\}$ and any $i, j \in \llbracket 1, K \rrbracket$*

$$\sup_{\theta \in \Theta} \sup_{t \in [0, T]} \left\| \sqrt{n} \left\{ \partial_{\theta_i}^p \partial_{\theta_j}^m \frac{1}{n} \lambda_s^{(n)}(\theta) - \partial_{\theta_i}^p \partial_{\theta_j}^m \bar{\lambda}(s, \theta, \theta_0) \right\} \right\|_2 = O_{\mathbb{P}(\theta_0)}(1)$$

Proof. We only consider the configuration $p = m = 0$ since the proof for other cases follow from the same arguments. Consider again the bound (49) from the proof of Corollary 2, whence we have some $u: t, \vartheta, \theta \mapsto u(t, \vartheta, \theta) \in \mathcal{M}_K(\mathbb{R})$ given by

$$u(t, \theta, \theta_0) = f(t, \theta) + f(\cdot, \theta) \star \langle F, \Psi \rangle(\cdot, \theta_0, \theta_0),$$

such that $u(\cdot, \theta, \theta_0) \in \mathcal{C}^1([0, T])$ for any $\theta \in \Theta$ and

$$(74) \quad \sqrt{n} \left\| \frac{1}{n} \lambda_t^{(n)}(\theta) - \bar{\lambda}(t, \theta, \theta_0) \right\|_2 \leq C \sup_{\theta \in \Theta} \sup_{t \in [0, T]} \left\| \int_{[0, t] \times \mathbb{R}^m} g(x, \theta) \frac{1}{\sqrt{n}} \mathbf{M}^{(n)}(du, dx) \right\|_2,$$

where

$$C = \sup_{\theta \in \Theta} \left(\|u(0, \theta, \theta_0)\|_2 + \int_0^T \|u'(s, \theta, \theta_0)\|_2 ds \right)$$

is finite under assumptions 3 and 7. Applying Lemma 12 to (74) yields Corollary 3. \square

Corollary 4 (A consequence of Lemmata 12 and 9). *Let $V, W \in \mathcal{C}^0(\mathbb{R})$, $k \in \llbracket 1, K \rrbracket$, $i, j \in \llbracket 1, d \rrbracket$, and*

$$U_{k,ij}^{(n)}(t, \theta) = V\left(\frac{1}{n} Y_{k,ij}^{(n)}(t, \theta)\right) W\left(\bar{Y}_{k,ij}(t, \theta, \theta_0)\right),$$

where we have used the notation of Lemma 9. Then, for any $q, r \in \{0, 1\}$,

$$\sup_{\theta \in \Theta} \sup_{t \in [0, T]} \left| \int_0^t U_{k,ij}^{(n)}(s, \theta) \sqrt{n} \left\{ \frac{1}{n} \partial_{\theta_i}^q \partial_{\theta_j}^r \lambda_{k,t}^{(n)}(\theta) - \partial_{\theta_i}^q \partial_{\theta_j}^r \bar{\lambda}_k(t, \theta, \theta_0) \right\} \frac{1}{n} dN_{k,s}^{(n)} \right| = O_{\mathbb{P}(\theta_0)}(1)$$

Proof. For any $t \in [0, T]$, $n \in \mathbb{N}^*$ and $\theta \in \Theta$, the expression in Corollary 4 is bounded by the product of the three terms

$$\sup_{\theta \in \Theta, t \in [0, T]} |U_{k,ij}^{(n)}(t, \theta)| \sup_{\theta \in \Theta, t \in [0, T]} \sqrt{n} \left\| \frac{1}{n} \partial_{\theta_i}^p \partial_{\theta_j}^m \lambda_t^{(n)}(\theta) - \partial_{\theta_i}^p \partial_{\theta_j}^m \bar{\lambda}(t, \theta, \theta_0) \right\|_1 \frac{1}{n} \|N_T^{(n)}\|_1,$$

which are all $O_{\mathbb{P}(\theta_0)}(1)$ as a result of Lemma 8, Lemma 12 and the classical law of large numbers, respectively. \square

7.2. Proof of Theorem 1. We are now ready to prove Theorem 1. As mentioned in the introduction, we plan to apply Theorem 4.4 of Geyer [19][p. 2003], which provides sufficient conditions for the epi-convergence in distribution of random functions. More precisely we require a set of four conditions to be verified.

First condition. The limit function $\bar{\Lambda}$ is twice differentiable and satisfies

$$(75) \quad \bar{\Lambda}(\theta) = (\theta - \theta_0)^\top V (\theta - \theta_0) + o(\|\theta - \theta_0\|^2)$$

for some positive definite $V \in \mathcal{M}_d(\mathbb{R})$. The regularity conditions above are immediately verified from $\bar{\Lambda}$ being a uniform limit of twice continuously differentiable functions, with the non degeneracy of $V = I(\theta_0)$ guaranteed by Assumption 10. It remains to show the gradient of $\bar{\Lambda}$ vanishes at $\bar{\Lambda}$, which does not come naturally as θ may lie outside of the interior of Θ . An immediate application of Proposition 4 does however yield, with probability tending to 1,

$$\theta_0 = \partial_\theta \bar{\Lambda}(\theta_0) = \lim_{n \rightarrow \infty} \frac{1}{\sqrt{n}} \left(\frac{1}{\sqrt{n}} \mathcal{S}(n, \theta_0) \right) = 0.$$

Condition (75) hence holds true.

Second condition. For any $n \in \mathbb{N}^*$ and θ in a neighbourhood of θ_0 ,

$$(76) \quad \Lambda_n(\theta_0, \theta) = \mathbf{D}_n^\top (\theta - \theta_0) + \mathbf{R}_n(\theta)^\top (\theta - \theta_0),$$

where the remainder $\mathbf{R}_n(\theta)$ is stochastically equicontinuous in the sense that, denoting by $\bar{\mathbf{R}}$ the limit in probability under $\mathbb{P}(\theta_0)$ of $n^{-1} \mathbf{R}_n$, for any $\varepsilon > 0$ and $\eta > 0$ we may find a neighbourhood K of θ_0 such that

$$(77) \quad \lim_{n \rightarrow \infty} \mathbb{P} \left[\sup_{\theta \in K} \sqrt{n} \left\| \frac{1}{n} \mathbf{R}_n(\theta) - \bar{\mathbf{R}}(\theta) \right\| > \eta \right] \leq \varepsilon.$$

The mean value theorem (or Lagrange form of the Taylor remainder) provides an expression for expansion (76) with remainder

$$(78) \quad \mathbf{R}_n(\theta)^\top = (\theta - \theta_0)^\top \partial_\theta \mathcal{S}(n, \theta^*)$$

where θ^* lies on the line between θ_0 and θ and is correctly defined on account of the convexity of Θ . A sufficient condition for the stochastic equicontinuity of the remainder is therefore that $\sqrt{n} \{n^{-1} \partial_\theta \mathcal{S}(n, \theta) - \partial_\theta \bar{\mathcal{S}}(\theta, \theta_0)\}$ be $O_{\mathbb{P}(\theta_0)}(1)$ uniformly in θ . We use again the notation of Lemma 9, for all $s, \theta \in [0, T] \times \Theta$,

$$Y_{k,ij}^{(n)}(s, \theta) = (\lambda_{k,s}^{(n)}, \partial_{\theta_i} \lambda_{k,s}^{(n)}(s, \theta), \partial_{\theta_j} \lambda_{k,s}^{(n)}, \partial_{\theta_i} \partial_{\theta_j} \lambda_{k,s}^{(n)}(\theta)).$$

From the definition of the score (30) and its derivative (31), we have some continuously differentiable function Φ such that,

$$\partial_\theta \mathcal{S}(n, \theta) = \left\{ \sum_{k=1}^K \int_0^T \Phi \left(\frac{1}{n} Y_{k,ij}^{(n)}(s, \theta) \right) dN_{k,s}^{(n)} - \int_0^T \partial_{\theta_i} \partial_{\theta_j} \lambda_{k,s}^{(n)}(\theta) ds \right\}_{1 \leq i, j \leq d},$$

so that following a re-arrangement of the terms reminiscent of the proof for Lemma 10, one has

$$\begin{aligned}
\sqrt{n}(\frac{1}{n}\partial_\theta \mathbf{S}(n, \theta) - \partial_\theta \bar{\mathbf{S}}(\theta, \theta_0))_{ij} &= \sqrt{n} \sum_{k=1}^K \int_0^T \{ \partial_{\theta_i} \partial_{\theta_j} \bar{\lambda}_k(s, \theta, \theta_0) - \partial_{\theta_i} \partial_{\theta_j} \frac{1}{n} \lambda_{k,s}^{(n)}(\theta) \} ds \\
(79) \quad &+ \sqrt{n} \sum_{k=1}^K \int_0^T \Phi(\bar{Y}_{k,ij}(s, \theta, \theta_0)) \{ \frac{1}{n} dN_{k,s}^{(n)} - h_k(s, \theta_0) \} ds \\
&+ \sqrt{n} \sum_{k=1}^K \int_0^T \Phi(\frac{1}{n} Y_{k,ij}^{(n)}(s, \theta)) - \Phi(\bar{Y}_{k,ij}(s, \theta, \theta_0)) \frac{1}{n} dN_{k,s}^{(n)}.
\end{aligned}$$

Consider the last term of (79). Re-arranging the terms again, for any $s \in [0, T]$,

$$\begin{aligned}
\sqrt{n} \{ \Phi(\frac{1}{n} Y_{k,ij}^{(n)}(s, \theta)) - \Phi(\bar{Y}_{k,ij}(s, \theta, \theta_0)) \} &= \\
&\frac{\frac{1}{n} \partial_{\theta_j} \lambda_{k,s}^{(n)}(\theta)}{\frac{1}{n} \lambda_{k,s}^{(n)}(\theta)} \sqrt{n} \{ \frac{1}{n} \partial_{\theta_i} \lambda_{k,s}^{(n)}(\theta) - \partial_{\theta_i} \bar{\lambda}_k(s, \theta, \theta_0) \} \\
&+ \frac{\partial_{\theta_i} \bar{\lambda}_k(s, \theta, \theta_0)}{\frac{1}{n} \lambda_{k,s}^{(n)}(\theta)} \sqrt{n} \{ \frac{1}{n} \partial_{\theta_j} \lambda_{k,s}^{(n)}(\theta) - \partial_{\theta_j} \bar{\lambda}_k(s, \theta, \theta_0) \} \\
(80) \quad &+ \frac{\frac{1}{n} \partial_{\theta_i} \bar{\lambda}_k(s, \theta, \theta_0) \partial_{\theta_j} \bar{\lambda}_k(s, \theta, \theta_0)}{\frac{1}{n} \lambda_{k,s}^{(n)}(\theta) \bar{\lambda}_k(s, \theta, \theta_0)} \sqrt{n} \{ \frac{1}{n} \lambda_{k,s}^{(n)}(\theta) - \bar{\lambda}_k(s, \theta, \theta_0) \} \\
&+ \frac{1}{\frac{1}{n} \lambda_{k,s}^{(n)}(\theta)} \sqrt{n} \{ \frac{1}{n} \partial_{\theta_i} \partial_{\theta_j} \lambda_{k,s}^{(n)}(s, \theta) - \partial_{\theta_i} \partial_{\theta_j} \bar{\lambda}_k(s, \theta, \theta_0) \} \\
&+ \frac{\partial_{\theta_i} \partial_{\theta_j} \bar{\lambda}_k(s, \theta, \theta_0)}{\frac{1}{n} \lambda_{k,s}^{(n)}(\theta) \bar{\lambda}_k(s, \theta, \theta_0)} \sqrt{n} \{ \frac{1}{n} \lambda_{k,s}^{(n)}(\theta) - \bar{\lambda}_k(s, \theta, \theta_0) \}.
\end{aligned}$$

All five terms on the right-hand side of equation (80) write as

$$U\left(\frac{1}{n} Y_{k,ij}^{(n)}(t, \theta)\right) V\left(\bar{Y}_{k,ij}(t, \theta)\right) \sqrt{n} \{ \partial_{\theta_i}^p \partial_{\theta_j}^m \frac{1}{n} \lambda_{k,t}^{(n)}(\theta) - \partial_{\theta_i}^p \partial_{\theta_j}^m \bar{\lambda}_k(t, \theta) \},$$

where U, V are some continuous functions, $p, m \in \{0, 1\}$, and $i, j \in \llbracket 1, d \rrbracket$. Applying Corollary 4 hence, one has that the third and last term on the right-hand side of equation (79) is uniformly $O_{\mathbb{P}}(1)$. Now, the second term on the right-hand side of (79) splits into

$$\begin{aligned}
(81) \quad &\sum_{k=1}^K \frac{1}{\sqrt{n}} \int_0^T \Phi(\bar{Y}_{ij,k}(s, \theta, \theta_0)) dM_{k,s}^{(n)} \\
&+ \int_0^T \Phi(\bar{Y}_{ij,k}(s, \theta, \theta_0)) \sqrt{n} (\frac{1}{n} \lambda_{k,s}^{(n)}(\theta_0) - \bar{\lambda}_k(s, \theta_0, \theta_0)) ds.
\end{aligned}$$

The last term of the sum (81) is uniformly $O_{\mathbb{P}}(1)$ as a direct consequence of Corollary 4. The deterministic function \bar{Y} being differentiable in t by virtue of Assumption 6 and (44), we may once again integrate by parts as in the proof of Corollary 2. Hence the first term of (81) is uniformly bounded by

$$\sup_{\theta \in \Theta} \sum_{k=1}^K |\Phi(\bar{Y}_{ij,k}(0, \theta, \theta_0))| + \int_0^T |\partial_s \bar{Y}_{ij,k}(s, \theta, \theta_0)| \Phi'(\bar{Y}_{ij,k}(s, \theta, \theta_0)) ds \frac{1}{\sqrt{n}} \sup_{t \in [0, T]} \|M_t^{(n)}\|_1,$$

which is also $O_{\mathbb{P}}(1)$ as per Lemma 7. Finally, the first and remaining integral appearing in the right-hand side of (79) is uniformly $O_{\mathbb{P}}(1)$ as a consequence of Corollary 3. Condition (77) is therefore verified.

Third condition The random vector \mathbf{D}_n in (76) verifies a central limit theorem

$$(82) \quad \frac{1}{\sqrt{n}} \mathbf{D}_n \xrightarrow[n \rightarrow \infty]{\mathcal{L}(\mathbb{P}(\theta_0))} \mathcal{N}(0, A)$$

Condition (82) is Proposition 4.

Fourth condition The MLE is consistent

$$(83) \quad \hat{\theta}_n = \theta_0 + o_{\mathbb{P}(\theta_0)}(1)$$

Condition (83) is Proposition 1.

The conditions for the constrained M-estimator master theorem are therefore satisfied. Writing

$$(84) \quad \begin{aligned} \Lambda_n &= 2(\mathcal{L}(n, \hat{\theta}_n) - \mathcal{L}(n, \hat{\theta}_n^0)) \\ &= 2(\mathcal{L}(n, \hat{\theta}_n) - \mathcal{L}(n, \theta_0)) - 2(\mathcal{L}(n, \hat{\theta}_n^0) - \mathcal{L}(n, \theta_0)), \end{aligned}$$

and applying Geyer's results [19][Theorem 4.4 and Remark p. 2004 to theorem 4.4], we obtain the weak convergence of Λ_n towards

$$(85) \quad \inf_{h \in \mathbb{H}} \{2h^T \mathbf{Z} - h^T I(\theta_0)h\} - \inf_{h \in \mathbb{H}_0} \{2h^T \mathbf{Z} - h^T I(\theta_0)h\},$$

where $\mathbf{Z} \sim \mathcal{N}(0, I(\theta_0))$. Re-arranging the terms as in Self & Liang [42][equ. (2.1) p. 606], one has

$$(86) \quad \begin{aligned} 2h^T \mathbf{Z} - h^T I(\theta_0)h &= 2h^T \mathbf{Z} - h^T \mathbf{Z} - \mathbf{Z}^T h + \mathbf{Z}^T I(\theta_0)^{-1} \mathbf{Z} - (I(\theta_0)^{-1} \mathbf{Z} - h)^T I(\theta_0) (I(\theta_0)^{-1} \mathbf{Z} - h), \\ &= \mathbf{Z}^T I(\theta_0)^{-1} \mathbf{Z} - (I(\theta_0)^{-1} \mathbf{Z} - h)^T I(\theta_0) (I(\theta_0)^{-1} \mathbf{Z} - h). \end{aligned}$$

Inserting (86) into (85) and simplifying redundant terms, the asymptotic distribution of Λ_n is the one of

$$(87) \quad \inf_{h \in \mathbb{H}_0} \{(\tilde{\mathbf{Z}} - h)^T I(\theta_0) (\tilde{\mathbf{Z}} - h)\} - \inf_{h \in \mathbb{H}} \{(\tilde{\mathbf{Z}} - h)^T I(\theta_0) (\tilde{\mathbf{Z}} - h)\},$$

where $\tilde{\mathbf{Z}} \sim \mathcal{N}(0, I(\theta_0)^{-1})$. Up to a matrix decomposition, this is the distribution of Theorem 1. Under the alternative, the weak convergence of the left-hand side of (84) towards the right-hand side of (87) is preserved, while the right-hand side of (84) is bounded from below by

$$n \inf_{\theta \in \Theta_0} \frac{1}{n} (\mathcal{L}(n, \theta) - \mathcal{L}(n, \theta_0)) = n \left(\inf_{\theta \in \Theta_0} \bar{\Lambda}(\theta, \theta_0) + o_{\mathbb{P}(\theta_0)}(1) \right)$$

as a result of Lemma 10. As $\inf_{\theta \in \Theta_0} \bar{\Lambda}(\theta, \theta_0) < 0$ since θ_0 is out of Θ_0 in the alternative, this ensures the consistency of the test.

7.3. Proof of proposition 3. From the previous discussion and Geyer [19][Theorem 4.4] still,

$$(88) \quad \sqrt{n}(\hat{\theta}_n - \theta_0) \xrightarrow[n \rightarrow \infty]{\mathcal{L}(\mathbb{P}(\theta_0))} \arg \min_{\theta \in \mathbb{H} + \theta_0} \{(\tilde{\mathbf{Z}} - (\theta - \theta_0))^T I(\theta_0) (\tilde{\mathbf{Z}} - (\theta - \theta_0))\},$$

which is the first assertion of Proposition 3, and

$$(89) \quad \sqrt{n}(\hat{\theta}_n^0 - \theta_0) \xrightarrow[n \rightarrow \infty]{\mathcal{L}(\mathbb{P}(\theta_0))} \arg \min_{\theta \in \mathbb{H}_0 + \theta_0} \{(\tilde{\mathbf{Z}} - (\theta - \theta_0))^T I(\theta_0) (\tilde{\mathbf{Z}} - (\theta - \theta_0))\},$$

where H is the Painlevé-Kuratowski limit of $\sqrt{n}(\Theta - \{\theta_0\})$ and H_0 that of $\sqrt{n}(\Theta_0 - \{\theta_0\})$. Now, for any u in some \mathbb{R}^k and $l \in \llbracket 1, k-1 \rrbracket$, we denote by $u_{\leq l}$ the sub-vector made of the first l coordinates of u . Similarly, for any $M \in \mathcal{M}_k(\mathbb{R})$ we denote by $M_{\leq l, \leq l}$ the first $l \times l$ principal submatrix of M , by $M_{\leq l, > l}$ the $l \times (k-l)$ submatrix on its right, and so on. The sparse MLE writes

$$\hat{\theta}_n^0 = \left[\hat{\theta}_{1,n}^0 \cdots \hat{\theta}_{d-p,n}^0, 0 \cdots 0 \right]^\top = \left[(\hat{\theta}_{n, \leq d-p}^0)^\top, 0 \cdots 0 \right]^\top.$$

Since $\theta_{0, \leq d-p}$ lies in the interior of its sub-parameter space, H_0 is a linear space and the arg min in (89) is attained at the critical point $h^* \in \mathbb{R}^{d-p}$ of

$$[h_1, \cdots, h_{d-p}]^\top \mapsto (\mathbf{Z}^\top - [h_1 \cdots h_{d-p}, 0 \cdots 0])I(\theta_0)(\mathbf{Z} - [h_1 \cdots h_{d-p}, 0 \cdots 0]^\top),$$

which expresses as

$$(90) \quad h^* = Z_{\leq d-p} - (I(\theta_0)_{\leq d-p, \leq d-p})^{-1} I(\theta_0)_{\leq d-p, > d-p} Z_{> d-p}.$$

After some elementary manipulation of bloc-matrix inverses, the variance of the Gaussian variable on the right-hand side of (90) simplifies as the inverse of $I(\theta_0)_{\leq d-p, \leq d-p}$ (see Appendix A.1). One finds

$$\sqrt{n}(\hat{\theta}_{n, \leq d-p}^0 - \theta_{0, \leq d-p}) \xrightarrow[n \rightarrow \infty]{\mathcal{L}(\mathbb{P}(\theta_0))} \mathcal{N}(0, (I(\theta_0)_{\leq d-p, \leq d-p})^{-1}).$$

This is the second and final assertion of Proposition (3).

8. ACKNOWLEDGEMENTS

The author would like to thank T. Deschatre, P. Gruet, and M. Hoffmann for helpful discussions and their fruitful input during the preparation of this work; and E. Cognéville and F. Trieu for the pre-processing of the German intraday power market order book.

A. APPENDIX

A.1. On the variance of the sparse MLE. Lemma 13 is part of the routine technicalities of likelihood-based model selection (see for instance Van der Vaart [50][Problem 5 p. 241]). Since its short proof is rarely stated in full detail we recall it here for the sake of completeness. The sub-vector notation is the one of section 7.3 and we suppose, without loss of generality, that the null adjacency coefficients constitute the last p coordinates of the true parameter θ_0 .

Lemma 13. *The asymptotic variance under $\mathbb{P}(\theta_0)$ of the first $d-p$ coordinates of $\sqrt{n}(\hat{\theta}_n^0 - \theta_0)$ is the inverse of $I(\theta_0)_{\leq d-p, \leq d-p}$, where $I(\theta_0)$ is the $d \times d$ asymptotic information matrix.*

Proof. Write

$$I(\theta_0) = \begin{bmatrix} A & B \\ B^\top & C \end{bmatrix}.$$

where A is a $(d-p) \times (d-p)$ submatrix. Then, using Schur's notation, one has

$$I(\theta_0)^{-1} = \begin{bmatrix} (I/D)^{-1} & (I/D)^{-1} B D^{-1} \\ D^{-1} B^\top (I/D)^{-1} & (I/A)^{-1} \end{bmatrix},$$

where the Schur complements

$$\begin{aligned} (I/D)^{-1} &= A - B D^{-1} B^\top, \\ (I/A)^{-1} &= D - B^\top A^{-1} B, \end{aligned}$$

verify

$$(91) \quad (I/D)^{-1} = A^{-1} + A^{-1}B(I/A)^{-1}B^T A^{-1},$$

$$(92) \quad (I/A)^{-1} = D^{-1} + D^{-1}B^T(I/D)^{-1}BD^{-1}.$$

From the proof of proposition 3 the asymptotic variance of

$$\sqrt{n}(\hat{\theta}_n^0 - \theta_{0, \leq d-p})$$

under $\mathbb{P}(\theta_0)$ is the variance of

$$\mathbf{Z}_{\leq d-p} - A^{-1}B^T \mathbf{Z}_{> d-p}.$$

whre $\mathbf{Z} \sim \mathcal{N}(0, I(\theta_0)^{-1})$. Following a straightforward simplification,

$$\text{Var}(\mathbf{Z}_{\leq d-p} - A^{-1}B^T \mathbf{Z}_{> d-p}) = (I/D)^{-1} - A^{-1}B(I/A)^{-1}B^T A^{-1}$$

which reformulates thanks to (91) as

$$A^{-1} = (I(\theta_0)_{\leq d-p, \leq d-p})^{-1}$$

□

A.2. On the Chi-Bar Distribution. The existence of the $\bar{\chi}^2$ distribution as a mixture of χ^2 distributions for regular enough models is a traditional result in non-standard likelihood theory. Proposition 1 recalls the correspondence between the Fisher information and the mixture weights of the chi-bar distribution for models parameterised on a closed orthant. This result was originally proposed by Kudo [33][Theorem 3.1 p. 414] and can also be regarded as a special case of Shapiro [43][Theorem 3.1 p. 138]. We provide a short alternative proof to Kudo's [33] to ensure the present paper remains self-contained. Let $d \in \mathbb{N}^*$ and define the orthant

$$(93) \quad C = \{\theta = (\theta_i) \in \mathbb{R}^d \mid \theta_i \geq 0 \text{ for all } i \in \llbracket 1, d \rrbracket\}.$$

For any $p \in \llbracket 1, d \rrbracket$ we denote by \mathcal{P}_k^p the set of all combinations of size k of $\llbracket 1, p \rrbracket$, or, up to a bijection,

$$\mathcal{P}_k^p = \{(i_1, \dots, i_k) \mid 1 \leq i_1 < \dots < i_k \leq p\},$$

and for any $I_k \in \mathcal{P}_k^p$ we abuse the notation to denote

$${}^c I_k = (j_1, \dots, j_{p-k}), \quad j_1 < \dots < j_{p-k}, \quad j_l \in \llbracket 1, p \rrbracket \setminus I_k.$$

For any $p \in \llbracket 0, p \rrbracket$, define also the sparse sub-spaces

$$C_p = \{\theta \in C \mid \text{there are exactly } p \text{ null coefficients in } \theta\},$$

and

$$C^{I_k} = \{\theta \in C \mid \theta_i = 0 \iff i \in I_k\},$$

so that for any $p \geq 1$, C_p is the union of the disjoint sets C^{I_k} with $I_k \in \mathcal{P}_k^p$, and $C_0 = C$. Finally we let A be a positive definite $d \times d$ real matrix. Picking $p \in \llbracket 1, d \rrbracket$, we let

$$m = [0 \quad \dots \quad 0 \quad m_{p+1} \quad \dots \quad m_d]^T \in C^{(1,2,\dots,p)} \subset C_p,$$

and $X \sim \mathcal{N}(m, A^{-1})$, and we define the quadratic form

$$q_X^A: z \in \mathbb{R}^d \mapsto (X - z)^T A (X - z).$$

Definition 11 (Chi-bar distribution). *The $\bar{\chi}_A^2(p)$ distribution is the law of*

$$(94) \quad \ell^A(X) = \min_{z \in C_p} q_X^A(z) - \min_{z \in C} q_X^A(z).$$

In the rest of this article we have omitted the variance subscript from the $\bar{\chi}^2$ notation to match conventions in force in the literature. Do note that regardless of this practice, a $\bar{\chi}^2$ distribution cannot be characterised by its degrees of freedom alone.

Proposition 1. *The Chi-bar law writes as a mixture of χ^2 distributions*

$$\bar{\chi}_{\mathcal{A}}^2(p) = \sum_{i=1}^p \omega_i^{\mathcal{A}} \chi^2(p-i),$$

where, defining $F_{I_k} \sim N(0, (\mathcal{A}_{I_k}^{-1})^{-1})$ and $G_{I_k} \sim N(0, (\mathcal{A}_{cI_k})^{-1})$ the mixing probabilities are given by

$$\omega_k^{\mathcal{A}} = \sum_{I_k \in \mathcal{P}_k^p} \mathbb{P}[F_{I_k} \in \mathbb{R}^{-k}] \mathbb{P}[G_{I_k} \in \mathbb{R}^{+p-k} \times \mathbb{R}^{K-p}].$$

Proof. In the rest of the proof, denote

$$\tilde{z} = \arg \min_{z \in C} (z - X)^T \mathcal{A} (z - X) \quad \text{and} \quad \tilde{z}^p = \arg \min_{z \in C_p} (z - X)^T \mathcal{A} (z - X).$$

Let us first clarify the distribution of the first term on the right of (94). By Karush-Kuhn-Tucker (KKT), there exist some random coefficients $\lambda^T \in \mathbb{R}^{+p}$ such that $\lambda_i \tilde{z}_i^p = 0$ for $i = 1 \dots k$, and

$$(95) \quad (\tilde{z}^p - X)^T \mathcal{A} = [\lambda_1 \quad \dots \quad \lambda_p \quad 0 \quad \dots \quad 0] = [\lambda \quad 0 \quad \dots \quad 0]$$

which given that $\tilde{z}_1^p = \tilde{z}_p^p = 0$ as $\tilde{z}^p \in C_p$ yields

$$(96) \quad \lambda = ((\mathcal{A}^{-1})_{\leq p, \leq p})^{-1} X_{\leq p} \sim N(0, (\mathcal{A}_{\leq p, \leq p}^{-1})^{-1}).$$

Plugging (95) back into $q_X^{\mathcal{A}}$,

$$\min_{z \in C_p} q_X^{\mathcal{A}}(z) = (\tilde{z}^p - X)^T \mathcal{A} (\tilde{z}^p - X) = [\lambda \quad 0 \quad \dots \quad 0] (\tilde{z}^p - X) = \lambda^T (\mathcal{A}^{-1})_{\leq p, \leq p} \lambda$$

which together with (96) shows that $\min_{z \in C_p} q_X^{\mathcal{A}}(z)$ follows a $\chi^2(p)$ distribution and writes as

$$\min_{z \in C_p} q_X^{\mathcal{A}}(z) = \sum_{k=1}^p \frac{X_k^2}{\mathcal{A}_{kk}^{-1}}.$$

We may actually proceed in a similar fashion for the second term of (96). Let $k \in \llbracket 1, p \rrbracket$. The event $\{\tilde{z} \in C_k\}$ decomposes into the mutually exclusive $\{\tilde{z} \in C^{I_k}\}$, $I_k \in \mathcal{P}_k^p$, upon each of which the behaviour of $\ell_X^{\mathcal{A}}$ is the same. We may assume $I_k = \llbracket 1, k \rrbracket$ without loss of generality. On the event $\{\tilde{z} \in C^{\llbracket 1, k \rrbracket}\}$ then, \tilde{z} coincides with the solution of the minimisation program of $q_X^{\mathcal{A}}$ over $C^{\llbracket 1, k \rrbracket}$, which is strongly dual, and verifies the KKT conditions (95) with p replaced by k . In particular,

$$(97) \quad Y^{(k)} = ((\mathcal{A}^{-1})_{\leq k, \leq k})^{-1} X_{\leq k} = -[\lambda_1 \dots \lambda_k]^T \in \mathbb{R}^{-k},$$

and, re-inserting the preceding expression for λ back into relation (95),

$$(98) \quad Z^{(k)} = X_{>k} - (\mathcal{A}^{-1})_{>k, \leq k} Y^{(k)} = \tilde{z}_{>k} \in \mathbb{R}^{+*p-k} \times \mathbb{R}^{K-p}.$$

From the same argument as for the first term, it follows from (97) that conditionally on $\{\tilde{z} \in C^{I_k}\}$,

$$\min_{z \in C} q_X^A(z) = \lambda^T (\mathcal{A}^{-1})_{\leq k \leq k} \lambda = \sum_{i \in I_k} \frac{X_i^2}{\mathcal{A}_{ii}^{-1}} \quad \text{and} \quad \ell^A(X) = \sum_{i \notin I_k} \frac{X_i^2}{\mathcal{A}_{ii}^{-1}} \sim \chi^2(p - i).$$

Note furthermore that relation (97) implies λ_i taking value 0 is a negligible event and the λ 's can be taken almost surely strictly positive. Reciprocally, when conditions (97) and (98) hold, that is, $Y^{(k)} < 0$ and $Z_{\leq p-k}^{(k)} > 0$ coordinate-wise, the pair $(x, \lambda) \in C^{I_k} \times \mathbb{R}^{+*k}$ defined by $\lambda = -(\mathcal{A}_{\leq k \leq k}^{-1})^{-1} X_{\leq k}$ and $\tilde{z} = [0 \ \cdots \ 0 \ Z^{(k)}]$ satisfy the KKT conditions for the minimisation problem of q_X^A over C . Its solution \tilde{z} then lies in C^{I_k} and therefore in C_k . We have shown the event $\{\tilde{z} \in C^{\llbracket 1, k \rrbracket}\}$ can be rewritten as $\{Y^{(k)} < 0, Z_{\leq p-k}^{(k)} > 0\}$. Remarking furthermore from simple matrix manipulations and Schur's formulas that

$$\begin{aligned} \text{Cov}(Y^{(k)}, Y^{(k)}) &= ((\mathcal{A}^{-1})_{\leq k \leq k})^{-1} \\ \text{Cov}(Z^{(k)}, Y^{(k)}) &= (\mathcal{A}^{-1})_{> k \leq k} ((\mathcal{A}^{-1})_{\leq k \leq k})^{-1} - (\mathcal{A}^{-1})_{> k \leq k} \text{Cov}(Y^{(k)}, Y^{(k)}) = 0 \\ \text{Cov}(Z^{(k)}, Z^{(k)}) &= (\mathcal{A}^{-1})_{> k > k} - (\mathcal{A}^{-1})_{> k \leq k} ((\mathcal{A}^{-1})_{\leq k \leq k})^{-1} (\mathcal{A}^{-1})_{\leq k > k} = (\mathcal{A}_{> k, > k})^{-1}, \end{aligned}$$

one observes $Y^{(k)}$ and $Z^{(k)}$ to be independent. Up to a re-arrangement of the coordinate of X , the same arguments hold for any other $I_k \in \mathcal{P}_k^n$. Summing over all possible configurations for I_k , this gives the announced expression for the mixture weight ω_k^A . There remains only the situation of \tilde{z} falling in the interior of C , in which case the second term of (94) vanishes and the distribution of ℓ_X^A is a $\chi^2(p)$ law, as would traditionally be observed in absence of positivity constraints. This happens with probability $\mathbb{P}[X_{\leq p} \in \mathbb{R}^{+p}]$. □

Remark 13. *In all generality, the expression for the weights have no closed form. In the case $K = p = 2$ however, $\mathbb{P}[F \in \mathbb{R}^- \times \mathbb{R}^-] = \mathbb{P}[H \in \mathcal{A}^{-1/2} \mathbb{R}^- \times \mathbb{R}^-]$ where $H \sim \mathcal{N}(0, I_2)$ and re-expresses as the portion of the unit ball intersecting with $\mathcal{A}^{-1/2} \mathbb{R}^- \times \mathbb{R}^-$, which is given by the angle between $\mathcal{A}^{-1/2} [-1 \ 0]$ and $\mathcal{A}^{-1/2} [0 \ -1]$, that is, foregoing a few matrix manipulation,*

$$\omega_2 = \frac{1}{2\pi} \cos^{-1}(\mathcal{A}_{12} / \sqrt{\mathcal{A}_{11} \mathcal{A}_{22}}) = \frac{1}{2\pi} \left\{ \pi - \cos^{-1}(\mathcal{A}_{12}^{-1} / \sqrt{\mathcal{A}_{11}^{-1} \mathcal{A}_{22}^{-1}}) \right\} = \frac{1}{2} - \omega_0.$$

This is example 7 from Self & Liang [42].

REFERENCES

- [1] P. Andersen, O. Borgan, R. Gill, and N. Keiding. Statistical models based on counting processes. Springer Series in statistics, 1992.
- [2] D. Andrews. Estimation when a parameter is on a boundary. Econometrica, 67(6), 1999.
- [3] D. Andrews. Testing when a parameter is on the boundary of the maintained hypothesis. Econometrica, 69(3):683–734, 2001.
- [4] D. W. K. Andrews and W. Ploberger. Admissibility of the likelihood ratio test when a nuisance parameter is present only under the alternative. The Annals of Statistics, 23(5):1609–1629, 1995.
- [5] E. Bacry, M. Bompain, S. Gaïffas, and J.F. Muzy. Sparse and low-rank multivariate Hawkes processes. J. Mach. Learn. Res., 21(1), jan 2020.
- [6] E. Bacry, S. Delattre, M. Hoffmann, and F. Muzy. Some limit theorems for Hawkes processes and application to financial statistics. Stochastic Processes and their Applications, 123(7):2475–2499, 2013.
- [7] A Bonnet, M. Martinez Herrera, and M Sangnier. Inference of multivariate exponential Hawkes processes with inhibition and application to neuronal activity. Statistics and Computing, 33(4), 2023.

- [8] P. Brémaud. *Point Process Calculus in Time and Space*, volume 98 of *Probability Theory and Stochastic Modelling*. Springer, 2020.
- [9] F. Chen and P. Hall. Inference for a nonstationary self-exciting point process with an application in ultra-high frequency financial data modeling. *Journal of Applied Probability*, 50(4):1006–1024, 2013.
- [10] F. Chen and W. Tan. Marked self exciting point process modelling of information propagation on twitter. *The Annals of Applied Statistics*, 12(4):2175–2196, 2018.
- [11] J. Chen, Michael P. Clements, and A. Urquhart. Modeling price and variance jump clustering using the marked Hawkes process. *Journal of Financial Econometrics*, 2023.
- [12] W-H Chiang, X. Liu, and G. Mohler. Hawkes process modeling of COVID-19 with mobility leading indicators and spatial covariates. *International Journal of Forecasting*, 38(2):505–520, 2022.
- [13] S. Dachian and Y. A. Kutoyants. Hypotheses testing: Poisson versus self-exciting. *Scandinavian Journal of Statistics*, 33(2):391–408, 2006.
- [14] Daley and Vere-Jones. *An Introduction to the Theory of Point Processes: elementary theory and methods*. Probability and its Applications. Springer, second edition, 2002.
- [15] S. Delattre, N. Fournier, and M. Hoffmann. Hawkes processes on large networks. *The Annals of Applied Probability*, 26(1):216–261, 2016.
- [16] T. Deschatre and P. Gruet. Electricity intraday price modelling with marked Hawkes processes. *Applied Mathematical Finance*, pages 227–260, 2023.
- [17] V. Didelez. Graphical models for marked point processes based on local independence. *Journal of the Royal Statistical Society. Series B (Statistical Methodology)*, 70(1):245–264, 2008.
- [18] P. Embrechts, T. Liniger, and L. Lin. Multivariate Hawkes processes: an application to financial data. *Journal of Applied Probability*, 48(A), 2011.
- [19] C. Geyer. On the Asymptotics of Constrained M-Estimation. *The Annals of Statistics*, 22(4):1993–2010, 1994.
- [20] M. Goda. Sparse estimation for generalized exponential marked Hawkes process. *Stat Inference Stoch Process*, 26, 2023.
- [21] G. Gripenberg, S. O. Londen, and O. Staffans. *Volterra Integral and Functional Equations*. Encyclopedia of Mathematics and its Applications. Cambridge University Press, 1990.
- [22] Bruce E. Hansen. Inference when a nuisance parameter is not identified under the null hypothesis. *Econometrica*, 64(2):413–430, 1996.
- [23] N. R. Hansen, P. Reynaud-Bouret, and V. Rivoirard. Lasso and probabilistic inequalities for multivariate point processes. *Bernoulli*, 21(1):83–143, 2015.
- [24] S. J. Hardiman, N. Bercot, and J-P. Bouchaud. Critical reflexivity in financial markets: a Hawkes process analysis. *The European Physical Journal B*, 86(10), 2013.
- [25] S. J. Hardiman and J-P. Bouchaud. Branching-ratio approximation for the self-exciting Hawkes process. *Phys. Rev. E*, 90:062807, 2014.
- [26] A. Hawkes and D. Oakes. A cluster process representation of a self-exciting process. *Journal of Applied Probability*, 11(3):493–503, 1974.
- [27] J. Jacod. Multivariate point processes: predictable projection, radon-nikodym derivatives, representation of martingales. *Zeitschrift für Wahrscheinlichkeitstheorie und Verwandte Gebiete*, 31:235—253, 1975.
- [28] J. Jacod and A.N. Shiryaev. *Limit theorems for stochastic processes*. volume 288 of *Grundlehren der Mathematischen Wissenschaften*. Springer-Verlag, 1987.
- [29] T. Jaisson and M. Rosenbaum. Limit theorems for nearly unstable Hawkes processes. *The Annals of Applied Probability*, 25(2), 2015.
- [30] P. Jusselin and M. Rosenbaum. No-arbitrage implies power-law market impact and rough volatility, 2020.
- [31] A. Kramer and R. Kiesel. Exogenous factors for order arrivals on the intraday electricity market. *Energy Economics*, 97, 2021.
- [32] V. Krishna. *Auction Theory*. 01 2002.
- [33] A. Kudo. A multivariate analogue of the one-sided test. *Biometrika*, 50(3):403—418, 1963.
- [34] K. Lee and B. Ki Seo. Marked Hawkes process modeling of price dynamics and volatility estimation. *Journal of Empirical Finance*, 40:174–200, 2017.
- [35] Y. Ogata. The asymptotic behaviour of maximum likelihood estimators for stationary point processes. *Ann Inst Stat Math*, 30:243–261, 1978.
- [36] Y. Ogata. Statistical models for earthquake occurrences and residual analysis for point processes. *Journal of the American Statistical Association*, 83(401):9–27, 1988.
- [37] D. Pollard. *Convergence of stochastic processes*. Springer Series in Statistics. Springer Verlag, 1984.

- [38] R. Rebolledo. Central limit theorems for local martingales. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete*, 51:269–286, 1980.
- [39] P. Reynaud-Bouret, V. Rivoirard, and C. Tuleau-Malot. Inference of functional connectivity in neurosciences via Hawkes processes. In *2013 IEEE Global Conference on Signal and Information Processing*, 2013.
- [40] C. Ritz and Ib M. Skovgaard. Likelihood ratio tests in curved exponential families with nuisance parameters present only under the alternative. *Biometrika*, 92(3):507–517, 09 2005.
- [41] F. Salehi, W. Trouleau, M. Grossglauser, and P. Thiran. Learning Hawkes processes from a handful of events. In *Advances in Neural Information Processing Systems*, volume 32. Curran Associates, Inc., 2019.
- [42] S. Self and K-Y. Liang. Asymptotic properties of maximum likelihood estimators and likelihood ratio tests under nonstandard conditions. *Journal of the American Statistical Association*, 82(398):605–610, 1987.
- [43] A. Shapiro. Asymptotic distribution of test statistics in the analysis of moment structures under inequality constraints. *Biometrika*, 72(1):133–144, 1985.
- [44] G. Shmueli, R. P. Russo, and W. Jank. The barista: A model for bid arrivals in online auctions. *The Annals of Applied Statistics*, 1(2), 2007.
- [45] M. Silvapulle and P.K. Sen. *Constrained statistical inference: Inequality, order and shape restrictions*. John Wiley & Sons, 1 edition, 2005.
- [46] D. Sulem, V. Rivoirard, and J. Rousseau. Bayesian estimation of nonlinear Hawkes processes. *Bernoulli*, 30(2):1257–1286, 2024.
- [47] E. Susko. Likelihood ratio tests with boundary constraints using data-dependent degrees of freedom. *Biometrika*, 100(4), 08 2013.
- [48] A. Takeuchi. Integration by parts formulas for marked Hawkes processes. *Statistics & Probability Letters*, 145:229–237, 2019.
- [49] S. van de Geer. Exponential Inequalities for Martingales, with Application to Maximum Likelihood Estimation for Counting Processes. *The Annals of Statistics*, 23(5):1779 – 1801, 1995.
- [50] A Van den Vaart. *Asymptotic statistics*. Cambridge Series in statistical and probabilistic mathematics, 1998.
- [51] H. Xu, M. Farajtabar, and H. Zha. Learning granger causality for Hawkes processes. volume 48 of *Proceedings of Machine Learning Research*, New York, New York, USA, 20–22 Jun 2016. PMLR.
- [52] K Zhou, H. Zha, and L. Song. Learning triggering kernels for multi-dimensional Hawkes processes. volume 28 of *Proceedings of Machine Learning Research*, Atlanta, Georgia, USA, 17–19 Jun 2013. PMLR.