Deciphering Solar Cycle Influence on Long-Term Orbital Deterioration of Space Debris in LEO and MEO orbits

Ayisha M Ashruf¹, Ankush Bhaskar¹, and Tarun Kumar Pant¹

¹Space Physics Laboratory, Vikram Sarabhai Space Centre, Thiruvananthapuram, Kerala, 695022, India

Key Points:

- first statistical study to show the long-term impact of solar cycles on the orbital decay of space debris even at MEO orbits
- rapid decay observed near the peak of each solar cycle bounded by $\sim 70\%$ from peak amplitude
- ionospheric impact can take over a year to manifest at higher altitudes

Corresponding author: Ayisha M. Ashruf, ayishamashruf@gmail.com

Abstract

The rapid increase in the number of space debris represents a substantial threat to the sustained viability of space operations. Monitoring these debris is critical for space situational awareness. Therefore, it has become of immense importance to understand the impact of solar activity on the space debris orbit. This study examines the effect of solar cycles, specifically Solar Cycles 22, 23, and 24, on the orbital decay of space debris. Utilizing TLE data for 95 objects in LEO and MEO dating back to the 1960s, we examined the orbital decay rates of these objects across the three solar cycles. Our analysis reveals a significant correlation between orbital decay and various indices serving as proxies for solar and geomagnetic activity. Notably, Solar Cycle 22 exhibited the highest decay rates, while Solar Cycle 24 showed the lowest. Moreover, we observed a similar impact at altitudes exceeding 2000 km, albeit with a time lag of over 14 months compared to the shorter lag observed at LEO orbits during peak solar activity. These findings underscore the crucial relationship between solar activity and its enduring influence on space debris.

Plain Language Summary

Space is becoming increasingly cluttered with debris, posing a challenge for longterm space activities. Our study investigates how solar activity, specifically during Solar Cycles 22, 23, and 24, impacts the orbital decay of space debris. We analyzed data for 95 space debris in LEO and MEO, all originating from the 1960s era. Our findings are intriguing: when the sun is more active, as observed in Solar Cycle 22, space debris experiences faster decay. Conversely, during Solar Cycle 24, characterized by lower solar activity, the decay rate was slower. Surprisingly, this influence was also evident at altitudes exceeding 2000 km, although it took more than a year for the effect to manifest compared to the shorter lag observed in orbits closer to Earth during peak solar activity which is intriguing. These findings highlight how the sun's behavior affects space junk on longer time scales.

1 Introduction

Space debris, also referred to as orbital debris, encompasses non-functional satellites, discarded rocket stages, and other artificial objects orbiting the Earth, and has the potential to cause damage to active spacecraft Graham et al. (1999); Li et al. (2002); Kleinig et al. (2022); Smith et al. (2020); Hayes & Caverly (2023). As the space industry expands rapidly, the volume of space debris also increases, presenting considerable hazards to functioning satellites, ongoing space missions, and astronaut safety. A pressing issue arises from the risk of cascading collisions, commonly termed the Kessler Syndrome Kessler et al. (2010), where a single collision can generate substantial additional debris, triggering a domino effect of subsequent collisions. Various factors influence the deterioration and decay of space debris, such as solar activity, geomagnetic storms, weathering of the debris, atmospheric compositional changes, and collisions, with the solar cycle's impact emerging as a significant contributor Nwankwo (2018); Klinkrad (2006); Walterscheid (1989). A recent event that sparked concerns in the scientific and space-tech sectors was the loss of 39 Starlink satellites, which was linked to mild geomagnetic disruptions Dang et al. (2022); Kataoka et al. (2022).

Furthermore, during severe space weather events, such as the historic Carrington-type extreme geomagnetic storms, satellites can experience significant drag, causing them to descend to lower altitudes and potentially reducing their operational lifespan Oliveira et al. (2020); Tsurutani et al. (2003); Hayakawa et al. (2022); Bruinsma et al. (2023). Typically, approximately every 11 years, the Sun undergoes active and quiet phases, constituting a solar cycle, resulting in notable variations in the emitted flux of electromagnetic and corpuscular radiation Lean (1987); Hathaway (2015); Richardson & Cane (2012); Gopalswamy (2006). This variation in solar activity drives changes in the Earth's magnetosphere and upper

atmosphere, specifically in density, temperature, and atmospheric scale height, which, in turn, impacts the orbital drag of satellites and space debris Harris & Priester (1962); Clette et al. (2015); Weng et al. (2020); Crisp et al. (2021).

Recent studies have delved into the effects of space weather, particularly solar transient events, on satellite drag. However, the short and long-term influence of these events on space debris remains a relatively uncharted area. As the significance of space situational awareness and mitigation measures continues to rise, there is an increasing demand for accurate monitoring of space debris and reliable orbit predictions. Moreover, considering the lifetime of space debris, which could span many years to decades, it becomes crucial to evaluate and quantify the impact of long-term variations in solar activity on space debris. This motivates the present study, which investigates the influence of solar cycles on the orbital decay of space debris in the terrestrial upper atmosphere, specifically focusing on Solar Cycles 22, 23, and 24. The research unequivocally demonstrates the influence of solar activity cycles on the degradation rate of space debris within Low-Earth Orbit (LEO) and Medium-Earth Orbit (MEO). The analysis covers 95 space debris objects that have been in orbit since the 1960s, and for which TLE data is available throughout the most recent three solar cycles, ensuring adequate time resolution.

2 Data and Methodology

The sunspot number (SN) constitutes a historical time series spanning from 1700 to the present, capturing the 11-year cyclic and long-term variations in solar activity and, consequently, space weather. Although contemporary observations offer more refined parameters for understanding space-weather impacts, the sunspot number represents the earliest direct record of solar activity. It serves as an essential link connecting past and present solar behavior Clette et al. (2014); Clette & Lefèvre (2016); Jayalekshmi et al. (2022).

Along with the SN data, the F10.7 index is also known to be a fairly good representative of solar activity Chen et al. (2011). The geomagnetic activity is represented by the Dst index, which is derived from ground-based magnetic observatories located in low-latitude regions that measure the intensity of the globally symmetrical equatorial electrojet or the ring current. Due to its ability to efficiently and consistently track the onset and evolution of magnetic storms, it stands out as one of the extensively used indices in geomagnetism and space physics Rostoker (1972); Zhang et al. (2023). The AL index describes the peak intensity of the westward electrojet current in the auroral zone and has been widely used to represent geomagnetic substorms Huang & Reeves (2005). Solar Cycle 22 spanned from September 1986 to August 1996, succeeded by Solar Cycle 23, which lasted until December 2008, and Solar Cycle 24, persisting until December 2019. The data for these indices are retrieved from the websites gfz-potsdam.de and wdc.kugi.kyoto-u.ac.jp respectively.

The orbital parameters of satellites and space debris are commonly summarized in Two-Line Elements or TLEs, which consist of a maximum of 69 alphanumeric characters. A detailed description of the TLE format can be found in Vallado & Cefola (2012). For this study, debris originating in the 1960s were selected, possessing TLE data with sufficient time resolution during Solar Cycles 22, 23, and 24. Notably, these objects continue to be in orbit as of June 2023. A total of 95 space debris objects from the 1960s were identified through Space-track (space-track.org), and the same website was utilized to obtain TLE data covering the period from September 1, 1986, to June 30, 2023. Subsequently, these objects were categorized into two groups based on their altitudes: out of the 95, 60 were LEO objects with altitudes up to 1000 km, and the remaining 35 were MEO objects with altitudes exceeding 1000 km.

The TLE elements of individual space debris were utilized to create a database of the orbital parameters of the objects at different epochs. This database was subsequently employed for further analyses. The semi-major axis 'a' of the orbit of each space debris at a given epoch is obtained from mean motion using the following equation,

$$a = \left(\frac{GM}{\left(\frac{2\pi n}{86400}\right)^2}\right)^{\frac{1}{3}}$$
(1)

where M is the mass of the Earth, G is the universal gravitational constant and n is the mean motion in revolutions per day. Following this, we plotted the rate of decay in meters per hour and SN as functions of the epoch. To handle outliers within the data points, we utilize z-scores, which provide information about the number of standard deviations a value deviates from the mean in a given distribution. For both the LEO and MEO objects, we exclude data points with a semi-major axis z-score exceeding 3 from the analysis.

To gain a more comprehensive and quantitative understanding of the influence of the solar cycle, we proceeded to compute the average rate of decay for all objects throughout the three solar cycles. The relevant data for these objects can be found in Table 1.

Following this, we investigated the relationship between initial altitude and the rate of decay near the peak of each solar cycle. Subsequently, we generated histograms to visually represent the mean decay rate for both LEO and MEO objects during different solar cycles. Additionally, we examined the correlation between orbital decay and indices such as SN, F10.7, Dst, and AL.

3 Observations

Figure 1 presents sample plots depicting the correlation between altitude and Sunspot number over the three solar cycles for two objects in LEO and two objects in MEO, serving as representative examples. In this context, 'apparent altitude' $(a - R_e)$ denotes a value derived by subtracting the Mean Earth Radius from the Mean Semi-Major Axes obtained from the TLE data. The decay profiles of the sample objects over the specified period are presented in the top two panels on the left and right. The color-blocked regions in these panels correspond to solar cycles 22, 23, and 24, respectively. Concurrently, figure panel (c) illustrates the cumulative sunspot number, figure panel (d) showcases the associated sunspot number, and the bottom panel depicts the F10.7 flux. Within figure panel (d), the blue curve represents the Gaussian curve fitted to the sunspot numbers. Cyan dashed lines are incorporated to facilitate the distinction between different solar cycles, improving visual clarity. The fitting of Gaussian curves to sunspot numbers in each solar cycle serves to estimate the percentage of the peak value at which sudden changes in decay rates occur. Additionally, the data points were fitted to both a log-normal distribution and a 12-month running mean. While the log-normal fit closely resembled the Gaussian fit presented here, the running mean proved unsuitable.

The vertical red dashed lines serve as markers for areas characterized by nearly linear and rapid orbit decay during each solar cycle. To position these red lines, the rate of decay is calculated for the entire dataset using two adjacent points. The points where a sharp change in the decay rate occurs is identified as the starting and ending points for slope calculations. Data points within these marked regions are then utilized in the computation of slopes. Regarding solar cycle 22, the red dashed lines intersect the Gaussian curve at 63% of the peak height during the ascending phase and 71% during the descending phase. For solar cycle 23, the red dashed lines intersect the Gaussian curve at 68% percent during the ascending phase and 75% during the descending phase. Within solar cycle 24, the red dashed lines intersect the Gaussian curve at 67% percent during both the ascending and descending phases. This suggests that beyond the threshold value of SN (~ 67%), there is a noticeable onset of rapid orbital decay for all cycles. Several significant observations can be




Figure 1. Apparent altitude, cumulative sunspot number, sunspot number, and F10.7 adjusted (s.f.u.) variation over the three solar cycles for two samples of LEO and MEO objects

gleaned from this figure. Notably, Solar Cycle 22 exhibits the highest level of solar activity, followed by Solar Cycle 23, with Solar Cycle 24 being the least active. As anticipated, the slopes of the plots are steeper during the peak of each solar cycle, reflecting an increased decay rate. The figures depict that the decay rate is more rapid during Solar Cycles 22 and 23, as indicated by the steeper slopes, a trend directly attributable to the higher solar activity levels represented by the sunspot numbers. Additionally, the decay rate becomes more gradual, almost leveling off as it approaches the solar minima.

Upon examining the cumulative sunspot number depicted in panel (c) figure 1, it becomes evident that it undergoes a rapid ascent during the peak of solar cycles. Subsequently, it reaches a saturation point and maintains an almost flat trend during the minima phase. Following this, there is another rapid rise as it enters the active phase of the next solar cycle. During periods of high solar activity, there is a rise in X-ray and ultraviolet radiation emitted by the Sun, leading to the expansion of the thermosphere as it absorbs this heightened energy influx. As the Sun approaches the solar minimum, the thermosphere undergoes a cooling and contraction phase due to the diminishing intensity of X-ray and ultraviolet radiation, as indicated by Mlynczak et al. (2018). Similarly, the orbital decay plots exhibit a comparable pattern, with rapid decay during the peak, followed by a period of flatter slope during periods of lower solar activity. Once again, a change in the rate of decay is noticeable during the peak of the subsequent cycle. Heightened solar activity leads to an increased scale height, causing a proportional rise in density at the orbital altitude. As the drag force experienced by the orbiting body depends on atmospheric density, this, in turn, accelerates orbital decay. The acceleration experienced by the object due to drag force can be expressed as,

$$a = -\frac{1}{2} \frac{C_D A}{m} \rho v^2 \tag{2}$$

where m is the mass of the orbiting body, C_D is the drag coefficient, A is the crosssectional area, ρ is the atmospheric density and v is the relative velocity of the body with respect to the atmosphere.

Another significant perturbative force that can affect an orbiting body is solar radiation pressure, arising from the exchange of momentum between photons and the object in orbit. The transferred momentum can be mathematically represented as,

$$\Delta p = \alpha \frac{\Delta U}{c} \tag{3}$$

where, ΔU is the change in energy, c is the speed of light and the coefficient α ranges from 1 to 2, varying from a perfect absorber to a perfect reflector.

The radiation pressure can then be expressed as,

$$P = \frac{1}{A} \frac{\Delta p}{\Delta t} \tag{4}$$

$$P = \frac{1}{A\Delta t} \left(\alpha \frac{\Delta U}{c} \right) \tag{5}$$

where, A is the area and Δt is the time interval. The equation can then be rearranged as

$$P = \frac{\alpha}{c} \left(\frac{\Delta U}{A \Delta t} \right) \tag{6}$$

where the term in the bracket is the average flux S. So the equation can be re-written as,

$$P = \alpha \frac{F}{c} \tag{7}$$

We know that S can be computed using the following equation,

$$S = \frac{L}{4\pi r^2} \tag{8}$$

where L is the luminosity of the Sun and r is the distance between the object and the Sun.

The radiation pressure experienced at a particular location varies inversely with distance, and so does the resulting force. Consequently, the ratio of forces experienced at altitudes 750 km and 3000 km, due to radiation pressure can be expressed as,

$$\frac{F_{750}}{F_{3000}} = \left(\frac{r_{3000}}{r_{750}}\right)^2 \tag{9}$$

Consider Figure 2, where D_{SE} is the Sun-Earth distance, R_E is the Earth radius and h is the altitude of the orbiting body. At position A,

$$\frac{F_{750}}{F_{3000}} = \left(\frac{D_{SE} - R_E - 3000}{D_{SE} - R_E - 3000}\right)^2 = 0.99997\tag{10}$$

Applying the same equation at position B,

$$\frac{F_{750}}{F_{3000}} = \left(\frac{D_{SE}^2 + (R_E + 3000)^2}{D_{SE}^2 + (R_E + 750)^2}\right) = 1.00000 \tag{11}$$

From equations (10) and (11), it is evident that the impact of radiation pressure would be nearly identical at both altitudes.




Figure 2. Illustration depicting the Sun, Earth, and a representative debris object at two positions in orbit labeled A and B; note that the figure is not drawn to scale.

To comprehend the reason behind the abrupt changes in slope during distinct periods across the three solar cycles, it is imperative to analyze the correlation between altitude and various indices such as SN, F10.7, Dst, and AL for the corresponding intervals. The normalized correlation coefficients are graphically represented against the time lag in days for the four sample debris objects collectively, as illustrated in Figure 3. The values of the maximum correlation coefficients are added in brackets next to the corresponding curve labels. As expected, the maximum correlation occurs at a negative value of lag since the response in altitude decay will occur after any changes in the indices occur. Additionally, the p-values are close to zero, indicating statistical significance. Notably, it is observed that SN exhibits the highest correlation. Table 1 provides details on the associated lag values and mean altitudes across the three solar cycles. If we examine time lags for various indices against mean altitude, coupled with the details in Table 1, a noticeable trend emerges: as altitude increases, the time lag extends, implying that it takes more time for the impact to reach higher altitudes in the thermosphere. Observing debris objects with SATCAT IDs 00732 and 00831, separated by nearly 1000 km, it is seen that the solar cycle variation takes approximately 14 months longer to impact debris object 00831 at an altitude above 2000 km.

In Figure 1, six regions are evident, each marked by unique slopes. Three of these segments, outlined by dashed red lines, showcase steeper slopes, while the other three exhibit




Figure 3. Normalized correlation coefficients plotted against time lag for SN, F10.7, Dst, and AL indices, with dashed lines indicating the respective peak locations.

| NORAD ID | Mean Alt (km) | SN Lag (Days) | F10.7 Lag (Days) | Dst Lag (Days) | AL Lag (Days) |
|----------|------------------|------------------|---------------------|-------------------|------------------|
| 22 | 653 | -1193 | -1230 | -1170 | -1222 |
| 115 | 659 | -958 | -897 | -863 | -715 |
| 732 | 1137 | -1291 | -1358 | -1295 | -1358 |
| 831 | 2175 | -1731 | -1884 | -1887 | -2001 |

 Table 1.
 Mean Altitude and Lag for the sample objects

a more gradual decline. Figure 4a illustrates the respective slopes from these six regions, along with the average F10.7 index for the corresponding period. Notably, a higher F10.7 index correlates with a steeper slope (region 1,3,5).

Figure 4b illustrates the slopes calculated by applying a linear fit to the points within the red-marked regions in Figure 1, plotted against the apparent altitude at the start of decay for the three solar cycles. The corresponding values of the fitted slopes are listed in Tables 1. It shows a decreasing trend in decay rate with altitude between 500-1000 km.

Figure 5 displays histograms portraying the average decay rate for both LEO and MEO objects across three solar cycles. The decay rates are initially calculated using adjacent epochs, and subsequently, these values are averaged over each of the solar cycles to determine the mean decay rate for the respective cycle. These histograms categorize the values into bins with an optimal bin width for each solar cycle determined by Scott's rule. Scott's rule minimizes the integrated mean squared error in the bin approximation under the assumption that the data is approximately Gaussian. Based on calculations, for LEO debris during Solar Cycle 22, there are 6 bins with a bin width of 1419.7 meters per hour, Solar Cycle 23 has 5 bins with a bin width of 163.3 meters per hour, and Solar Cycle 24 has 5 bins with a bin width of 83.4 meters per hour. Significantly, higher decay rates are observed in Solar Cycles 22 and 23. The maximum decay rate occurred during Solar Cycle 22, specifically within the bin of -8940 to -7520 m/h. For Solar Cycle 23, the maximum decay was recorded in the bin -924 to -761 m/h, and for Solar Cycle 24, the maximum decay was observed in the

bin -490 to -407 m/h. It appears that Solar Cycle 24 does not have a significant impact on the higher-value bins in these histograms. This suggests that it has less of an effect on the orbital decay of space debris.




Figure 4. (a) Relationship between F10.7 solar flux and decay rate (b) Slope vs apparent altitude for three solar cycles

Following Scott's rule for optimized bin width of the histogram, in the case of MEO space debris, Solar Cycle 22 has 3 bins with a width of 264.9 meters per hour, Solar Cycle 23 also has 3 bins but with a width of 68.5 meters per hour, and Solar Cycle 24 comprises 4 bins with a width of 65.6 meters per hour. Notably, higher decay rates are evident in Solar Cycle 22. The maximum decay rate occurred during Solar Cycle 22, specifically within the bin of -1050 to -785 m/h. For Solar Cycle 23, the maximum decay was recorded in the bin -233 to -164 m/h, and for Solar Cycle 24, the maximum decay was observed in the bin -286 to -220 m/h. Figure 4c clearly illustrates that the mean rate of decay is higher in Low Earth Orbits compared to Medium Earth Orbits. This discrepancy arises because atmospheric density directly influences drag, and as atmospheric density significantly decreases beyond LEO, there is less drag force acting on MEO objects, resulting in a lower rate of decay. Nevertheless, it is crucial to note that even MEO orbit space debris exhibits notable decay.

4 Discussion

With the rapid growth in the space sector and an exponential increase in satellites and associated space debris, monitoring and mitigation of space debris have become of paramount importance. Many space debris have lifetimes ranging from years to decades, making the impact of long-term changes in solar activity on their orbit a highly timely topic for investigation.

The present study conducts a systematic analysis of the solar cycle's influence on the orbits of 95 space debris that have been tracked since 1967. Our analysis reveals several key observations. First and foremost, a clear and compelling high correlation between solar activity and the rate of orbital decay emerges. Solar Cycle 22 stands out as the most effective, followed by Solar Cycle 23, while Solar Cycle 24 registers as the least effective in terms of impacting the orbital decay of space debris. This decrease in solar activity from Cycle 22 to 24 is reflected in overall thermospheric cooling Ogawa et al. (2014). This direct relationship significantly impacts the steepness of the decay curves, resulting in more pronounced decay during the peak periods of Solar Cycles 22 and 23. These trends closely mirror the patterns observed in sunspot numbers. The rapid decay is observed near the peak of each solar cycle bounded by $\sim 70\%$ from the peak amplitude of the solar cycle seen in SN. During the active phase of a solar cycle, characterized by a higher number of sunspots, the increased energy reaching the Earth heats the atmosphere, causing it to expand beyond its quiet sun state,




Figure 5. Distribution of average decay rates: upper panel - LEO objects, lower panel - MEO objects

leading to an order of magnitude increase in neutral density in the thermosphere (Marcos et al. (2005)). Consequently, an object will experience an accelerated orbital decay rate due to the increased drag Stansbery & Foster (2004). Therefore, this observation reinforces the fundamental role of solar activity in shaping the dynamics of neutral atmosphere expansion and contraction and its impact on the orbital decay of space debris.

Furthermore, as anticipated, the low atmospheric densities at MEO altitudes lead to a reduction in drag force, consequently resulting in lower mean decay rates for MEO objects when compared to LEO objects. This distinction is a crucial factor in understanding the differential rates of orbital decay in these two orbital regimes. However, the influence of the solar cycle extends to altitudes exceeding 2000 km, making it a thought-provoking observation. The solar radiation pressure also exerts additional force on the debris, our estimates show that LEO and MEO orbital debris should experience almost same force and instantaneously. Whereas, these effects are seen to be more apparent at higher altitudes with a discernible time delay of over 14 months compared to their manifestation at lower altitudes. This unequivocally confirms that the thermospheric effects of the solar activity cycle endure in the thermosphere for a duration exceeding a year, progressively becoming more evident as altitude increases. It is imperative that future modeling studies are conducted to thoroughly decode these phenomena.

5 Concluding Remarks

This novel study examines the long-term impact of solar activity on 95 orbital debris objects spanning three solar cycles. The findings revealed in this study provide valuable insights into the significant influence of solar cycles on the orbital decay of space debris, with a particular focus on both LEO and MEO objects. The direct impact of solar activity in influencing the orbital decay of debris objects is seen and how the rates of orbital decay vary in these two orbital regimes is seen. Additionally, we have seen that thermospheric effects of the solar cycle persist in the thermosphere for well over a year, gradually manifesting with increasing altitude. In the absence of in-situ measurements of atmospheric density or drag force at altitudes above 2000 km, we can employ this as a proxy to comprehend the impact and dynamics of the solar cycle in this region. The insights from these observations hold significance for future space missions, satellite planning, and specifically strategies for mitigating space debris. It underscores the continual importance of monitoring the impact of solar activity on space debris in the context of ongoing and future rapid growth in space exploration and satellite operations.

Acknowledgements

The authors would like to acknowledge the Space-Track website (www.space-track .org) for providing access to the TLE data of the debris objects.

References

- Bruinsma, S., Dudok de Wit, T., Fuller-Rowell, T., Garcia-Sage, K., Mehta, P., Schiemenz, F., ... Elvidge, S. (2023). Thermosphere and satellite drag. Advances in Space Research. doi: https://doi.org/10.1016/j.asr.2023.05.011
- Chen, Y., Liu, L., & Wan, W. (2011). Does the f10. 7 index correctly describe solar euv flux during the deep solar minimum of 2007–2009? Journal of Geophysical Research: Space Physics, 116(A4).
- Clette, F., Cliver, E., Lefèvre, L., Svalgaard, L., & Vaquero, J. (2015, 08). Revision of the sunspot number(s). Space Weather, 13.
- Clette, F., & Lefèvre, L. (2016, nov). The new sunspot number: Assembling all corrections. Solar Physics, 291 (9-10), 2629–2651.
- Clette, F., Svalgaard, L., Vaquero, J. M., & Cliver, E. W. (2014, aug). Revisiting the sunspot number. Space Science Reviews, 186(1-4), 35-103.
- Crisp, N., Roberts, P., Romano, F., Smith, K., Oiko, V., Sulliotti-Linner, V., ... Seminari, S. (2021). System modelling of very low earth orbit satellites for earth observation. Acta Astronautica, 187, 475-491. doi: https://doi.org/10.1016/j.actaastro.2021.07.004
- Dang, T., Li, X., Luo, B., Li, R., Zhang, B., Pham, K., ... Wang, Y. (2022). Unveiling the space weather during the starlink satellites destruction event on 4 february 2022. *Space Weather*, 20(8), e2022SW003152. doi: https://doi.org/10.1029/2022SW003152
- Gopalswamy, N. (2006). Properties of interplanetary coronal mass ejections. Space Science Reviews, 124, 145–168.
- Graham, G., Kearsley, A., Grady, M., Wright, I., Griffiths, A., & McDonnell, J. (1999). Hypervelocity impacts in low earth orbit: Cosmic dust versus space debris. Advances in Space Research, 23(1), 95-100.
- Harris, I., & Priester, W. (1962). Theoretical models for the solar-cycle variation of the upper atmosphere. Journal of Geophysical Research, 67(12), 4585–4591.
- Hathaway, D. H. (2015). The solar cycle. Living reviews in solar physics, 12, 1–87.
- Hayakawa, H., Nevanlinna, H., Blake, S. P., Ebihara, Y., Bhaskar, A. T., & Miyoshi, Y. (2022). Temporal variations of the three geomagnetic field components at colaba observatory around the carrington storm in 1859. *The Astrophysical Journal*, 928(1), 32.
- Hayes, A. D., & Caverly, R. J. (2023). Model predictive tracking of spacecraft deorbit

trajectories using drag modulation. Acta Astronautica, 202, 670-685. doi: https://doi.org/10.1016/j.actaastro.2022.10.057

- Huang, C.-S., & Reeves, G. D. (2005). Periodic substorms: A new periodicity of 2-3 hours in the magnetosphere. In A. Lui, Y. Kamide, & G. Consolini (Eds.), *Multiscale* coupling of sun-earth processes (p. 265-279). Amsterdam: Elsevier Science B.V. doi: https://doi.org/10.1016/B978-044451881-1/50019-8
- Jayalekshmi, G. L., Pant, T. K., & Prince, P. R. (2022, July). Sunspot-Cycle Evolution of Major Periodicities of Solar Activity. Solar Physics, 297(7), 85. doi: 10.1007/s11207-022 -02017-1
- Kataoka, R., Shiota, D., Fujiwara, H., Jin, H., Tao, C., Shinagawa, H., & Miyoshi, Y. (2022, 12).
 Unexpected space weather causing the reentry of 38 starlink satellites in february 2022. Journal of Space Weather and Space Climate, 12. doi: 10.1051/swsc/2022034
- Kessler, D. J., Johnson, N. L., Liou, J., & Matney, M. (2010). The kessler syndrome: implications to future space operations. Advances in the Astronautical Sciences, 137(8), 2010.
- Kleinig, T., Smith, B., & Capon, C. (2022). Collision avoidance of satellites using ionospheric drag. Acta Astronautica, 198, 45-55. doi: https://doi.org/10.1016/j.actaastro.2022.03 .017
- Klinkrad, H. (2006). Space debris: models and risk analysis. Springer Science & Business Media.
- Lean, J. (1987). Solar ultraviolet irradiance variations: A review. Journal of Geophysical Research: Atmospheres, 92(D1), 839–868.
- Li, C., Ouyang, Z., & Du, H. (2002, 01). Space debris and space environment. Quaternary Sciences, 22, 540-551.
- Marcos, F. A., Wise, J. O., Kendra, M. J., Grossbard, N. J., & Bowman, B. R. (2005). Detection of a long-term decrease in thermospheric neutral density. *Geophysical Research Letters*, 32(4).
- Mlynczak, M., Hunt, L., Marshall, B., & Russell III, J. (2018, 10). Infrared radiation in the thermosphere near the end of solar cycle 24. *Geophysical Research Letters*, 45. doi: 10.1029/2018GL080389
- Nwankwo, V. U. (2018). Space weather: Response of the atmosphere to solar activity and its implications for leo satellites aerodynamic drag. In *Exploring the universe: From near space to extra-galactic: A collection of research reviews on contemporary astrophysics and space science* (pp. 637–644).
- Ogawa, Y., Motoba, T., Buchert, S. C., Häggström, I., & Nozawa, S. (2014). Upper atmosphere cooling over the past 33 years. *Geophysical Research Letters*, 41(15), 5629–5635.
- Oliveira, D. M., Zesta, E., Hayakawa, H., & Bhaskar, A. (2020). Estimating satellite orbital drag during historical magnetic superstorms. Space weather, 18(11), e2020SW002472.
- Richardson, I. G., & Cane, H. V. (2012). Near-earth solar wind flows and related geomagnetic activity during more than four solar cycles (1963–2011). Journal of Space Weather and Space Climate, 2, A02.
- Rostoker, G. (1972). Geomagnetic indices. Reviews of Geophysics, 10(4), 935–950.
- Smith, B., Capon, C., Brown, M., & Boyce, R. (2020). Ionospheric drag for accelerated deorbit from upper low earth orbit. Acta Astronautica, 176, 520-530. doi: https:// doi.org/10.1016/j.actaastro.2020.07.007
- Stansbery, E. G., & Foster, J. L. (2004). Monitoring the low earth orbit debris environment over an 11-year solar cycle. Advances in Space Research, 34(5), 878-883. (Space Debris)
- Tsurutani, B. T., Gonzalez, W. D., Lakhina, G., & Alex, S. (2003). The extreme magnetic storm of 1–2 september 1859. *Journal of Geophysical Research: Space Physics*, 108(A7).
- Vallado, D., & Cefola, P. (2012, 01). Two-line element sets practice and use. Proceedings of the International Astronautical Congress, IAC, 7, 5812-5825.
- Walterscheid, R. (1989). Solar cycle effects on the upper atmosphere-implications for satellite drag. Journal of spacecraft and rockets, 26(6), 439–444.

- Weng, L., Lei, J., Zhong, J., Dou, X., & Fang, H. (2020). A machine-learning approach to derive long-term trends of thermospheric density. *Geophysical Research Letters*, 47(6), e2020GL087140.
- Zhang, J., Feng, Y., Zhang, J., & Li, Y. (2023, 10). The short time prediction of the dst index based on the long-short time memory and empirical mode decomposition-long-short time memory models. *Applied Sciences*, 13, 11824. doi: 10.3390/app132111824

Appendix A List of objects used in the analysis

Table A1: Debris Objects

| | Mean Rate of Decay (m/h) | | | Slope at Peak (m/h) | | |
|-------------------|----------------------------|------------------------|------------------------------------|------------------------|------------------------|------------------------|
| NORAD ID | SC 22 | SC 23 | SC 24 | SC 22 | SC 23 | SC 24 |
| 22 | -4.86E+03 | -9.37E+01 | -1.14E+02 | -1.42E+00 | -1.40E+00 | -8.13E-01 |
| 29 | -1.76E+03 | -2.47E+01 | -6.48E+01 | -5.24E-01 | -4.61E-01 | -1.73E-01 |
| 45 | -2.17E+02 | -5.72E+02 | -4.90E+01 | -3.92E-01 | -2.99E-01 | -1.11E-01 |
| 46 | -1.14E+02 | -2.46E+02 | -7.06E+01 | -7.30E-01 | -6.05E-01 | -2.26E-01 |
| 85 | -2.54E+03 | -5.98E+02 | -8.64E+00 | -4.50E+00 | -4.43E+00 | -2.67E+00 |
| 107 | -9.54E+00 | -5.27E+01 | -4.53E+02 | -1.71E+00 | -1.44E+00 | -6.50E-01 |
| 115 | -8.94E+01 | -2.20E+01 | -2.93E+02 | -9.03E-01 | -9.31E-01 | -4.40E-01 |
| 130 | -0.34L+03 -1.41E+03 | -2.20E+00 -8.97E+01 | -2.94E-02 | -1.33E-01 | -9.87E-01 | -3.68E-02 |
| 144 | -9.08E+02 | -8.83E+01 | -2.54E-02 -4.60E-02 | -2.59E-01 | -3.07E-02 -2.12E-01 | -3.00E-02 -8.21E-02 |
| 150 | -3.00E + 02 -1.10E + 03 | -1.29E+01 | -9.37E+01 | -4.32E-01 | -3.31E-01 | -0.21E-02 -1.35E-01 |
| 162 | -1.66E+03 | -1.29E+01 -1.59E+02 | -3.57E+01 -1.57E+01 | -4.32E-01 -2.36E-01 | -3.31E-01 -1.81E-01 | -5.90E-02 |
| 162 | -1.00E+03 -2.18E+03 | -1.59E+02 -6.43E+01 | $-1.57 \pm +01$ -8.18 ± -02 | -2.56E-01 -2.65E-01 | -1.96E-01 | -5.90E-02 -6.61E-02 |
| 223 | -2.76E+03 | -0.43E+01 -9.24E+02 | -0.18E-02 -1.28E-02 | -2.05E-01 -7.21E-02 | -1.90E-01 -5.24E-02 | -0.01E-02 -1.95E-02 |
| $\frac{223}{227}$ | -2.76E+02 -1.63E+00 | -9.24E+02 -1.65E-01 | -1.28E-02 -5.55E-02 | -7.21E-02 -4.23E-01 | -5.24E-02 -3.26E-01 | -1.95E-02 -1.09E-01 |
| | | | | | | -1.09E-01 -7.13E-01 |
| 228 | -6.27E-01 | -5.41E-01 | -2.99E-01 -5.11E-02 | -1.04E+00 -2.72E-01 | -1.20E+00 | |
| 232 | -2.51E-01 | -1.07E-01 | | | -1.93E-01 | -7.54E-02 |
| 252 | -6.46E+02 | -6.49E+01 | -6.04E-02 | -2.65E-01 | -2.08E-01 | -8.69E-02 |
| 262 | -3.35E+03 | -4.32E+00 | -3.16E-01 | -1.15E+00 | -1.10E+00 | -6.31E-01 |
| 309 | -5.41E+00 | -2.04E-01 | -9.82E-01 | -5.35E-01 | -4.46E-01 | -1.60E-01 |
| 323 | -2.84E-01 | -2.38E-01 | -1.25E-01 | -5.14E-01 | -4.07E-01 | -1.49E-01 |
| 397 | -2.96E+00 | -2.97E-01 | -1.29E-01 | -6.60E-01 | -6.31E-01 | -2.67E-01 |
| 405 | -4.57E+02 | -7.47E+02 | -5.47E+01 | -1.52E-01 | -1.17E-01 | -4.97E-02 |
| 420 | -2.43E+02 | -2.07E+01 | -1.22E+01 | -6.59E-01 | -5.82E-01 | -2.20E-01 |
| 424 | -6.76E-02 | -3.87E-02 | -1.98E-02 | -7.83E-02 | -5.92E-02 | -2.51E-02 |
| 426 | -1.41E-01 | -2.58E-02 | -1.61E-02 | -5.05E-02 | -3.81E-02 | -1.73E-02 |
| 465 | -4.97E+03 | -3.21E+01 | -1.03E+01 | -9.98E-01 | -9.71E-01 | -3.60E-01 |
| 510 | -4.63E-02 | -2.96E-02 | -1.44E-02 | -6.12E-02 | -4.60E-02 | -1.90E-02 |
| 511 | -1.15E-01 | -4.26E-02 | -1.68E-02 | -6.10E-02 | -4.45E-02 | -1.89E-02 |
| 614 | -3.91E + 00 | -3.54E + 00 | -3.00E + 00 | -6.56E + 00 | -6.58E + 00 | -5.14E + 00 |
| 716 | -2.42E-01 | -1.76E-01 | -6.73E-02 | -4.72E-01 | -4.02E-01 | -1.45E-01 |
| 720 | -3.19E-01 | -1.96E-01 | -7.17E-02 | -5.16E-01 | -4.29E-01 | -1.49E-01 |
| 724 | -6.09E + 01 | -2.80E+00 | -1.11E + 00 | -4.19E+00 | -3.99E+00 | -2.48E+00 |
| 727 | -9.07E-02 | -5.12E-02 | -1.98E-02 | -7.88E-02 | -5.76E-02 | -2.13E-02 |
| 728 | -4.35E-01 | -5.61E-02 | -2.26E-02 | -1.13E-01 | -8.32E-02 | -3.13E-02 |
| 729 | -2.11E-01 | -3.92E-02 | -1.40E-02 | -6.17E-02 | -4.49E-02 | -1.66E-02 |
| 730 | -6.03E-02 | -4.68 E - 02 | -1.74E-02 | -8.22E-02 | -5.98E-02 | -2.20E-02 |
| 731 | -2.27E-01 | -5.07E-02 | -1.79E-02 | -8.75E-02 | -6.41E-02 | -2.37E-02 |
| 733 | -5.25E-01 | -8.40E-02 | -3.64E-02 | -2.50E-01 | -1.78E-01 | -7.06E-02 |
| 734 | -9.08E-02 | -6.41E-02 | -2.44E-02 | -1.67E-01 | -1.20E-01 | -3.56E-02 |
| 735 | -9.41E-02 | -5.98E-02 | -2.38E-02 | -1.56E-01 | -1.13E-01 | -3.31E-02 |
| 801 | -2.09E-01 | -1.86E-01 | -7.08E-02 | -3.92E-01 | -3.19E-01 | -1.15E-01 |
| 805 | -7.32E + 00 | -2.09E+00 | -1.79E-01 | -8.88E-01 | -7.800E-01 | -3.53E-01 |
| 809 | -7.13E-01 | -1.23E-01 | -7.23E-02 | -2.71E-01 | -2.21E-01 | -9.12E-02 |
| 812 | -3.26E-01 | -6.42E-02 | -2.25E-02 | -1.47E-01 | -9.85E-02 | -3.72E-02 |
| 813 | -2.04E-01 | -1.62E + 00 | -2.10E-02 | -1.35E-01 | -9.06E-02 | -3.26E-02 |
| 815 | -1.10E-01 | -8.11E-02 | -3.08E-02 | -2.29E-01 | -1.72E-01 | -5.29E-02 |
| 870 | -1.01E-01 | -5.21E-02 | -2.28E-02 | -1.23E-01 | -9.16E-02 | -3.30E-02 |
| 871 | -1.45E-01 | -1.14E-01 | -5.13E-02 | -2.84E-01 | -2.28E-01 | -8.52E-02 |

Continued on the next page.

| | Mean Rate of Decay (m/h) | | | Slope at Peak (m/h) | | |
|----------|--------------------------|-------------|-------------|---------------------|-------------|-------------|
| NORAD ID | SC 22 | SC 23 | SC 24 | SC 22 | SC 23 | SC 24 |
| 876 | -4.94E-01 | -1.47E-01 | -3.29E + 00 | -3.64E-01 | -2.73E-01 | -8.90E-02 |
| 877 | -1.63E-01 | -1.21E-01 | -7.20E-01 | -3.18E-01 | -2.53E-01 | -8.52E-02 |
| 899 | -1.18E-01 | -6.38E-02 | -3.06E-02 | -1.68E-01 | -1.22E-01 | -4.88E-02 |
| 900 | -2.69E-01 | -2.60E-01 | -1.09E-01 | -4.65E-01 | -3.81E-01 | -1.65E-01 |
| 903 | -1.03E+02 | -2.25E-01 | -1.17E-01 | -5.43E-01 | -4.57E-01 | -1.99E-01 |
| 907 | -7.32E-02 | -5.26E-02 | -2.43E-02 | -1.22E-01 | -9.02E-02 | -3.38E-02 |
| 976 | -6.07E-01 | -2.00E-01 | -9.18E-02 | -4.88E-01 | -3.92E-01 | -1.44E-01 |
| 1943 | -1.30E + 03 | -9.39E+01 | -9.32E + 01 | -7.33E-01 | -6.51E-01 | -2.52E-01 |
| 2097 | -2.63E-01 | -5.02E + 00 | -1.01E-01 | -4.50E-01 | -3.70E-01 | -1.66E-01 |
| 2986 | -4.40E+00 | -7.69E-02 | -4.08E-02 | -2.17E-01 | -1.16E-01 | -5.43E-02 |
| 3741 | -8.92E+01 | -5.56E + 00 | -4.48E-01 | -2.30E+00 | -2.22E+00 | -9.68E-01 |
| 13513 | -4.12E+03 | -6.84E + 01 | -7.99E + 01 | -6.89E-01 | -5.71E-01 | -2.53E-01 |
| 53 | -1.05E+03 | -8.91E + 01 | -9.26E + 01 | -3.04E-02 | -2.43E-02 | -1.13E-02 |
| 58 | -5.04E+01 | -2.33E+02 | -5.01E + 01 | -2.14E-02 | -1.57E-02 | -6.26E-03 |
| 59 | -9.07E+01 | -2.26E + 02 | -2.86E + 02 | -3.67E-02 | -2.76E-02 | -1.06E-02 |
| 82 | -4.35E+02 | -6.72E + 01 | -9.36E-01 | -2.27E-01 | -2.46E-01 | -2.61E-02 |
| 202 | -1.30E-01 | -3.72E-02 | -1.53E-02 | -4.89E-02 | -3.54E-02 | -1.50E-02 |
| 204 | -1.85E+00 | -3.47E-02 | -1.56E-02 | -3.92E-02 | -2.91E-02 | -1.14E-02 |
| 205 | -1.25E-01 | -3.20E-02 | -1.36E-02 | -3.19E-02 | -2.27E-02 | -9.17E-03 |
| 558 | -1.03E+03 | -2.20E + 02 | -6.79E-01 | -1.44E-01 | -1.05E-01 | -4.76E-02 |
| 669 | -1.87E-02 | -1.49E-02 | -9.38E-03 | -2.78E-02 | -2.12E-02 | -9.16E-03 |
| 670 | -3.10E-02 | -2.44E-02 | -1.37E-02 | -6.11E-02 | -4.66E-02 | -2.09E-02 |
| 671 | -3.54E-02 | -2.70E-02 | -1.54E-02 | -6.60E-02 | -5.04E-02 | -2.28E-02 |
| 694 | -9.41E-01 | -5.35E-01 | -2.38E-01 | -1.47E + 00 | -1.23E + 00 | -5.34E-01 |
| 700 | -2.16E-01 | -1.50E-01 | -4.61E-02 | -3.44E-01 | -2.60E-01 | -9.11E-02 |
| 703 | -2.33E-02 | -1.49E-02 | -1.29E-02 | -3.08E-02 | -2.28E-02 | -9.91E-03 |
| 704 | -3.47E-02 | -2.26E-02 | -2.23E-02 | -5.95E-02 | -3.62E-02 | -1.56E-02 |
| 705 | -3.27E-02 | -2.39E-02 | -1.32E-02 | -6.06E-02 | -4.61E-02 | -1.89E-02 |
| 706 | -1.07E-01 | -9.00E-02 | -4.38E-02 | -2.10E-01 | -1.62E-01 | -7.32E-02 |
| 721 | -1.87E-01 | -1.61E-01 | -9.18E-02 | -2.29E-01 | -3.18E-01 | -5.77E-02 |
| 732 | -4.46E + 00 | -1.63E + 00 | -7.40E-01 | -2.79E + 00 | -3.18E + 00 | -1.72E + 00 |
| 739 | -2.99E-01 | -2.15E-01 | -7.56E-02 | -4.97E-01 | -4.18E-01 | -1.58E-01 |
| 741 | -2.25E-02 | -2.35E-02 | -1.69E-02 | -3.15E-02 | -2.24E-02 | -9.83E-03 |
| 742 | -4.10E-02 | -2.50E-02 | -1.53E-02 | -5.83E-02 | -4.41E-02 | -2.12E-02 |
| 743 | -4.21E-02 | -2.85E-02 | -1.81E-02 | -6.86E-02 | -5.13E-02 | -2.35E-02 |
| 745 | -1.36E-01 | -9.03E-02 | -4.45E-02 | -1.97E-01 | -1.58E-01 | -6.86E-02 |
| 746 | -3.24E+00 | -1.09E+01 | -5.86E + 01 | -2.29E+00 | -1.83E+00 | -8.38E-01 |
| 750 | -5.90E + 01 | -2.55E+01 | -2.25E+02 | -1.06E+01 | -9.16E + 00 | -4.77E + 00 |
| 753 | -9.49E-02 | -8.29E-02 | -4.14E-02 | -1.93E-01 | -1.48E-01 | -6.54E-02 |
| 829 | -5.18E + 00 | -2.58E + 01 | -2.87E + 00 | -2.33E+00 | -1.95E+00 | -8.60E-01 |
| 831 | -2.58E + 01 | -5.51E + 00 | -2.13E + 01 | -1.11E + 01 | -9.30E+00 | -4.90E+00 |
| 893 | -2.40E-02 | -1.65E-02 | -9.90E-03 | -3.41E-02 | -2.53E-02 | -1.08E-02 |
| 897 | -3.37E-02 | -2.09E-02 | -1.26E-02 | -5.17E-02 | -3.80E-02 | -1.62E-02 |
| 901 | -4.65E-02 | -3.45E-02 | -1.84E-02 | -8.30E-02 | -6.26E-02 | -2.90E-02 |
| 902 | -2.60E-02 | -1.64E-02 | -1.05E-02 | -3.81E-02 | -2.80E-02 | -1.25E-02 |
| 977 | -1.35E-01 | -1.39E+00 | -5.39E-02 | -2.61E-01 | -2.13E-01 | -8.38E-02 |
| 3750 | -4.74E+01 | -1.55E+01 | -9.63E-01 | -4.72E + 00 | -4.20E+00 | -1.69E+00 |