

OPTIMAL ASYMPTOTIC VOLUME RATIO FOR NONCOMPACT 3-MANIFOLDS WITH ASYMPTOTICALLY NONNEGATIVE RICCI CURVATURE AND A UNIFORMLY POSITIVE SCALAR CURVATURE LOWER BOUND

XIAN-TAO HUANG AND SHUAI LIU

ABSTRACT. In this paper, we study 3-dimensional complete non-compact Riemannian manifolds with asymptotically nonnegative Ricci curvature and a uniformly positive scalar curvature lower bound. Our main result is that, if this manifold has k ends and finite first Betti number, then it has at most linear volume growth, and furthermore, if the negative part of Ricci curvature decays sufficiently fast at infinity, then we have an optimal asymptotic volume ratio $\limsup_{r \rightarrow \infty} \frac{\text{Vol}(B(p, r))}{r} \leq 4k\pi$. In particular, our results apply to 3-dimensional complete non-compact Riemannian manifolds with nonnegative Ricci curvature and a uniformly positive scalar curvature lower bound.

Keywords: scalar curvature, asymptotically nonnegative Ricci curvature, 3-dimensional manifold, linear volume growth.

1. INTRODUCTION

In this paper, we investigate the influence of positive scalar curvature on the geometric properties of 3-dimensional complete non-compact Riemannian manifolds with asymptotically nonnegative Ricci curvature.

Definition 1.1. We say a Riemannian manifold having asymptotically nonnegative Ricci curvature if for any fixed point $p \in M$, there exists a positive function $f(r)$ such that $\lim_{r \rightarrow \infty} f(r) = 0$ and for any point $x \in M$, it holds that $\text{Ric}(x) \geq -f(d(x, p))$.

It is believed that the uniformly positive lower bound of scalar curvature has an influence on the geometric properties of a manifold. In [8], Gromov raised the following conjecture:

Conjecture 1.2. *Let (M^n, g) be an n -dimensional complete non-compact Riemannian manifold with nonnegative Ricci curvature and $\text{Sc}_g \geq 1$. Do we have*

$$(1.1) \quad \limsup_{r \rightarrow \infty} \frac{\text{Vol}(B(p, r))}{r^{n-2}} < \infty?$$

Gromov's conjecture is solved in some special cases.

In [27], Zhu proves that, (1.1) holds on (M^n, g) with nonnegative Ricci curvature and $\text{Sc}_g \geq 1$ provided M has a positive lower bound of the injective radius. In the 3-dimensional case, there are more progresses. Zhu ([27]) verified (1.1) holds on 3-manifolds with nonnegative Ricci curvature and $\text{Sc}_g \geq 1$ under the additional assumption that $\inf_{q \in M} \text{Vol}(B(q, 1)) \geq v_0 > 0$. Later on, Munteanu and Wang (see [15]) solved completely the 3-dimensional case by analyzing certain properties of the Green's function. More precisely, the following result has been established in [15]:

Theorem 1.3 ([15]). *Let (M^3, g) be a 3-dimensional complete noncompact Riemannian manifold with nonnegative curvature and $\text{Sc}_g \geq 1$. Then there exists a universal constant*

$C > 0$ such that for any fixed point $p \in M$ and $r > 0$,

$$(1.2) \quad \frac{\text{Vol}(B(p, r))}{r} < C.$$

We remark that in [15] Munteanu and Wang also study 3-manifolds with certain type of asymptotically nonnegative Ricci curvature.

Later, Chodosh, Li and Stryker ([6]) use the theory of μ -bubbles and Cheeger-Colding's almost splitting theorem to recover Theorem 1.3, with the exception that the constant C in [6] is not universal, but depends on M (on the other hand, we remark that both in [15] [6], the authors consider the growth rate of volume of geodesic ball in the case that the positive scalar curvature lower bound decays compared to some power of the distance function, and the class handled in [6] is wider than that of [15]). In a recent paper [21], by a more detailed exploration of the method in [6], Wang is able to recover Theorem 1.3 completely.

Note that Theorem 1.3 says that the manifold has at most linear volume growth. We remark that Calabi [2] and Yau [25] independently proved that every noncompact manifold with nonnegative Ricci curvature has at least linear volume growth.

In [26], Yau proposed the following question.

Question 1.4. *Let (M^n, g) be an n -dimensional complete non-compact Riemannian manifold with nonnegative Ricci curvature. Then for any fixed point $p \in M$, do we have*

$$(1.3) \quad \limsup_{r \rightarrow \infty} r^{n-2} \int_{B(p, r)} \text{Sc}_g < \infty?$$

It is easy to see that a positive answer of Question 1.4 will give an answer to Conjecture 1.2.

In [27], Zhu proved the correctness of Question 1.4 for 3-dimensional manifolds with a pole (recall that M has a pole $p \in M$ means that the exponential map at p is a diffeomorphism) and $\text{Ric}_g \geq 0$. In [24], Xu proved the following sharp result:

Theorem 1.5. *Let (M, g) be a 3-dimensional Riemannian manifold with a pole. Suppose $\text{Ric}_g \geq 0$, then we have*

$$(1.4) \quad \lim_{r \rightarrow \infty} \frac{\int_{B(p, r)} \text{Sc}_g}{r} = 8\pi(1 - \lim_{r \rightarrow \infty} \frac{\text{Vol}(B(p, r))}{\omega_3 r^3}),$$

where ω_3 is the volume of unit ball in $\mathbb{R}^3$.

The aim of the present paper consists of two parts. Firstly, in Theorem 1.3, the universal constant C is not explicit. Motivated by Theorem 1.5, we want to find an optimal volume ratio at infinity. Secondly, we want to explore the influence of positive scalar curvature on the geometry on a larger class of manifolds which may have negative Ricci curvature somewhere.

The following theorem is our main result.

Theorem 1.6. *Let (M^3, g) be a 3-dimensional complete noncompact Riemannian manifold with k ends and finite first Betti number. Suppose $S_g \geq 2$ outside a compact set, and for a fixed pointed $p \in M$ there is a positive function with $\lim_{r \rightarrow \infty} f(r) = 0$, so that*

$$\text{Ric}_g(x) \geq -f(r(x)).$$

Then the following conclusions were justified.

(1) The macroscopic dimension of M is equal to 1, and there exists $C > 0$ such that

$$(1.5) \quad \limsup_{r \rightarrow \infty} \frac{\text{Vol}(B(p, r))}{r} \leq C.$$

(2) If $f(r) = \frac{1}{r^{3+\alpha}}$ for some $\alpha > 0$, then there

$$(1.6) \quad \limsup_{r \rightarrow \infty} \frac{\text{Vol}(B(p, r))}{r} \leq 4k\pi.$$

The notion of macroscopic dimension is introduced by Gromov (see [9]):

Definition 1.7. We say a metric space has macroscopic dimension at most k if there exist a constant $C > 0$, a k -dimensional polyhedron P and a proper continuous map $\phi : X \rightarrow P$ such that $\text{diam}(\phi^{-1}(p)) \leq C$ for every $p \in P$.

Remark 1.8. Recall that any manifolds with nonnegative Ricci curvature has at most 2 ends. In addition, 3-dimensional manifolds with nonnegative Ricci curvature have been classified by Liu in [13], and it is known that every such manifold must have finite first Betti number. Therefore, Theorem 1.6 holds automatically on 3-manifolds with nonnegative Ricci curvature and positive scalar curvature. In this case, (2) in Theorem 1.6 gives a sharp upper bound. If $k = 2$, then according to Cheeger-Gromoll's splitting theorem, M is isometric to $\mathbb{R} \times S$ with S a compact manifold with sectional curvature ≥ 1 , and it is easy to see that the equality in (1.6) holds only when S is isometric to $\mathbb{S}^2(1)$. If $k = 1$, then the equality in (1.6) holds on $M = (\mathbb{R} \times \mathbb{S}^2(1))/\mathbb{Z}_2$, where $\mathbb{Z}_2 = \langle 1, \sigma \rangle$, and $\sigma : \mathbb{R} \times \mathbb{S}^2(1) \rightarrow \mathbb{R} \times \mathbb{S}^2(1)$, $\sigma(r, s) = (-r, \tau(s))$, where τ is the antipodal map on $\mathbb{S}^2(1)$.

(1) of Theorem 1.6 can be proved by the same ideas in [6], while (2) of Theorem 1.6 needs some new observations. In the proof of (2) in Theorem 1.6, we first prove the following proposition, which generalizes the famous result of Calabi [2] and Yau [25].

Proposition 1.9. *Let (M^n, g) be an n -dimensional complete noncompact Riemannian manifold. Given $q \in M$, denote by $r(x) = d(x, q)$. Suppose*

$$(1.7) \quad \text{Ric}_g(x) \geq -\frac{1}{r^{3+\alpha}(x)},$$

where $\alpha > 0$. Then there exists a constant $c > 0$ such that

$$(1.8) \quad \liminf_{r \rightarrow \infty} \frac{\text{Vol}(B_r(q))}{r} > c.$$

In the proof of Proposition 1.9, we utilize the method presented by Sormani in [18].

Now we sketch a proof of (1.6) under the assumptions that M has nonnegative Ricci curvature and $\text{Sc}_g \geq 2$ **everywhere**, and $k = 1$. Firstly, we choose a sequence of points p_i lying on a ray with p_i going to infinity. Then by Cheeger-Colding's theory, up to a subsequence, (M, d_g, p_i) converges to a Ricci limit space (X, d_X) , which is isometric to $(\mathbb{R} \times Y, d_{\text{Eucl}} \otimes d_Y)$ for some metric space (Y, d_Y) . We need to understand more properties of Y . The properties of noncollapsed Ricci limit spaces coming from manifolds with a uniformly positive scalar curvature lower bound was studied in the recent papers [22] [28] etc. The limit spaces which are studied by [22] [28] are assumed to be noncollapsed everywhere, i.e. the volumes of **any** unit ball have a uniformly positive lower bound. In addition, according to the proof in [28], using the theory of Ricci flow coming from 3-manifolds with possibly unbounded curvature (see [17] and [1] etc.), it can be shown that if the above X is noncollapsed everywhere, then the above $(Y, d_Y, \mathcal{H}^2)$ is in fact a

non-collapsed $\text{RCD}(1, 2)$ space (in the sense of [7]), then we have an optimal upper bound of $\mathcal{H}^2(Y)$, which is a key in the proof of (1.6).

In the proof of (1.6), we prove the following proposition:

Proposition 1.10. *Under the assumption of (2) in Theorem 1.6, there is $\delta > 0$ such that for all $q \in M$,*

$$(1.9) \quad \text{Vol}(B(q, 1)) > \delta.$$

Proposition 1.10 enables us to apply the theory of Ricci flow, as what is done in [28]. Our proof of (1.6) under the general assumptions in Theorem 1.6 still uses Ricci flow. Because lower bound of Ricci curvature may be negative some where and the positive lower bound of scalar curvature is not global, we must make some adjustment to the proof in [28].

Remark 1.11. It seems that the decay condition $f(r) = \frac{1}{r^{3+\alpha}}$ is not optimal to ensure the validity of (1.6). It is an interesting question to find other more general decay conditions.

Remark 1.12. The results of this paper are part of the second author's thesis. During the process of writing this paper, Wei, Xu and Zhang have posted their paper [23] on arXiv. In the independent work [23], the sharp volume ratio at infinity has also been obtained on manifolds with $\text{Ric} \geq 0$ and $\text{Sc} \geq 2$. The methods used in [23] and our paper are different. In addition, the corresponding rigidity problem and examples are discussed in [23], while we consider manifolds with asymptotically nonnegative Ricci curvature in the present paper.

In the following we give the outline of this paper. Section 2 contains some Preliminaries on distance functions and Busemann functions. In Section 3, we prove Proposition 1.9. In Section 4, we give the proof of part (1) of Theorem 1.6. In Section 5, we prove Proposition 1.10. In Section 6, we finish the proof of part (2) of Theorem 1.6.

Acknowledgments.

The authors would like to thank Profs. Hongzhi Huang and Shaochuang Huang for discussions. The authors are partially supported by National Key R&D Program of China(2021YFA1002100) and National Natural Science Foundation of China (Nos. 12271531 and 12025109).

2. PRELIMINARIES

The manifold we are considering in this section is always complete and noncompact. All geodesics are parametrized by arclength.

A geodesic $\gamma : (-\infty, \infty) \rightarrow M$ is called a ray if for any interval $[a, b] \subset (-\infty, \infty)$, $\gamma|_{[a, b]}$ is a minimizing geodesic. A ray is a half-line $\gamma : [0, \infty) \rightarrow M$.

Given a ray γ , for any $t \geq 0$, denote by $b_t(x) := t - d(x, \gamma_t)$. It is easy to check that, given any $x \in M$, the function $t \mapsto b_t(x)$ is non-decreasing, and $b_t(x) \leq d(x, p)$ for all $t \geq 0$, hence

$$(2.1) \quad b_\gamma(x) = \lim_{t \rightarrow \infty} b_t(x)$$

is well-defined, which is called the Busemann function associated to γ , and is abbreviated as $b(x)$ if there is no confusion on γ .

A ray γ_x emanating from x is called a Busemann ray associated with γ if it is the limit of a sequence of minimal geodesics, σ_i , from x to $\gamma(R_i)$ in the following sense,

$$(2.2) \quad \gamma'_x(0) = \lim_{R_i \rightarrow \infty} \sigma'_i(0).$$

We reparametrize γ_x by arc length so that $\gamma_x(b(x)) = x$.

Let K be a compact set contained in M . Then we define

$$(2.3) \quad \Omega(K) = \{x | \exists z \in K, \exists t \geq b(z), \exists \gamma_z \text{ such that } x = \gamma_z(t)\}.$$

Furthermore, let $\Omega_R(K) = \Omega(K) \cap b^{-1}(R)$. It is not hard to check that $\Omega(K)$ is closed and $\Omega_R(K)$ is compact. The following lemma shows that the Busemann function can be regarded as a distance function.

Lemma 2.1 (Lemma 6 in [18]). *Fix $r < R$ and K be a subset of $b^{-1}((-\infty, r))$. Then*

$$(2.4) \quad d(x, b^{-1}(R)) = R - b(x), \quad \forall x \in b^{-1}((-\infty, R]).$$

Furthermore, if $z \in \Omega(K) \cap b^{-1}([r, R])$, then there is only one point, $y \in b^{-1}(R)$, such that $d(z, y) = d(z, b^{-1}(R))$. Thus $y = \gamma_z(R)$ and

$$(2.5) \quad d(z, \Omega_R(K)) = R - b(z), \quad \forall z \in \Omega(K) \cap b^{-1}([r, R]).$$

Now we are proving that under certain conditions, the distance function and the Busemann function can control each other.

Lemma 2.2. *Let γ be a ray emitting from $p \in M$ ($\gamma(0) = p$), and let $b(x)$ be the Busemann function with respect to γ . Then*

(a) $b^{-1}(R) \subset \overline{M \setminus B(p, R)}$, $\forall R > 0$;

(b) Let E be an end of M such that $\gamma(t) \in E$ for every sufficiently large t . Then for every sufficiently large $R > 0$, $b^{-1}(R) \subset E$. In addition, if there exists a positive constant $C > 0$ such that $\text{diam}(E \cap \partial B(p, r)) < C$ for every sufficiently large $r > 0$, then for any sufficiently large R it holds that

$$(2.6) \quad b^{-1}(R) \subset \overline{E \cap B(p, R + C) \setminus B(p, R)},$$

and in particular,

$$(2.7) \quad \text{diam}(b^{-1}(R)) \leq 3C.$$

Proof. For (a), suppose there exists a point $x \in b^{-1}(R) \cap B(p, R)$, then $d(p, x) < R$. Since $b(p) = 0$, by Lemma 2.1 we know that

$$(2.8) \quad d(p, b^{-1}(R)) = R,$$

which leads to

$$(2.9) \quad R > d(p, x) \geq d(p, b^{-1}(R)) = R,$$

resulting in a contradiction.

In the following we prove (b).

Claim 1. Given any R . Then any $x, y \in b^{-1}(R)$ can be connected by a curve contained in $b^{-1}([R, \infty))$.

Proof of Claim 1. We only need to prove that any $x \in b^{-1}(R)$ can be connected to $\gamma(R + 1)$ by a curve $\sigma \subset b^{-1}([R, \infty))$. Firstly, $\gamma_x(t) \subset b^{-1}([R, \infty))$ connects x and $y = \gamma_x(R + 1)$. Since $R + 1 = b(y) = \lim_{t \rightarrow \infty} b_t(y)$, we choose $\tilde{t}$ sufficiently large such that $b_{\tilde{t}}(y) = \tilde{t} - d(y, \gamma(\tilde{t})) > R + \frac{1}{2}$. Let $\tilde{\sigma} : [0, d(y, \gamma(\tilde{t}))] \rightarrow M$ be a unit speed geodesic connecting y to $\gamma(\tilde{t})$. Then for any $r \in [0, d(y, \gamma(\tilde{t}))]$, we have $b(\tilde{\sigma}(r)) \geq b_{\tilde{t}}(\tilde{\sigma}(r)) = b_{\tilde{t}}(y) + r > R + \frac{1}{2}$. Hence $\tilde{\sigma} \subset b^{-1}([R, \infty))$. Finally, γ itself is a curve connecting $\gamma(\tilde{t})$ and

$\gamma(R+1)$. Glue the above curves to form a desired curve and this completes the proof of Claim 1.

Now we assume R is large enough so that a unbounded connected component of $M \setminus B(p, R)$, denoted by N , is contained in E . According to (a) and Claim 1, every point in $b^{-1}(R)$ can be connected to $\gamma(R)$ by a curve contained in $M \setminus B(p, R)$. Since $\gamma(R) \in N$, we have $b^{-1}(R) \subset N \subset E$.

Claim 2. Suppose $R > 0$ is sufficiently large, then $E \cap \partial B(p, R) \subset b^{-1}([R-C, R])$.

Proof of Claim 2. By (a), we have $\partial B(p, R) \subset b^{-1}((-\infty, R])$. Suppose the claim does not hold, then there exists $S < R-C$ such that $E \cap \partial B(p, R) \cap b^{-1}(S) \neq \emptyset$. Take $y \in E \cap \partial B(p, R) \cap b^{-1}(S)$. Since $\text{diam}(E \cap \partial B(p, R)) < C$ and $\gamma(R) \in \partial B(p, R) \cap b^{-1}(R)$, we know that

$$(2.10) \quad d(y, b^{-1}(R)) \leq d(y, \gamma(R)) < C.$$

On the other hand, by Lemma 2.1 we know that $d(y, b^{-1}(R)) = R - S > C$, which leads to a contradiction. This completes the proof of Claim 2.

Now, let's prove that $\sup_{x \in b^{-1}(R)} d(p, x) \leq R + C$. Suppose the conclusion does not hold,

then there exists a point $z \in b^{-1}(R) \subset E$ and $\varepsilon > 0$ such that $d(p, z) = R + C + \varepsilon$. Hence $E \cap \partial B(p, R + C + \varepsilon) \cap b^{-1}(R) \neq \emptyset$. On the other hand, according to Claim 2, we have $E \cap \partial B(p, R + C + \varepsilon) \subset b^{-1}([R + \varepsilon, R + C + \varepsilon])$, which is a contradiction. Therefore, $b^{-1}(R) \subset E \cap \overline{B(p, R + C) \setminus B(p, R)}$. Hence,

$$(2.11) \quad \begin{aligned} \text{diam}(b^{-1}(R)) &\leq \text{diam}(E \cap \overline{B(p, R + C) \setminus B(p, R)}) \\ &\leq C + \text{diam}(E \cap \partial B(p, R)) + \text{diam}(E \cap \partial B(p, R + C)) \\ &< 3C. \end{aligned}$$

The proof is completed. □

Given any $\delta > 0$ and any $r_1 < r_2 < R$, let

$$(2.12) \quad S_{\delta, r_1, r_2} = S_{\delta, r_1, r_2}(\Omega_R(K))$$

be the set of points x with $d(x, \Omega_R(K)) \in [R - r_2, R - r_1]$ such that there exists a minimal geodesic σ from $\Omega_R(K)$ to x with

$$(2.13) \quad \text{Length}(\sigma) = d(x, \Omega_R(K)) \quad \text{and} \quad g(\sigma'(0), -\nabla b) \geq 1 - \delta.$$

In the following we collect some notions and results from [18].

Lemma 2.3 (Lemma 8 and Corollary 9 in [18]). *Fix $R > r_2 > r_1$, $\delta > 0$ and K be a compact subset of $b^{-1}((-\infty, r_1))$. Then*

$$(2.14) \quad \bigcap_{\delta > 0} S_{\delta, r_1, r_2}(\Omega_R(K)) = \Omega(K) \cap b^{-1}([r_1, r_2])$$

and

$$(2.15) \quad \lim_{\delta \rightarrow 0} \text{Vol}(S_{\delta, r_1, r_2}(\Omega_R(K))) = \text{Vol}(\Omega(K) \cap b^{-1}([r_1, r_2])).$$

Lemma 2.4 (Lemma 10 in [18]). *Let $K \subset b^{-1}((-\infty, r_1])$ and $R > r_3 > r_2 > r_1$. Given any $\delta > 0$, there exists*

$$(2.16) \quad h_1(\delta) = h_1(\delta, r_1, r_2, r_3, R, K, M) < \delta$$

such that for any $x \in S_{h_1(\delta), r_2, r_3}(\Omega_R(K))$, every minimal geodesic, σ , from $\sigma(0) \in \Omega_R(K)$ to x with $d(x, \sigma(0)) = d(x, \Omega_R(K))$, it holds

$$(2.17) \quad g(\sigma'(0), -\nabla b) \geq 1 - \delta.$$

Given any $\varepsilon > 0$, there exists a set

$$(2.18) \quad \Omega_R^\varepsilon(K) = \{p_1, \dots, p_{N_\varepsilon}\} \subset \Omega_R(K)$$

such that the tubular neighborhood, $T_\varepsilon(\Omega_R^\varepsilon(K))$, contains $\Omega_R(K)$. In this paper, for a set $X \subset M$ and $0 \leq a < b$, we use the notations

$$T_{a,b}(X) := \{y | a \leq d(y, X) \leq b\},$$

and

$$T_b(X) := T_{0,b}(X).$$

Given $\Omega_R^\varepsilon(K)$ and $p \in \Omega_R^\varepsilon(K)$, the star-shaped set $V_{p,\varepsilon}$ and the star-shaped wedge set $U_{p,\delta}$ are defined to be

$$(2.19) \quad V_{p,\varepsilon} = \{x | d(x, p) < d(x, q), \forall q \in \Omega_R^\varepsilon(K) \text{ such that } q \neq p\},$$

$$(2.20) \quad U_{p,\delta} = \{x | \exists \text{ a minimal geodesic } \sigma \text{ from } p \text{ to } x \text{ such that } g(\sigma'(0), -\nabla b) \geq 1 - \delta\}.$$

In addition, denote by

$$(2.21) \quad U_{\varepsilon,\delta} = \bigcup_{i=1}^{N_\varepsilon} (U_{p_i,\delta} \cap V_{p_i,\varepsilon}).$$

Lemma 2.5 (Lemma 14 in [18]). *Fix $r_1 < R$ and the compact set, $K \subset b^{-1}((-\infty, r_1])$. For all $\varepsilon > 0$, there exists $h_2(\varepsilon, r_1, R, K, M) > 0$ such that, for all $p \in \Omega_R(K)$ and all $\delta < h_2$ we have*

$$(2.22) \quad d(\gamma_x(R), p) < \varepsilon, \quad \forall x \in U_{p,\delta} \cap B(p, R - r_1 - \varepsilon).$$

Thus

$$(2.23) \quad b(x) \in [R - d(x, p), R - d(x, p) + \varepsilon], \quad \forall x \in U_{p,\delta} \cap B(p, R - r_1 - \varepsilon).$$

Lemma 2.6 (Lemma 15 in [18]). *Fix $r_1 < r_2 < R$ and $\delta > 0$. Let $\varepsilon_0 > 0$, $K \subset b^{-1}((-\infty, r_1])$, there exists $h_3(M, \delta, r_1, r_2, R) > 0$ sufficiently small that for all $\varepsilon < \min\{\varepsilon_0, h_3\}$,*

$$(2.24) \quad \Omega(K) \cap b^{-1}([r_1, r_2]) \subset T_{R-r_2-\varepsilon_0, R-r_1-\varepsilon_0}(\Omega_R^\varepsilon(K)) \cap \bar{U}_{\varepsilon,\delta}.$$

where $\bar{U}$ is the closure of U . So

$$(2.25) \quad \text{Vol}(\Omega(K) \cap b^{-1}([r_1, r_2])) \leq \text{Vol}(T_{R-r_2-\varepsilon_0, R-r_1-\varepsilon_0}(\Omega_R^\varepsilon(K)) \cap U_{\varepsilon,\delta}).$$

Lemma 2.7 (Lemma 16 in [18]). *Fix $r_1 < r_2 < r_3 < R$ and $K \subset b^{-1}((-\infty, r_1])$. Given any $\delta > 0$, let $h_1(\delta) < \delta$ be the constant defined in Lemma 2.4. Then there exists $h_4 = h_4(M, h_1(\delta), R, r_1, r_2, r_3) > 0$ such that given any $\varepsilon_0 > 0$, for all $\varepsilon < \min\{h_4, \varepsilon_0\}$, we have*

$$(2.26) \quad T_{R-r_3+\varepsilon_0, R-r_2-\varepsilon_0}(\Omega_R^\varepsilon(K)) \cap U_{\varepsilon, h_1(\delta)} \subset S_{2\delta, r_2, r_3}(\Omega_R(K)).$$

3. PROOF OF PROPOSITION 1.9

Lemma 3.1. *Let (M^n, g) be an n -dimensional complete noncompact Riemannian manifold with a given ray γ emitting from q and its associated Busemann function, $b = b_\gamma$ with $b(\gamma(t)) = t$. Suppose*

$$(3.1) \quad \text{Ric}_g(x) \geq -\frac{C}{b^{3+\alpha}(x)}, \quad \forall x \in b^{-1}((1, \infty))$$

for some $C > 0$ and $\alpha > 0$. Let $K \subset b^{-1}((0, \infty))$ be a compact set, then there exists some $k_0 > 0$ such that for any $k > k_0$, it holds

$$(3.2) \quad \text{Vol}(\Omega(K) \cap b^{-1}([k+3, k+4])) \geq \frac{1}{1 + 2^n(k+2)^{-(1+\alpha)}} \text{Vol}(\Omega(K) \cap b^{-1}([k+1, k+2])).$$

Furthermore, there exists a constant $C_1 > 0$ such that

$$(3.3) \quad \text{Vol}(\Omega(K) \cap b^{-1}([k, k+1])) > C_1$$

for any sufficiently large k .

Proof. Let

$$(3.4) \quad R > r_4 > r_3 > r_2 > r_1 > \sup_{x \in K} b^{-1}(x).$$

Set $\varepsilon_0 < \frac{1}{10} \min\{r_2 - r_1, r_4 - r_3, R - r_4\}$, by Lemma 2.5, there exists h_2 , such that if $p \in \Omega_R^\varepsilon(K)$ and $0 < \delta < h_2$, then for all $0 < h < \delta$ and $x \in U_{p,h} \cap B(p, R - r_1 - \varepsilon_0)$, we have

$$(3.5) \quad b(x) \in [R - d(x, p), R - d(x, p) + \varepsilon_0].$$

Take $h_1 < \delta$ from Lemma 2.4, then for every $x \in U_{p,h_1} \cap B(p, R - r_1 - \varepsilon_0)$, (3.5) holds, and

$$(3.6) \quad \text{Ric}_g(x) \geq -\frac{C}{(R - d(x, p))^{3+\alpha}}$$

holds for these x .

Now we construct a radially symmetric model space.

Let $J_R(t) = t(1 + (R - t)^{-(1+\alpha)})$, then

$$(3.7) \quad J'_R(t) = 1 + (R - t)^{-(1+\alpha)} + (1 + \alpha)t(R - t)^{-(2+\alpha)},$$

$$(3.8) \quad J''_R(t) = 2(1 + \alpha)(R - t)^{-(2+\alpha)} + (2 + \alpha)(1 + \alpha)t(R - t)^{-(3+\alpha)}.$$

Suppose C, t are sufficiently large, we have

$$\begin{aligned} -\frac{J''_R(t)}{J_R(t)} &= -\frac{2(1 + \alpha)(R - t)^{-(2+\alpha)} + (2 + \alpha)(1 + \alpha)t(R - t)^{-(3+\alpha)}}{t(1 + (R - t)^{-(1+\alpha)})} \\ &\leq -C(R - t)^{-(3+\alpha)}, \end{aligned}$$

and

$$(3.9) \quad (n - 2) \frac{1 - J'^2_R}{J_R^2} - \frac{J''_R}{J_R} \leq -\frac{J''_R}{J_R} \leq -C(R - t)^{-(3+\alpha)}.$$

Let $(M^J, g_0 = dt^2 + J^2 g_{\mathbb{S}^{n-1}})$, where $g_{\mathbb{S}^{n-1}}$ is the standard metric on $\mathbb{S}^{n-1}$, $t(z) = d(z \in M^J, 0 \in M^J)$. When $t \geq 1$, $J = J_R$. Therefore, for sufficiently large $C, t > 0$, we have

$$(3.10) \quad \text{Ric}_{g_0}(\partial t, \partial t) = -(n - 1) \frac{J''_R(t)}{J_R(t)} \leq -C(R - t)^{-(3+\alpha)}.$$

For the unit vector v orthogonal to ∂t , we have

$$(3.11) \quad \text{Ric}_{g_0}(v, v) = (n - 2) \frac{1 - J'^2_R}{J_R^2} - \frac{J''_R}{J_R} \leq -C(R - t)^{-(3+\alpha)}.$$

So, it holds that

$$(3.12) \quad \text{Ric}_{g_0}(z) \leq -\frac{C}{(R - t)^{3+\alpha}} \leq \text{Ric}_g(y),$$

for all $z \in B^{M^J}(0, R) \setminus B^{M^J}(0, 1)$ with $t(z) = d(z, 0)$ and $y \in U_{p,h_1} \cap B(p, R - r_1 - \varepsilon)$ with $t(y) = d(y, p)$.

Choose h_3 and h_4 from Lemmas 2.6 and 2.7 respectively. Let

$$(3.13) \quad \varepsilon < \min\{h_3, h_4, \varepsilon_0\}.$$

Set

$$(3.14) \quad V_p = V_{p,\varepsilon} \cap U_{p,h_1} \cap B(p, R - r_1 - \varepsilon_0).$$

Then, by (2.21), we have the following disjoint union,

$$(3.15) \quad \bigcup_{p \in \Omega_R^\varepsilon(K)} V_p = T_{R-r_1-\varepsilon_0}(\Omega_R^\varepsilon(K)) \cap U_{\varepsilon,h_1}.$$

Let $\text{Ann}_p(r, s) = B(p, s) \setminus B(p, r)$.

Since

$$(3.16) \quad (\text{Ann}_p(r, s) \cap V_{p_i}) \cap (\text{Ann}_p(r, s) \cap V_{p_j}) \subset V_{p_i} \cap V_{p_j} = \emptyset,$$

for $p_i, p_j \in \Omega_R^\varepsilon(K)$, $i \neq j$, and by (3.14), (3.15) we have

$$(3.17) \quad T_{r,s}(\Omega_R^\varepsilon(K)) \cap U_{\varepsilon,h_1} = \bigcup_{p \in \Omega_R^\varepsilon(K)} (\text{Ann}_p(r, s) \cap V_p).$$

Therefore,

$$(3.18) \quad \text{Vol}(T_{r,s}(\Omega_R^\varepsilon(K)) \cap U_{\varepsilon,h_1}) = \sum_{p \in \Omega_R^\varepsilon(K)} \text{Vol}(\text{Ann}_p(r, s) \cap V_p).$$

Imitating [3] to apply the volume comparison theorem to starlike set, we obtain

$$(3.19) \quad \frac{\text{Vol}(\text{Ann}_p(R - r_4, R - r_3) \cap V_p)}{\text{Vol}(\text{Ann}_p(R - r_2, R - r_1 - \varepsilon_0) \cap V_p)} \geq \frac{V(r_3, r_4, R)}{V(r_1 + \varepsilon_0, r_2, R)},$$

where

$$(3.20) \quad V(s_1, s_2, R) = \int_{R-s_2}^{R-s_1} J_R^{n-1} dt.$$

Thus,

$$(3.21) \quad \begin{aligned} & \frac{\text{Vol}(T_{R-r_4, R-r_3}(\Omega_R^\varepsilon(K)) \cap U_{\varepsilon,h_1})}{\text{Vol}(T_{R-r_2, R-r_1-\varepsilon_0}(\Omega_R^\varepsilon(K)) \cap U_{\varepsilon,h_1})} \\ &= \frac{\sum_{p \in \Omega_R^\varepsilon(K)} \text{Vol}(\text{Ann}_p(R - r_4, R - r_3) \cap V_p)}{\sum_{p \in \Omega_R^\varepsilon(K)} \text{Vol}(\text{Ann}_p(R - r_2, R - r_1 - \varepsilon_0) \cap V_p)} \\ &\geq \frac{\sum_{p \in \Omega_R^\varepsilon(K)} \text{Vol}(\text{Ann}_p(R - r_2, R - r_1 - \varepsilon_0) \cap V_p) \frac{V(r_3, r_4, R)}{V(r_1 + \varepsilon_0, r_2, R)}}{\sum_{p \in \Omega_R^\varepsilon(K)} \text{Vol}(\text{Ann}_p(R - r_2, R - r_1 - \varepsilon_0) \cap V_p)} \\ &= \frac{V(r_3, r_4, R)}{V(r_1 + \varepsilon_0, r_2, R)}. \end{aligned}$$

We can now apply Lemma 2.6 and Lemma 2.7, therefore,

$$(3.22) \quad \frac{\text{Vol}(S_{2\delta, r_3-\varepsilon_0, r_4+\varepsilon_0}(\Omega_R(K)))}{\text{Vol}(\Omega_R(K) \cap b^{-1}([r_1 + 2\varepsilon_0, r_2 - \varepsilon_0]))} \geq \frac{\text{Vol}(T_{R-r_4, R-r_3}(\Omega_R^\varepsilon(K)) \cap U_{\varepsilon,h_1})}{\text{Vol}(T_{R-r_2, R-r_1-\varepsilon_0}(\Omega_R^\varepsilon(K)) \cap U_{\varepsilon,h_1})}.$$

Thus combining this inequality with (3.21) means that

$$(3.23) \quad \frac{\text{Vol}(S_{2\delta, r_3-\varepsilon_0, r_4+\varepsilon_0}(\Omega_R(K)))}{\text{Vol}(\Omega_R(K) \cap b^{-1}([r_1 + 2\varepsilon_0, r_2 - \varepsilon_0]))} \geq \frac{V(r_3, r_4, R)}{V(r_1 + \varepsilon_0, r_2, R)}.$$

Note that (3.23) does not depend on h_1 or ε .

Taking the limit as $\delta \rightarrow 0$ and applying (2.15), we obtain

$$(3.24) \quad \frac{\text{Vol}(\Omega(K) \cap b^{-1}([r_3 - \varepsilon_0, r_4 + \varepsilon_0]))}{\text{Vol}(\Omega(K) \cap b^{-1}([r_1 + \varepsilon_0, r_2 - \varepsilon_0]))} \geq \frac{V(r_3, r_4, R)}{V(r_1 + \varepsilon_0, r_2, R)}.$$

From (3.24) and (3.20) we have

$$\begin{aligned}
(3.25) \quad & \frac{\text{Vol}(\Omega(K) \cap b^{-1}([r_3 - \varepsilon_0, r_4 + \varepsilon_0]))}{\text{Vol}(\Omega(K) \cap b^{-1}([r_1 + \varepsilon_0, r_2 - \varepsilon_0]))} \\
& \geq \frac{V(r_3, r_4, R)}{V(r_1 + \varepsilon_0, r_2, R)} \\
& = \frac{\int_{R-r_4}^{R-r_3} t^{n-1} (1 + (R-t)^{-(1+\alpha)})^{n-1} dt}{\int_{R-r_2}^{R-r_1-\varepsilon_0} t^{n-1} (1 + (R-t)^{-(1+\alpha)})^{n-1} dt} \\
& \geq \frac{\int_{R-r_4}^{R-r_3} t^{n-1} dt}{\int_{R-r_2}^{R-r_1-\varepsilon_0} t^{n-1} (1 + 2^n (R-t)^{-(1+\alpha)}) dt} \\
& = \frac{(R-r_3)^n - (R-r_4)^n}{(R-r_1-\varepsilon_0)^n - (R-r_2)^n + n2^n \int_{R-r_2}^{R-r_1-\varepsilon_0} t^n (R-t)^{-(1+\alpha)} dt},
\end{aligned}$$

where

$$\begin{aligned}
(3.26) \quad & \int_{R-r_2}^{R-r_1-\varepsilon_0} t^{n-1} (R-t)^{-(1+\alpha)} dt \\
& \leq r_2^{-(1+\alpha)} \int_{R-r_2}^{R-r_1-\varepsilon_0} t^{n-1} dt \\
& = \frac{r_2^{-(1+\alpha)}}{n} ((R-r_1-\varepsilon_0)^n - (R-r_2)^n).
\end{aligned}$$

Therefore,

$$\begin{aligned}
(3.27) \quad & \frac{\text{Vol}(\Omega(K) \cap b^{-1}([r_3 - \varepsilon_0, r_4 + \varepsilon_0]))}{\text{Vol}(\Omega(K) \cap b^{-1}([r_1 + \varepsilon_0, r_2 - \varepsilon_0]))} \\
& \geq \frac{(R-r_3)^n - (R-r_4)^n}{((R-r_1-\varepsilon_0)^n - (R-r_2)^n)(1 + 2^n r_2^{-(1+\alpha)})}.
\end{aligned}$$

Let $r_i = k + i$, $R \rightarrow +\infty$, and $\varepsilon_0 \rightarrow 0$ we have

$$\begin{aligned}
(3.28) \quad & \frac{\text{Vol}(\Omega(K) \cap b^{-1}([k+3, k+4]))}{\text{Vol}(\Omega(K) \cap b^{-1}([k+1, k+2]))} \\
& \geq \lim_{R \rightarrow +\infty} \frac{(R-k-3)^n - (R-k-4)^n}{((R-k-1)^n - (R-k-2)^n)(1 + 2^n (k+2)^{-(1+\alpha)})} \\
& = \lim_{R \rightarrow +\infty} \frac{nR^{n-1}(k+4) - nR^{n-1}(k+3) + O(R^{n-2})}{(nR^{n-1}(k+2) - nR^{n-1}(k+1) + O(R^{n-2}))(1 + 2^n (k+2)^{-(1+\alpha)})} \\
& = \frac{1}{1 + 2^n (k+2)^{-(1+\alpha)}}.
\end{aligned}$$

Now we go to prove (3.3). For every $k \in \mathbb{Z}^+$, denote by $B_k = \ln(\text{Vol}(\Omega(K) \cap b^{-1}([k+3, k+4])))$. According to (3.2), for $k \geq k_0 + 4$, we have

$$\begin{aligned}
(3.29) \quad & B_k \geq B_{k-2} - \ln\left(1 + \frac{2^n}{(k-2)^{1+\alpha}}\right) \\
& \geq B_{k-2} - \frac{2^n}{(k-2)^{1+\alpha}},
\end{aligned}$$

where we use the basic inequality $\ln(1+x) \leq x$ for every $x \geq 0$. By induction, we have

$$(3.30) \quad B_k > \min\{B_{k_0+1}, B_{k_0}\} - \sum_{k_0}^{k-2} \frac{2^n}{(k-2)^{1+\alpha}}.$$

Since for every $\alpha > 0$, the series $\sum_{i=1}^{\infty} \frac{1}{i^{1+\alpha}}$ converges, whose sum is denoted by C_α , we have

$$B_k > \min\{B_{k_0+1}, B_{k_0}\} - 2^n C_\alpha.$$

Hence (3.3) holds for $k \geq k_0$. The proof is completed. $\square$

Now we give the proof of Proposition 1.9.

Proof. Given a ray γ emitting from q and $b(x)$ denotes its associated Busemann function. By (a) of Lemma 2.2 we know that $r(x) \geq b(x)$ for $x \in b^{-1}((0, +\infty))$, hence

$$(3.31) \quad \text{Ric}_g(x) \geq -\frac{1}{r^{3+\alpha}(x)} \geq -\frac{1}{b^{3+\alpha}(x)}.$$

According to Lemma 3.1, there exist $c_0 > 0$ and $k_0 > 0$ such that

$$(3.32) \quad \text{Vol}(\Omega(K) \cap b^{-1}([k, k+1])) > c_0$$

for any $k > k_0$.

Set $K = \bar{B}(q, 1)$, then for any $y \in \Omega(K) \cap b^{-1}([k_0, k_0 + r])$, there exists $y' \in K$ such that $y = \gamma_{y'} \cap b^{-1}(t)$, for some $t \in [k_0, k_0 + r]$. Therefore $d(y, y') \leq k_0 + 1 + r$, and hence $d(y, q) \leq k_0 + 2 + r$, and

$$(3.33) \quad \Omega(K) \cap b^{-1}([k_0, k_0 + r]) \subset B(q, k_0 + 2 + r).$$

Combine (3.33) with (3.32), it holds that

$$(3.34) \quad \text{Vol}(B(q, k_0 + 2 + r)) \geq \text{Vol}(\Omega(K) \cap b^{-1}([k_0, k_0 + r])) \geq c_0(r - 1).$$

For sufficiently large r , we have

$$(3.35) \quad \text{Vol}(B(q, r)) > \frac{c_0}{2}r.$$

The proof is completed. $\square$

4. PART (1) OF THEOREM 1.6

Recall the following basic lemma, which is a consequence of the theory of μ -bubbles due to Gromov [10].

Lemma 4.1 (Lemma 2.1 in [6]). *Let (N^3, g) be a 3-manifold with boundary satisfying $\text{Sc}_g \geq 1$. Then there are universal constants $L > 0$ and $c > 0$ such that if there is a $p \in N$ with $d_N(p, \partial N) > L/2$, then there is an open set $\Omega \subset B(\partial N, L/2) \cap N$ and a smooth surface Σ^2 such that $\partial\Omega = \Sigma \sqcup \partial N$ and each component of Σ has diameter at most c .*

In the following we fix the positive constants L and c so that $L \gg c$ and they satisfies the conclusion in Lemma 4.1.

Lemma 4.2. *Let (M^3, g) be a complete noncompact 3-dimensional manifold with $\text{Sc}_g \geq 2$ outside a compact set K . Suppose that M has finitely many ends and its first Betti number $b_1(M) < \infty$. Fix a point $p \in M$, $r(x) = d(x, p)$. Suppose there is a function $f : \mathbb{R}^+ \rightarrow \mathbb{R}^+$, such that $\lim_{r \rightarrow \infty} f(r) = 0$ and*

$$(4.1) \quad \text{Ric}_g(x) \geq -f(r(x)).$$

Then there is an $r_0(p, M, g, f) > 0$ and a universal constant $C > 0$ so that the following holds. For every end $E \subset M$ such that E contains a unbounded connected component of $M \setminus B(p, r_0)$, any $r' > r_0$, and $a_1, a_2 \in [L, 2L]$, let E_1 and E_2 be the unbounded components of $E \setminus \bar{B}(p, r' + a_1)$ and $E \setminus \bar{B}(p, r' + a_1 + a_2)$ respectively, then $\bar{E}_1 \setminus E_2$ consists of one connected component, and

$$(4.2) \quad \text{diam}(\bar{E}_1 \setminus E_2) \leq C.$$

In particular, we have $\text{diam}(\partial E_1) \leq C$.

Lemma 4.2 slightly generalizes the results in [6], and its proof follows the same line as that of [6]. We give a sketch of its proof as follows.

Sketch of the proof of Lemma 4.2: Let E_0 be the unbounded component of $E \setminus \bar{B}(p, r')$. Since $b_1(M)$ is finite, we can prove that there is a $r_0(p, M, g) > 0$ such that ∂E_k , ($k = 0, 1, 2$) are connected when $r' > r_0$, see Proposition 3.2 in [5] for a proof. In particular, we conclude that $\bar{E}_1 \setminus E_2$ consists of one connected component. Furthermore, we assume $r_0(p, M, g)$ is large so that $\text{Sc}_g \geq 2$ on $M \setminus B(p, r_0)$.

Let γ be a ray emitting from p so that $\gamma(t) \in E$ for every large t . Then for all $r' > r_0$, $\gamma \cap \partial B(p, r')$ lies on the boundary of the unique unbounded component of $E \setminus B(p, r')$. We apply Lemma 4.1 to E_k for $k = 0, 1$. According to Lemma 5.4 in [5], we obtain a connected surface Σ_k in $B(\partial E_k, L/2) \cap E_k$ with $\text{diam}(\Sigma_k) \leq c$ that separates ∂E_k from E . Take $t_k \in \mathbb{R}_+$ with $\gamma(t_k) \in \Sigma_k$ for $k = 0, 1$. Since $d(\gamma(t_0), \gamma(t_1)) \leq a_1 + a_2 \leq 4L$, we have $d_g(x_0, x_1) \leq 4L + 2c$ for any $x_0 \in \Sigma_0$ and $x_1 \in \Sigma_1$.

As in [6], denote by $D := 4L + 2c$, $b := 3c$, $A := 2\sqrt{b^2 + D^2}$. Fix $R \in \mathbb{R}$ such that

$$(4.3) \quad R \geq A + 4L, \quad \sqrt{b^2 + (2R + D)^2} + 1 < 2D + 2R.$$

Then take $\delta \in (0, 1)$ small enough such that

$$(4.4) \quad \delta < \sqrt{b^2 + D^2}, \quad 14\delta + 6\sqrt{\delta(D + R)} < c, \quad 2c + 22\delta + 6\sqrt{\delta(D + R)} < \frac{L}{2}.$$

All the constants above are independent of (M, g) and r_0 .

Since (4.1) holds for $\lim_{r \rightarrow \infty} f(r) = 0$, by Cheeger-Colding's almost splitting theorem ([4]), assuming $r_0(D, R, \delta, f)$ is sufficiently large, there is a length space (Y, d_Y) with

$$(4.5) \quad d_{GH}(B(\gamma(t_1), D + R) \subset (M, d_g), B((y, 0), D + R) \subset (Y \times \mathbb{R}, d_Y \otimes d_{\text{Eucl}})) < \delta.$$

The above is the only one occasion where the almost nonnegative Ricci curvature condition is used. Once we have obtained (4.5), together with the setting of constants in (4.3) and (4.4) and their geometric meanings, the remaining argument in the proof of Claims 1, 2, 3 in [6] can be carried out. In particular, we can prove that, $\bar{E}_1 \setminus E_2$ is contained in the tubular neighborhood $T_A(\gamma|_{|t-t_1| < D+R})$ (this is just Claim 3 in [6]). Therefore, we have

$$(4.6) \quad \text{diam}(\bar{E}_1 \setminus E_2) \leq 2A + 2D + 2R.$$

This finishes the proof. $\square$

With Lemma 4.2, we can finish the proof of (1) of Theorem 1.6.

Proof of (1) in Theorem 1.6. Let $E^{(1)}, \dots, E^{(k)}$ be the unbounded connected components of $M \setminus \bar{B}(p, r_0 + 2L)$, where r_0 is from Lemma 4.2.

For each $s \in \mathbb{Z}^+$, let $E_s^{(i)}$ denote the unbounded connected components of $E^{(i)} \setminus \bar{B}(p, r_0 + sL)$. Then for every r with $r_0 + (s+1)L \leq r < r_0 + (s+2)L$, we have

$$(4.7) \quad B(p, r) \subset (M \setminus \bigcup_{i=1}^k E_1^{(i)}) \bigcup (\bigcup_{i=1}^k \bigcup_{j=2}^s (E_{j-1}^{(i)} \setminus E_j^{(i)}))$$

By Lemma 4.2 and the Bishop-Gromov volume comparison theorem, it holds that

$$(4.8) \quad \text{Vol}(\bar{E}_i \setminus E_{i+1}) < \tilde{C}.$$

Thus,

$$(4.9) \quad \begin{aligned} \text{Vol}(B(p, r)) &\leq \text{Vol}(M \setminus \bigcup_{i=1}^k E_1^{(i)}) + k\tilde{C}(s-1) \\ &\leq \text{Vol}(M \setminus \bigcup_{i=1}^k E_1^{(i)}) + k\tilde{C}(\frac{1}{L}(r - r_0) - 2). \end{aligned}$$

i.e. (M, g) has at most linear volume growth.

In the next we prove the macroscopic dimension of M is 1.

For simplicity of notation, we assume E is one of $E^{(1)}, \dots, E^{(k)}$. For every $R \geq r_0 + 2L$, $E \cap \partial B(p, R)$ may consists of many connected components, which is denoted by $A_0, A_1, \dots, A_\alpha$. We assume A_0 is the (unique) boundary component of $E \setminus B(p, r)$. Then by Lemma 4.2, $\text{diam}(A_0) \leq C$. For $i \in \{1, \dots, \alpha\}$, let

$$\mu_i := \inf\{s | A_i \text{ is contained in the unbounded component of } E \setminus B(p, s)\}.$$

Then $\mu_i \geq r_0 + 2L$. We apply Lemma 4.2 to $r' = \mu_i - 1 - L$, $a_1 = a_2 = L$ (recall that $L \gg 1$) to conclude that every two points in A_i and ∂B_i respectively has a distant at most C , and

$$\text{diam}(\partial B_i) \leq C.$$

where B_i denotes the unbounded components of $E \setminus \bar{B}(p, \mu_i - 1)$. Then we have $\mu_i \geq R - C + 1$. Let $i_0 \in \{1, \dots, \alpha\}$ is one such that μ_{i_0} is the minimal one of $\{\mu_1, \dots, \mu_\alpha\}$. From the construction it is easy to see that every point $x \in A_i$ can be connected by a geodesic to a point $y \in \partial B_{i_0}$, and $d(x, y) \leq C$. On the other hand, basing on Lemma 4.2, by an induction argument we can prove that every two points in A_0 and ∂B_{i_0} respectively has a distance at most $(\frac{C}{L} + 1)C$. Thus we conclude that for every $R \geq r_0 + 2L$, $E \cap \partial B(p, R) = \bigcup_{i=0}^\alpha A_i$ satisfies

$$(4.10) \quad \text{diam}(E \cap \partial B(p, R)) \leq C_2 := (\frac{C}{L} + 2)C.$$

Since $M \setminus \bigcup_{i=1}^k E^{(i)}$ is compact, it is obvious that there exists R_0 such that for $R \geq R_0$, every connect component of $\partial B(p, R)$ is contained in $\bigcup_{i=1}^k E^{(i)}$.

Let P be the radial rays obtained by gluing the end points of k copies of rays $[R_0, \infty)$. Let y_0 denote the center point of P . Define a map $h : M \rightarrow P$ as follows. For $x \in B(p, R_0)$, $h(x) = y_0$; for $x \in E^{(i)} \cap \partial B(p, R)$, where $R > R_0$, x is mapped by h to a point in the i -th copy of $[R_0, \infty)$ correspond to $d(x, p)$. It is easy to see that there is a constant $C_3 > 0$ such that $\text{diam}(h^{-1}(y)) \leq C_3$ holds for every $y \in P$. Since M is noncompact, we conclude that the macroscopic dimension of M is 1. $\square$

Corollary 4.3. *Under the assumption of (1) in Theorem 1.6, let $\gamma(t)$ be a ray emanating from $p \in M$, and let $b(x)$ be the Busemann function with respect to $\gamma(t)$, then there exists a large constant $R_0 > 0$ such that*

$$(4.11) \quad \text{diam}(b^{-1}(R)) \leq 3C_2, \quad \forall R \geq R_0.$$

Proof. Let E be the end of M such that $\gamma(t) \in E$ for every large t . Recall that we have proved (4.10) for every sufficiently large R . Then by (b) of Lemma 2.2, (4.11) holds. $\square$

5. PROOF OF PROPOSITION 1.10

Proof of Proposition 1.10. It suffices to prove (1.9) holds for every $q \in E$, where E is any fixed end of M . Consider a ray $\gamma(t)$ emanating from p , such that $\gamma(t) \in E$ for every large t , and let $b(x)$ denote the Busemann function with respect to $\gamma(t)$.

Now, suppose there exists a sequence of points $q_i \in E$ such that $\text{Vol}(B(q_i, 1)) \rightarrow 0$ as $i \rightarrow \infty$. Let $d(p, q_i) = R_i$, then $R_i \rightarrow \infty$. Since

$$(5.1) \quad d(b^{-1}(R_k), b^{-1}(R_k + 1)) = 1,$$

for every sufficiently large k , we have

$$(5.2) \quad \begin{aligned} & \text{diam}(b^{-1}([R_k, R_k + 1])) \\ & \leq \text{diam}(b^{-1}(R_k)) + \text{diam}(b^{-1}(R_{k+1})) + d(b^{-1}(R_k), b^{-1}(R_k + 1)) \\ & < 6C_2 + 1, \end{aligned}$$

where we use (4.11) in the last inequality.

On the other hand, combining (4.10) with the facts that $q_k \in \partial B(p, R_k)$, $\partial B(p, R_k) \cap b^{-1}([R_k, R_k + 1]) \neq \emptyset$, we have

$$(5.3) \quad b^{-1}([R_k, R_k + 1]) \subset B(q_k, 7C_2 + 1).$$

Because the Ricci curvature of the manifold is asymptotically nonnegative, there exists $K \geq 0$ such that $\text{Ric}_g \geq -(n-1)K$. By the Bishop-Gromov volume comparison theorem, there exists a constant $\tilde{C}$ depending only on C_2 and K such that

$$(5.4) \quad \begin{aligned} & \text{Vol}(B(q_k, 1)) \\ & \geq \tilde{C} \text{Vol}(B(q_k, 7C_2 + 1)) \\ & \geq \tilde{C} \text{Vol}(b^{-1}([R_k, R_k + 1])). \end{aligned}$$

By Lemma 3.1,

$$(5.5) \quad \text{Vol}(b^{-1}([R_k, R_k + 1])) > C_1 > 0,$$

holds for every sufficiently large k , then

$$(5.6) \quad \text{Vol}(B(q_k, 1)) \geq \tilde{C}C_1,$$

leading to a contradiction. The proof is completed. $\square$

6. PART (2) OF THEOREM 1.6

Firstly, let's recall some results on Ricci flow. The following theorem is proved in [1] and [17].

Theorem 6.1. *Given any $\alpha_0 > 0$, $v_0 > 0$, there exist positive constants $T = T(\alpha_0, v_0)$, $C = C(v_0)$ such that the following holds. Let (M, g) be a 3-dimensional Riemannian manifold, $B_g(x_0, s) \subset\subset M$ for some $s \geq 4$. Suppose*

$$(6.1) \quad \text{Ric}_g \geq -\alpha_0 \quad \text{on } B_g(x_0, s),$$

$$(6.2) \quad \text{Vol}(B_g(x, 1)) \geq v_0 \quad \text{for all } x \in B_g(x_0, s-1).$$

Then there exists a Ricci flow $(g(t))_{t \in [0, T]}$ on $B_g(x_0, s-2)$, with $g(0) = g$, so that

$$(6.3) \quad \text{Ric}_{g(t)} \geq -C\alpha_0 \quad \text{on } B_g(x_0, s-2),$$

$$(6.4) \quad |\text{Rm}_{g(t)}| \leq \frac{C}{t} \quad \text{on } B_g(x_0, s-2),$$

for all $t \in (0, T]$.

Recall the shrinking balls lemma and the expanding balls lemma from [16] as follows.

Lemma 6.2 (The shrinking balls lemma, Corollary 3.3 in [16]). *Suppose $(M, g(t))_{t \in [0, T]}$ is a Ricci flow on a manifold M of any dimension n . Then there exists $\beta = \beta(n) \geq 1$ such that the following is true. Suppose $x_0 \in M$ and that $B_{g(0)}(x_0, r) \subset\subset M$ for some $r > 0$, and $|\text{Rm}|_{g(t)} \leq \frac{c_0}{t}$ and some $c_0 > 0$. Then for any $0 \leq s \leq t \leq T$, we have*

$$(6.5) \quad B_{g(0)}(x_0, r) \supset B_{g(s)}(x_0, r - \beta\sqrt{c_0 s}) \supset B_{g(t)}(x_0, r - \beta\sqrt{c_0 t}),$$

Lemma 6.3 (The expanding balls lemma, Lemma 3.1 in [16]). *Suppose $(M, g(t))$ is a Ricci flow for $t \in [-T, 0]$, $T > 0$, on a manifold M of any dimension. Suppose that $x_0 \in M$ and that $B_{g(0)}(x_0, R) \subset\subset M$ and $\text{Ric}_{g(t)} \geq -K < 0$ on $B_{g(0)}(x_0, R) \cap B_{g(t)}(x_0, Re^{Kt})$ for each $t \in [-T, 0]$. Then for all $t \in [-T, 0]$, we have*

$$(6.6) \quad B_{g(0)}(x_0, R) \supset B_{g(t)}(x_0, Re^{Kt})$$

The following theorem provides a local maximum principles along Ricci flow.

Theorem 6.4 (Theorem 1.1 in [14]). *Let $(M^n, g(t))$, $t \in [0, T]$, be a smooth solution to the Ricci flow which is possibly incomplete. Suppose*

$$(6.7) \quad \text{Ric}_{g(t)} \leq \frac{\alpha}{t}$$

on $M \times (0, T]$ for some $\alpha > 0$. Let $\varphi(x, t)$ be a continuous function on $M \times [0, T]$ which satisfies $\varphi(x, t) \leq \frac{\alpha}{t}$ on $M \times (0, T]$ and

$$(6.8) \quad \left(\frac{\partial}{\partial t} - \Delta_{g(t)} \right) \varphi|_{(x_0, t_0)} \leq L(x_0, t_0) \varphi(x_0, t_0)$$

whenever $\varphi(x_0, t_0) > 0$ in the sense of barrier, for some continuous function $L(x, t)$ on $M \times [0, T]$ with $L(x, t) \leq \frac{\alpha}{t}$. Suppose $p \in M$ such that $B_{g(0)}(p, 2) \subset\subset M$ and $\varphi(x, 0) \leq 0$ on $B_{g(0)}(p, 2)$. Then for any $l > \alpha + 1$, there exists $\hat{T} = \hat{T}(n, \alpha, l) > 0$ such that for $t \in [0, \min\{T, \hat{T}\}]$,

$$(6.9) \quad \varphi(p, t) \leq t^l.$$

Corollary 6.5. *Let $(M^n, g(t))$, $t \in [0, T]$ be a smooth solution to the Ricci flow which is possibly incomplete. Suppose*

$$(6.10) \quad \text{Ric}_{g(t)} \leq \frac{\alpha}{t}$$

on $M \times (0, T]$ for some $\alpha > 0$. Suppose $p \in M$ such that $B_{g(0)}(p, 2) \subset\subset M$ and $\text{Sc}(p) \geq 2$ on $B_{g(0)}(p, 2)$. Then there exists $\hat{T} = \hat{T}(n, \alpha) > 0$ such that for $t \in [0, \min\{T, \hat{T}\}]$,

$$(6.11) \quad \text{Sc}_{g(t)}(p) \geq 2 - t^{n\alpha+3}.$$

Proof. Take $\varphi(x, t) = 2 - \text{Sc}_{g(t)}$, and $L(x, t) = \frac{4+2\text{Sc}_{g(t)}}{n}$, then $\varphi(x, t) \leq \frac{n\alpha}{t}$, $L(x, t) \leq \frac{n\alpha+1}{t}$, and

$$(6.12) \quad \left(\frac{\partial}{\partial t} - \Delta_{g(t)}\right)\varphi = -2|\text{Ric}_{g(t)}|^2 \leq -2\frac{\text{Sc}_{g(t)}^2}{n} < L\varphi.$$

According to Theorem 6.4, (6.11) holds for suitable $\hat{T} = \hat{T}(n, \alpha)$. $\square$

The following theorem, which will be used in our proof of Theorem 1.6, slightly generalising Theorem 1.8 in [17]. Such kind of result may be known to experts, and it is proof is a slight modification of that of [17]. We include the proof for completeness.

Theorem 6.6. *Let (M_i, g_i, p_i) be a sequence of 3-dimensional Riemannian manifolds (possibly incomplete). Suppose $U_i = B_{g_i}(p_i, s_i) \subset\subset M_i$ for a sequence of $s_i \rightarrow \infty$, and $\text{Ric}_{g_i} \geq -\alpha_0$ (where $\alpha_0 \in [0, 1]$) on $B_{g_i}(p_i, s_i)$, and $\text{Vol}_{g_i} B(p, 1) > v_0 > 0$ for all $p \in B_{g_i}(p_i, s_i - 1)$ and all $i \in \mathbb{N}$. Then there exists a smooth manifold M , a point $p_\infty \in M$, a complete Ricci flow $g(t)$ on M for $t \in (0, T]$, where $T > 0$ depends only on α_0 and v_0 , and a continuous distance metric d_0 on M such that $d_{g(t)} \rightarrow d_0$ locally uniformly as $t \downarrow 0$, and up to passing to a subsequence we have that (U_i, g_i, p_i) converges in the pointed Gromov-Hausdorff sense to (M, d_0, p_∞) . Furthermore, the Ricci flow satisfies*

$$(6.13) \quad \text{Ric}_{g(t)} \geq -C\alpha_0 \quad \text{on } M \times (0, T],$$

$$(6.14) \quad |\text{Rm}_{g(t)}| \leq \frac{C}{t} \quad \text{on } M \times (0, T],$$

$$(6.15) \quad \text{inj}_{g(t)} \geq \rho_0 \sqrt{t} \quad \text{for all } t \in (0, T],$$

$$(6.16) \quad d_{g(t_1)}(x, y) - \beta \sqrt{C}(\sqrt{t_2} - \sqrt{t_1}) \leq d_{g(t_2)}(x, y) \leq e^{C\alpha_0(t_2-t_1)} d_{g(t_1)}(x, y).$$

for any $0 \leq t_1 \leq t_2 \leq T$ and $x, y \in M$, where C depends only on v_0 , and $\beta > 0$ is a universal constant.

Proof. By Theorem 6.1, there exists a Ricci flow $(g_i(t))_{t \in [0, T]}$ on $B_{g_i}(p_i, s_i - 2)$, with $g_i(0) = g$, so that

$$(6.17) \quad \text{Ric}_{g_i(t)} \geq -C\alpha_0 \quad \text{on } B_{g_i}(p_i, s_i - 2),$$

$$(6.18) \quad |\text{Rm}_{g_i(t)}| \leq \frac{C}{t} \quad \text{on } B_{g_i}(p_i, s_i - 2),$$

for some $T = T(\alpha_0, v_0)$, $C = C(v_0)$.

By the shrinking balls lemma 6.2 and the expanding balls lemma 6.3, up to replacing T by a smaller one, we have

$$(6.19) \quad B_{g_i(0)}(p_i, s_i - 2) \supset B_{g_i(t)}(p_i, s_i - 4)$$

and

$$(6.20) \quad B_{g_i(t)}(p_i, \frac{1}{2}s_i - 2) \supset B_{g_i(s)}(p_i, \frac{1}{4}s_i - 1)$$

for any $0 \leq s \leq t \leq T$.

Denote by $U'_i = B_{g_i(0)}(p_i, \frac{1}{4}s_i - 2)$ for short. For any $0 \leq t \leq T$ and $x, y \in U'_i$, by (6.20), the shortest curve with respect to $g_i(t)$ connecting x and y is contained in $B_{g_i(t)}(p_i, s_i - 4) \subset B_{g_i(0)}(p_i, s_i - 2)$. Making use of (6.17) and (6.18), it is well-known that, there exists $\beta > 0$ such that for any $0 \leq t_1 \leq t_2 \leq T$ and $x, y \in U'_i$ (see e.g. [16]), we have

$$(6.21) \quad d_{g_i(t_1)}(x, y) - \beta\sqrt{C}(\sqrt{t_2} - \sqrt{t_1}) \leq d_{g_i(t_2)}(x, y) \leq e^{C\alpha_0(t_2-t_1)}d_{g_i(t_1)}(x, y).$$

By (6.18), according to Shi's derivative estimates, for any integer $k \geq 1$,

$$(6.22) \quad |\nabla^k \text{Rm}_{g_i(t)}| \leq \frac{C_k}{t^{1+\frac{k}{2}}} \quad \text{on } U'_i,$$

where C_k depends on C and k . By Lemma 4.1 in [16], there exists a positive constant ρ_0 depending only on v_0 so that

$$(6.23) \quad \text{inj}_{g_i(t)} \geq \rho_0\sqrt{t} \quad \text{on } U'_i.$$

By Hamilton's compactness theorem, up to choosing a subsequence, $(U'_i, g_i(t), p_i)_{t \in (0, T]}$ converges in the smooth Cheeger-Gromov sense to a Ricci flow solution $(M, g(t), p_\infty)_{t \in (0, T]}$, where M is a three-manifold. By the shrinking lemma, it is easy to see that for any $t \in (0, T]$, $(M, g(t))$ is complete. By the smooth convergence for $t \in (0, T]$, it is obvious that (6.13)-(6.15) hold. Note that (6.16) holds for any $0 < t_1 \leq t_2 \leq T$ and $x, y \in M$, which ensures the existence of a metric d_0 on M so that $d_{g(t)}$ converges to d_0 locally uniformly as $t \rightarrow 0$, and for any $x, y \in M$ and $0 < t \leq T$, we have

$$(6.24) \quad d_0(x, y) - \beta\sqrt{C}t \leq d_{g(t)}(x, y) \leq e^{C\alpha_0 t}d_0(x, y).$$

By (6.24), (6.21) and the Cheeger-Gromov convergence of $(U'_i, g_i(t))$ to $(M, g(t))$ for every small $t > 0$ (6.24), by the triangle inequality it is easy to check that $(U'_i, g_i(0))$ converges to (M, d_0) in Gromov-Hausdorff sense. $\square$

Corollary 6.7. *If in Theorem 6.6, we assume in addition that $\text{Sc}_{g_i} \geq 2$ on $U_i = B_{g_i}(p_i, s_i)$, then on $(M, g(t))_{t \in (0, T]}$, it holds*

$$(6.25) \quad \text{Sc}_{g(t)} \geq 2.$$

Proof. Since (6.18) holds, according to Corollary 6.5, for any $t \in (0, T]$ (here we can choose T to a smaller one), we have

$$(6.26) \quad \text{Sc}_{g_i(t)} \geq 2 - t^\tau \quad \text{on } B_{g_i}(p_i, s_i - 4)$$

for some $\tau > 0$ depending only on v_0 . Then by the pointed Cheeger-Gromov convergence of $(U'_i, g_i(t))$ to $(M, g(t))$, we know

$$(6.27) \quad \text{Sc}_{g(t)} \geq 2 - t^\tau \quad \text{on } M \times (0, T].$$

For any $t_0 \in (0, T]$, since the Ricci flow $(M, g(t))_{t \in [t_0, T]}$ is complete and has uniformly bounded curvature, by the standard maximum principle, the lower bound of scalar curvature is preserved, i.e.

$$(6.28) \quad \text{Sc}_{g(t)} \geq 2 - t_0^\tau \quad \text{on } M \times [t_0, T].$$

By the arbitrariness of t_0 , we conclude that (6.25) holds. $\square$

Proof of (2) of Theorem 1.6: By an argument by contradiction, we will prove that on every end E with a ray $\gamma : [0, \infty) \rightarrow E$, it holds:

$$(6.29) \quad \limsup_{r \rightarrow \infty} \frac{\text{Vol}(E \cap B(p, r))}{r} \leq 4\pi.$$

Let b_γ be the Busemann function associated with γ . We will prove that, for any $\epsilon > 0$, there exists some sufficiently large $R_0 > 0$ so that for any $r > R_0$, it holds

$$(6.30) \quad \text{Vol}(\{-1 \leq b_\gamma - r \leq 1\}) < 8\pi + \epsilon.$$

If (6.30) is proved, then for every $r > r_0$,

$$(6.31) \quad \begin{aligned} \text{Vol}(E \cap B(p, r)) &\leq \text{Vol}(E \cap b^{-1}([0, r])) \\ &\leq \text{Vol}(E \cap b^{-1}([0, r_0])) + \text{Vol}(E \cap b^{-1}([r_0, r])) \\ &\leq \text{Vol}(E \cap b^{-1}([0, r_0])) + (8\pi + \epsilon)\left(\frac{r - r_0}{2} + 1\right), \end{aligned}$$

and hence $\limsup_{r \rightarrow \infty} \frac{\text{Vol}(E \cap B(p, r))}{r} \leq 4\pi + \frac{\epsilon}{2}$. Then (6.29) holds by the arbitrariness of ϵ .

We prove (6.30) by contradiction. Suppose on the contrary, there is some $\epsilon_0 > 0$ such that there is a sequence of $r_i \uparrow +\infty$ and

$$(6.32) \quad \text{Vol}(\{-1 \leq b_\gamma - r_i \leq 1\}) \geq 8\pi + \epsilon_0.$$

Denote by $p_i = \gamma(r_i)$, $s_i = \frac{r_i}{2}$, $\epsilon_i = f(s_i)$, $V_i = B_g(p_i, s_i)$.

Note that on V_i the Ricci curvature has a lower bound $-\epsilon_i$. In addition, by Proposition 1.10, there exists some $v_0 > 0$ such that $\text{Vol}(B(p, 1)) > v_0 > 0$ for all $p \in V_i$. By Theorem 6.6 and Corollary 6.7, there exists a smooth manifold X , a point $p_\infty \in X$, a complete Ricci flow $g(t)$ on X for $t \in (0, T]$, and a continuous distance metric d_0 on X such that $d_{g(t)} \rightarrow d_0$ locally uniformly as $t \downarrow 0$, and up to passing to a subsequence we have that (V_i, g, p_i) converges in the pointed Gromov-Hausdorff sense to (X, d_0, p_∞) , and (6.25) holds on $X \times (0, T]$. Furthermore, since $\text{Ric}_g \geq -\epsilon_i$ ($\epsilon_i \rightarrow 0$) on V_i , from the smooth convergence for $t \in (0, T]$ in the proof of Theorem 6.6, it is easy to see that, for each $t \in (0, T]$, we have

$$(6.33) \quad \text{Ric}_{g(t)} \geq 0$$

on X , and for any $x, y \in X$ and $0 < t \leq T$, we have

$$(6.34) \quad d_0(x, y) - \beta\sqrt{Ct} \leq d_{g(t)}(x, y) \leq d_0(x, y).$$

On the other hand, by Cheeger-Colding's theory, up to choosing a subsequence, (V_i, g, p_i) converges in the measure Gromov-Hausdorff sense to $(X, d_0, \mathcal{H}^3, p_\infty)$. Note that $(X, d_\infty, \mathcal{H}^3)$ is a non-collapsed RCD(0, 3) space (we say ncRCD(0, 3) for short) in the sense of [7]. On the other hand, by the choice of V_i , it is obvious that X contains a line, hence by the splitting theorem of Gigli (see [11]), there exists a ncRCD(0, 2) space $(Y, d_Y, \mathcal{H}^2)$ such that $(X, d_0, \mathcal{H}^3)$ is isomorphic to $(\mathbb{R} \times Y, d_{\text{Eucl}} \otimes d_Y, \mathcal{L}^1 \otimes \mathcal{H}^2)$.

Denote by $f_i = b_\gamma - r_i$, then each f_i is 1-Lipschitz with $f_i(p_i) = 0$. According to Proposition 27.20 of [20], up to a subsequence, f_i converges locally uniformly to a 1-Lipschitz function P on (X, d_0) . By Corollary 4.3, there exist positive constants C_1 and R_0 such that

$$(6.35) \quad \text{diam}(b^{-1}(R)) \leq C_1 \quad \text{for every } R \geq R_0.$$

Combining this fact with Lemma 2.1, it is not hard to prove that

$$(6.36) \quad \text{diam}(P^{-1}(R)) \leq C_1 \quad \text{for every } R.$$

On the other hand, for any $\mu_0 \in \mathbb{R}$, denote by $F_i := f_i^{-1}([\mu_0, \infty))$. According to Lemma 2.1, $\mu_0 - f_i(x) = d(x, F_i)$, for every $x \in V_i \setminus F_i$. Since $\mu_0 - f_i$ converges to $\mu_0 - P$ uniformly, similar to Lemma 3.25 in [12], one can prove that $\mu_0 - P(x) = d(x, P^{-1}([\mu_0, \infty)))$ for every

$x \in X$ satisfying $P(x) < \mu_0$. By the arbitrariness of μ_0 , it is easy to see that there exists a line $\eta \subset X$ such that

$$(6.37) \quad P(\eta(t_1)) - P(\eta(t_2)) = t_1 - t_2 \text{ for every } t_1, t_2 \in \mathbb{R}.$$

Note that the projection of η to the Y factor must be a point. For otherwise the projection on Y is line, and by the splitting theorem of Gigli, (X, d_0) is isometric to $(\mathbb{R}^2 \times Z, d_{\text{Eucl}} \otimes d_Z)$, contradicting to (6.36) and (6.37). Thus the map P is just the projection of $\mathbb{R} \times Y$ onto the $\mathbb{R}$ -factor, and hence

$$(6.38) \quad \text{diam}(Y) \leq C_1.$$

Claim: for every $t \in (0, T]$, $(X, g(t))$ contains a line.

Proof of the claim: We will prove the claim holds for every sufficiently small $t > 0$, then the Ricci flow solution ensures the validity of the claim for larger t . Let $\eta : (-\infty, +\infty)$ be the line on $\mathbb{R} \times Y$ with $\eta(0) = p_\infty$. Now we fix a small t . For every large $R > 0$, denote by $x_\infty = \eta(-R)$, $y_\infty = \eta(R)$. By (6.34), $|d_{g(t)}(x_\infty, y_\infty) - 2R| \leq C_2\sqrt{t}$, $|d_{g(t)}(p_\infty, x_\infty) - R| \leq C_2\sqrt{t}$ and $|d_{g(t)}(p_\infty, y_\infty) - R| \leq C_2\sqrt{t}$. Let $z^{(R)}$ be the mid-point of a geodesic, $\eta^{(R)} : [-\frac{1}{2}d_{g(t)}(x_\infty, y_\infty), \frac{1}{2}d_{g(t)}(x_\infty, y_\infty)] \rightarrow X$, with respect to $g(t)$, which connecting x_∞ and y_∞ , then by (6.34) again, $|d_0(x_\infty, z^{(R)}) - R| \leq \frac{3}{2}C_2\sqrt{t}$ and $|d_0(y_\infty, z^{(R)}) - R| \leq \frac{3}{2}C_2\sqrt{t}$. Combining them with the product structure $\mathbb{R} \times Y$ and (6.38), it is easy to see that there exists a $C_3 > 0$ depending only on C_1 and $C_2\sqrt{t}$ (we assume R is sufficiently large, and it is easy to see that C_3 is independent of R) so that $d_0(p_\infty, z^{(R)}) \leq C_3$. Now we take a sequence of $R_i \rightarrow +\infty$, and take geodesics $\eta^{(R_i)}X$ with respect to $g(t)$, whose length $\approx 2R_i$, and with midpoint $z^{(R_i)}$ contained in a compact set. After taking a converging subsequence, we obtain a line in $(X, g(t))$. $\square$

By the above claim and (6.33), $(M_\infty, g_\infty(t))_{t \in (0, T]}$ is isometric to $(\mathbb{R}, g_{\text{Eucl}}) \otimes (\tilde{S}, \tilde{g}(t))_{t \in (0, T]}$, where $\tilde{g}(t)$, $t \in (0, T]$, is a Ricci flow solution on a smooth manifold $\tilde{S}$. By (6.25), it is easy to see that

$$(6.39) \quad \sec_{\tilde{g}(t)} \geq 1.$$

Hence $\tilde{S}$ is either diffeomorphic to S^2 or RP^2 , and each $(\tilde{S}, \tilde{g}(t))$ is a ncRCD(1, 2) space. Clearly, the pointed Gromov-Hausdorff convergence from $(X, g(t), p_\infty)$ (where $t \downarrow 0$) to (X, d_0, p) induces the Gromov-Hausdorff convergence from $(S^2, \tilde{g}(t))$ to (Y, d_Y) . So the limit $(Y, d_Y, \mathcal{H}^2)$ satisfies ncRCD(1, 2) condition. By the generalized relative volume comparison theorem for CD-spaces (see [19]), it is easy to see that

$$(6.40) \quad \mathcal{H}^2(Y) \leq 4\pi.$$

Since $f_i = b_\gamma - r_i$ locally uniformly converges to the projection $P : \mathbb{R} \times Y \rightarrow \mathbb{R}$, and (6.35) (6.38) hold, we have that (see e.g. Proposition 2.10 in [29]), for every large i ,

$$(6.41) \quad \begin{aligned} & |\text{Vol}(\{-1 \leq b_\gamma - r_i \leq 1\}) - 2\mathcal{H}^2(Y)| \\ &= |\text{Vol}(\{-1 \leq b_\gamma - r_i \leq 1\}) - \mathcal{H}^3(\{-1 \leq P \leq 1\})| \\ &< \frac{\epsilon_0}{2}. \end{aligned}$$

Combining it with (6.40), we have

$$(6.42) \quad \text{Vol}(\{-1 \leq b_\gamma - r_i \leq 1\}) < 8\pi + \frac{\epsilon_0}{2},$$

contradicting to (6.32). This finishes the proof of Theorem 1.6. $\square$

Remark 6.8. The argument to derive (6.40) is motivated by [28], where the authors is concentrated in the optimal diameter upper bound of Y . Some technical differences, such as the possible negative Ricci curvature and the non-global positive scalar curvature lower bound, require some adjustment in our proof. In addition, our proof of the Claim is different from [28]. In fact, the argument in [28] to ensure the splitting of $(X, g(t))$ can be carried to our setting, but we provide such a different proof because we think that it may be used in other related problems.

REFERENCES

- [1] R. Bamler, E. Cabezas-Rivas, B. Wilking, The Ricci flow under almost non-negative curvature conditions, *Invent. Math.*, 217 (2019)
- [2] E. Calabi, On manifolds with non-negative Ricci-curvature II. *Notices Amer. Math. Soc.* 22 (1975), A205.
- [3] J. Cheeger, Critical points of distance functions and applications to geometry, in *Geometric Topology and Recent Developments*, Montecatini Terme, 1990:1-38; *Lecture Notes in Mathematics*, Springer, Berlin, 1991, 1504.
- [4] J. Cheeger, T. H. Colding, Lower bounds on Ricci curvature and the almost rigidity of warped products. *Ann. of Math.*, 144 (1996), 189-237.
- [5] O. Chodosh, C. Li, D. Stryker, Complete stable minimal hypersurfaces in positively curved 4-manifolds, to appear in *Journal of European Mathematical Society*, arXiv:2202.07708.
- [6] O. Chodosh, C. Li, D. Stryker, Volume growth of 3-manifolds with scalar curvature lower bounds, *Proc. Amer. Math. Soc.* Vol(151) (2023), 4501-4511.
- [7] G. De Philippis, N. Gigli, From volume cone to metric cone in the nonsmooth setting. *Geom. Funct. Anal.*, 26 (2016), 1526-1587.
- [8] M. Gromov, Large Riemannian manifolds, Curvature and topology of Riemannian manifolds, *Lecture Notes in Mathematics*, Vol.1201, Springer, Berlin, 1986:108-121.
- [9] M. Gromov, Positive Curvature, Macroscopic Dimension, Spectral Gaps and Higher Signatures, *Functional Analysis on the Eve of the 21st Century* (Birkhäuser, Boston, MA, 1996), Vol. 2, pp. 1-213.
- [10] M. Gromov, Metric inequalities with scalar curvature, *Geom. Funct. Anal.* 28 (2018), no. 3, 645-726.
- [11] N. Gigli, The splitting theorem in non-smooth context. arXiv:1302.5555, (2013).
- [12] X. T. Huang, An almost rigidity theorem and its applications to noncompact $\mathrm{RCD}(0, N)$ spaces with linear volume growth, *Communications in Contemporary Mathematics*, Vol. 22, No. 4 (2020).
- [13] G. Liu, 3-manifolds with nonnegative Ricci curvature, *Inventiones Mathematicae*, 2013, 193 (2013), 367-375.
- [14] M.-C. Lee, L.-F. Tam, Some local maximum principles along Ricci flows. *Canad. J. Math.*, 74 (2022), 329-348.
- [15] O. Munteanu, J. Wang, Geometry of three-dimensional manifolds with scalar curvature lower bound, (2022), arxiv.2201.05595v2.
- [16] M. Simon, P. M. Topping, Local control on the geometry in 3D Ricci flow, *J. Differential Geom.* 122 (2022), no. 3, 467-518
- [17] M. Simon, P. M. Topping, Local mollification of Riemannian metrics using Ricci flow, and Ricci limit spaces, *Geom. Topol.* 25 (2021), no. 2, 913-948.
- [18] C. Sormani, Busemann functions on manifolds with lower bounds on Ricci curvature and minimal volume growth, *Journal Differential Geometry*, 48 (1998), 557-585
- [19] K. T. Sturm, On the geometry of metric measure spaces II, *Acta Math.* 196 (2006), 133-177.
- [20] C. Villani, Optimal transport, Old and new. *Grundlehren der Mathematischen Wissenschaften*, Springer-Verlag, Berlin, vol. 338 (2009).
- [21] Y. Wang, Effective volume growth of three-manifolds with positive scalar curvature, arXiv:2405.03023 (2024).
- [22] J. Wang, Z. Xie, B. Zhu, X. Zhu, Positive Scalar Curvature Meets Ricci Limit Spaces, arXiv:2212.10416 (2022).
- [23] G. Wei, G. Xu, S. Zhang, Volume growth and positive scalar curvature, arXiv:2405.04001 (2024)
- [24] G. Xu, Integral of scalar curvature on manifolds with a pole, to appear in *Proceedings of the American Mathematical Society*, arXiv:2306.07460v1. (2023)
- [25] S. Yau, Some function-theoretic properties of complete Riemannian manifolds and their applications to geometry. *Indiana Math. J.*, 25 (1976), 659-670.

- [26] S. Yau, Open problems in geometry, Chern—a great geometer of the twentieth century, Int. Press, Hong Kong, 1992:275-319.
- [27] B. Zhu, Geometry of positive scalar curvature on complete manifold, Journal für die Reine und Angewandte Mathematik, 791 (2022), 225-246.
- [28] B. Zhu, X. Zhu, Optimal diameter estimates of three-dimensional Ricci limit spaces, Proc. Amer. Math. Soc., 152 (2024), 815-821.
- [29] X. Zhu, On the geometry at infinity of manifolds with linear volume growth and nonnegative Ricci curvature. 2023, arXiv:2310.00640.

SCHOOL OF MATHEMATICS SUN YAT-SEN UNIVERSITY GUANGZHOU 510275
Email address: `hxiant@mail2.sysu.edu.cn`; `huangxt28@mail.sysu.edu.cn`

SCHOOL OF MATHEMATICS SUN YAT-SEN UNIVERSITY GUANGZHOU 510275
Email address: `liush328@mail2.sysu.edu.cn`