

A motivic integral identity for (-1) -shifted symplectic stacks

Chenjing Bu

Abstract

We prove a motivic integral identity relating the motivic Behrend function of a (-1) -shifted symplectic stack to that of its stack of graded points. This generalizes analogous identities for moduli stacks of objects in a 3-Calabi–Yau category obtained by Kontsevich–Soibelman [40] and Joyce–Song [35], which are crucial in proving wall-crossing formulae for Donaldson–Thomas invariants. We expect our identity to be useful in generalizing motivic Donaldson–Thomas theory to general (-1) -shifted symplectic stacks.

Contents

1	Introduction	2
2	Motivic vanishing cycles	6
2.1	Rings of motives	6
2.2	Descent of motives	11
2.3	Motivic vanishing cycles for schemes	12
2.4	Motivic vanishing cycles for stacks	15
2.5	The motivic Behrend function	20
3	Graded and filtered points	24
3.1	Definition and deformation theory	24
3.2	Local structure	28
4	The main results	30
4.1	The local version	30
4.2	The global version	35
4.3	The numerical version	39
	References	42

1 Introduction

1.1.1. Let $\mathbb{K}$ be an algebraically closed field of characteristic zero, and let $\mathfrak{X}$ be a (-1) -shifted symplectic derived algebraic stack over $\mathbb{K}$, in the sense of Pantev–Toën–Vaquié–Vezzosi [47].

One of the motivating examples of such a stack is the moduli stack $\mathfrak{X} = \mathfrak{M}_{\mathcal{A}}$ of objects in a $\mathbb{K}$ -linear 3-Calabi–Yau category $\mathcal{A}$, such as the category of coherent sheaves on a smooth projective Calabi–Yau threefold. See Brav–Dyckerhoff [12] for more details on this.

Starting from $\mathfrak{X}$, one can consider the derived mapping stacks

$$\mathrm{Grad}(\mathfrak{X}) = \mathfrak{Map}([*/\mathbb{G}_m], \mathfrak{X}), \quad (1.1)$$

$$\mathrm{Filt}(\mathfrak{X}) = \mathfrak{Map}([\mathbb{A}^1/\mathbb{G}_m], \mathfrak{X}), \quad (1.2)$$

called the *stack of graded points* and the *stack of filtered points* of $\mathfrak{X}$, respectively, as in Halpern-Leistner [26]. For example, if $\mathfrak{X} = \mathfrak{M}_{\mathcal{A}}$, then $\mathrm{Grad}(\mathfrak{X})$ is the moduli stack of $\mathbb{Z}$ -graded objects in $\mathcal{A}$, and $\mathrm{Filt}(\mathfrak{X})$ is the moduli stack of $\mathbb{Z}$ -filtered objects in $\mathcal{A}$. In particular, there are inclusions

$$\mathfrak{M}_{\mathcal{A}} \times \mathfrak{M}_{\mathcal{A}} \hookrightarrow \mathrm{Grad}(\mathfrak{M}_{\mathcal{A}}), \quad (1.3)$$

$$\mathfrak{M}_{\mathrm{Exact}(\mathcal{A})} \hookrightarrow \mathrm{Filt}(\mathfrak{M}_{\mathcal{A}}) \quad (1.4)$$

as closed and open substacks, i.e. disjoint unions of connected components, where $\mathfrak{M}_{\mathrm{Exact}(\mathcal{A})}$ is the moduli stack of short exact sequences in $\mathcal{A}$. These substacks can be given by, for example, graded and filtered objects that only have non-trivial factors in the degrees 0 and 1.

1.1.2. We show in Theorem 3.1.5 that there is a (-1) -shifted Lagrangian correspondence

$$\mathrm{Grad}(\mathfrak{X}) \xleftarrow{\mathrm{gr}} \mathrm{Filt}(\mathfrak{X}) \xrightarrow{\mathrm{ev}_1} \mathfrak{X}. \quad (1.5)$$

When $\mathfrak{X} = \mathfrak{M}_{\mathcal{A}}$, the morphisms gr and ev_1 send a filtered object to its associated graded object and its total object, respectively. In particular, restricting it to the substacks (1.3)–(1.4), this gives the (-1) -shifted Lagrangian correspondence

$$\mathfrak{M}_{\mathcal{A}} \times \mathfrak{M}_{\mathcal{A}} \xleftarrow{(p_1, p_3)} \mathfrak{M}_{\mathrm{Exact}(\mathcal{A})} \xrightarrow{p_2} \mathfrak{M}_{\mathcal{A}} \quad (1.6)$$

as in Brav–Dyckerhoff [12], where $p_1, p_2, p_3 : \mathfrak{M}_{\mathrm{Exact}(\mathcal{A})} \rightarrow \mathfrak{M}_{\mathcal{A}}$ send a short exact sequence to its three respective terms.

The correspondence (1.6) has proved to be useful in enumerative geometry. It lies in the heart of the construction of Hall-algebra-type algebraic structures, including *motivic Hall algebras* studied by Joyce [29], *cohomological Hall algebras* introduced by Kontsevich–Soibelman [41], and *Joyce vertex algebras* constructed by Joyce [32; 33]. These structures are closely related to the construction of *Donaldson–Thomas invariants* and other enumerative invariants.

1.1.3. Following a series of work [7; 10; 11; 14; 31] by Joyce and his collaborators, it is known that a (-1) -shifted symplectic stack $\mathfrak{X}$ can be locally modelled as *derived critical loci* of functions on smooth stacks. When $\mathfrak{X}$ is equipped with *orientation data*, one can define an element $v_{\mathfrak{X}}^{\text{mot}}$ in the ring of monodromic motives on $\mathfrak{X}$, which we call the *motivic Behrend function* of $\mathfrak{X}$. It is locally modelled by the *motivic vanishing cycle* defined by Denef–Loeser [21], and is a motivic enhancement of the *Behrend function* $v_{\mathfrak{X}}: \mathfrak{X} \rightarrow \mathbb{Z}$ introduced by Behrend [6] and extended by Joyce–Song [35, §4.1] to algebraic stacks.

When $\mathfrak{X} = \mathfrak{M}_{\mathcal{A}}$ as above, the element $v_{\mathfrak{M}_{\mathcal{A}}}^{\text{mot}}$ was considered by Kontsevich–Soibelman [40], and is important in the Donaldson–Thomas theory of $\mathcal{A}$. Given a stability condition on $\mathcal{A}$, if $\mathfrak{M}_{\alpha} \subset \mathfrak{M}_{\mathcal{A}}$ is a component, then the *motivic Donaldson–Thomas invariant* of the class α is the monodromic motive given by the *motivic integral*

$$\text{DT}_{\alpha}^{\text{mot}} = \int_{\mathfrak{M}_{\alpha}} (\mathbb{L}^{1/2} - \mathbb{L}^{-1/2}) \cdot \epsilon_{\alpha} \cdot v_{\mathfrak{M}_{\mathcal{A}}}^{\text{mot}}, \quad (1.7)$$

where ϵ_{α} is a *weight function* encoding the data of the stability condition. The numerical *Donaldson–Thomas invariant* DT_{α} is then equal to the Euler characteristic of this monodromic motive, so that

$$\text{DT}_{\alpha} = \int_{\mathfrak{M}_{\alpha}} (\mathbb{L}^{1/2} - \mathbb{L}^{-1/2}) \cdot \epsilon_{\alpha} \cdot v_{\mathfrak{M}_{\mathcal{A}}} d\chi. \quad (1.8)$$

See also Joyce–Song [35] for more on this numerical version.

1.1.4. The main result of this paper, [Theorem 4.2.2](#), states that the motivic Behrend functions of $\mathfrak{X}$ and $\text{Grad}(\mathfrak{X})$ are related via the correspondence (1.5), by the identity

$$\text{gr}_! \circ \text{ev}_1^*(v_{\mathfrak{X}}^{\text{mot}}) = \mathbb{L}^{\text{vdim Filt}(\mathfrak{X})/2} \cdot v_{\text{Grad}(\mathfrak{X})}^{\text{mot}}, \quad (1.9)$$

as an identity of monodromic motives on $\text{Grad}(\mathfrak{X})$, where $\text{vdim Filt}(\mathfrak{X})$ is the virtual dimension of $\text{Filt}(\mathfrak{X})$. In particular, evaluating this at a graded point $\gamma \in \text{Grad}(\mathfrak{X})$ gives the motivic integral identity

$$\int_{\varphi \in \text{gr}^{-1}(\gamma)} v_{\mathfrak{X}}^{\text{mot}}(\text{ev}_1(\varphi)) = \mathbb{L}^{\text{vdim Filt}(\mathfrak{X})/2} \cdot v_{\text{Grad}(\mathfrak{X})}^{\text{mot}}(\gamma), \quad (1.10)$$

as an identity of monodromic motives over $\mathbb{K}$.

We prove this identity by first proving a local version of it in [Theorem 4.1.1](#), which is, roughly speaking, the special case of (1.9) when $\mathfrak{X} = [\text{Crit}(f)/\mathbb{G}_m]$ is a derived critical locus, where f is a $\mathbb{G}_m$ -invariant function on a $\mathbb{G}_m$ -equivariant smooth $\mathbb{K}$ -variety. Our proof of this local version involves the theory of *nearby and vanishing cycles* for rings of motives on algebraic stacks, which we develop in §2.4, with [Theorem 2.4.4](#) as a main result. Then, we prove the global version of the identity by gluing together the local models.

1.1.5. We explain the relation between the identity (1.9) and known results and conjectures in the literature.

Firstly, these identities can be seen as a global version and a generalization of an integral identity conjectured by Kontsevich–Soibelman [40, Conjecture 4], later proved by Lê [44]. The local version of our identity, Theorem 4.1.1, is stated in a form similar to Kontsevich–Soibelman’s identity, and generalizes it by removing the assumption that the torus action only has weights -1 , 0 , and 1 . It is crucial that this assumption is removed in order for the identity to serve as a local model for (1.9) for general (-1) -shifted symplectic stacks, not only for stacks of the form $\mathfrak{M}_{\mathcal{A}}$.

Kontsevich–Soibelman then used their identity to prove [40, Theorem 8], which can be seen as a special case of our identity (1.9) when $\mathfrak{X} = \mathfrak{M}_{\mathcal{A}}$ as above. Their theorem is a key ingredient in proving *wall-crossing formulae* of motivic Donaldson–Thomas invariants, governing the behaviour of these invariants under changes of stability conditions.

Secondly, by taking the Euler characteristic of our identity, we obtain numerical integral identities in Theorem 4.3.3. These identities are direct generalizations of the Behrend function identities of Joyce–Song [35, Theorem 5.11] to general (-1) -shifted symplectic stacks.

Thirdly, the identity (1.9) is related to a conjecture on perverse sheaves, sometimes known as the *Joyce conjecture*, formulated in Joyce–Safronov [34, Conjecture 1.1], also discussed in Amorim–Ben-Bassat [5, §5.3]. One form of the conjecture states that for an oriented (-1) -shifted Lagrangian correspondence

$$\mathfrak{X} \xleftarrow{f} \mathfrak{L} \xrightarrow{g} \mathfrak{Y}, \quad (1.11)$$

under certain assumptions, there should exist a natural morphism

$$\mu_{\mathfrak{L}}: f_! \circ g^*(\mathcal{P}_{\mathfrak{Y}}) \longrightarrow \mathcal{P}_{\mathfrak{X}}[-\mathrm{vdim} \mathfrak{L}], \quad (1.12)$$

satisfying certain properties, where $\mathcal{P}_{\mathfrak{X}}$ and $\mathcal{P}_{\mathfrak{Y}}$ are the perverse sheaves constructed in Ben-Bassat–Brav–Bussi–Joyce [7, Theorem 4.8], sometimes called the *Donaldson–Thomas perverse sheaves*. They can be seen as analogues of the motivic Behrend functions $v_{\mathfrak{X}}^{\mathrm{mot}}$ and $v_{\mathfrak{Y}}^{\mathrm{mot}}$ in cohomological Donaldson–Thomas theory.

In the special case when the correspondence (1.11) is taken to be the correspondence (1.5), a recent result of Kinjo–Park–Safronov [37] shows that there is a natural isomorphism of the form (1.12), strengthening the Joyce conjecture in this special case. In this sense, the identity (1.9) can be seen as a motivic analogue of this version of the Joyce conjecture.

1.1.6. In future work, the author and his collaborators plan to extend the definition of the weight functions ϵ_{α} mentioned in §1.1.3 to a more general class of algebraic stacks, which will allow us to define motivic Donaldson–Thomas invariants for these general stacks. The integral identity (1.9) will hopefully be helpful in the study of these invariants, such as formulating wall-crossing formulae for these generalized invariants.

For example, the author [13] defines these weight functions for the moduli stack of *self-dual objects* in a *self-dual $\mathbb{K}$ -linear category*, such as a certain compactification of the stack of principal orthogonal or symplectic bundles on a smooth projective variety. Combined with the contents of the present work, it will become possible to define motivic Donaldson–Thomas invariants for type B/C/D structure groups on a Calabi–Yau threefold, and write down wall-crossing formulae for these invariants. The author plans to report on this in a future paper.

1.1.7. Acknowledgements. The author would like to thank Andrés Ibáñez Núñez, Dominic Joyce, and Tasuki Kinjo, for helpful discussions and comments.

The author is grateful to the Mathematical Institute, University of Oxford, for its support during the preparation of this paper.

1.1.8. Conventions. Throughout this paper, we will use the following notations, terminology, and conventions.

- $\mathbb{K}$ is an algebraically closed field of characteristic zero.
- A $\mathbb{K}$ -variety is a separated $\mathbb{K}$ -scheme of finite type.
- A *reductive group* over $\mathbb{K}$ is a linear algebraic group over $\mathbb{K}$ that is linearly reductive, and is allowed to be disconnected.
- All $\mathbb{K}$ -schemes, *algebraic spaces* over $\mathbb{K}$, and *algebraic stacks* over $\mathbb{K}$ are assumed to be quasi-separated and locally of finite type. Algebraic stacks are assumed to have separated diagonal.
- A *derived algebraic stack* over $\mathbb{K}$ is a derived stack over $\mathbb{K}$ that has an open cover by *geometric stacks* in the sense of Toën–Vezzosi [51, §1.3.3]. Its *virtual dimension* refers to the rank of its cotangent complex, whenever this is defined.
- An *s-shifted symplectic stack* over $\mathbb{K}$, where $s \in \mathbb{Z}$, is a locally (homotopically) finitely presented derived algebraic stack over $\mathbb{K}$, equipped with an *s-shifted symplectic structure* in the sense of Pantev–Toën–Vaquié–Vezzosi [47, §1].

2 Motivic vanishing cycles

We provide background material on rings of motives in §2.1, and discuss their descent properties in §2.2. Then, in §2.3, we recall the construction of the *motivic Milnor fibre* from Denef–Loeser [20–22] and Looijenga [45], as well as its generalization as *nearby and vanishing cycle maps* on rings of motives defined by Bittner [9]. In §2.4, we extend this construction to define motivic nearby and vanishing cycle maps for algebraic stacks, and prove results that will be used in the proof of Theorem 4.1.1. Finally, in §2.5, we introduce the *motivic Behrend function* introduced by Bussi–Joyce–Meinhardt [14] and Ben-Bassat–Brav–Bussi–Joyce [7], slightly generalizing their construction by weakening the assumptions on the stack.

2.1 Rings of motives

2.1.1. We recall the construction of rings of motives over schemes, algebraic spaces, and algebraic stacks. The case of stacks was first due to Joyce [30], where the ring of motives was called the ring of *stack functions*.

To avoid repetition, we only state the majority of definitions and results for stacks, with the understanding that schemes and algebraic spaces are special cases of algebraic stacks.

2.1.2. Stacks with affine stabilizers. Let $\mathcal{X}$ be an algebraic stack over $\mathbb{K}$. We say that $\mathcal{X}$ has *affine stabilizers*, if for any field-valued point $x \in \mathcal{X}$, the stabilizer group of $\mathcal{X}$ at x is an affine algebraic group.

2.1.3. The ring of motives. Let $\mathcal{X}$ be an algebraic stack over $\mathbb{K}$ with affine stabilizers. Define the *Grothendieck ring of varieties* over $\mathcal{X}$ to be the abelian group

$$K_{\text{var}}(\mathcal{X}) = \bigoplus'_{Z \rightarrow \mathcal{X}} \mathbb{Z} \cdot [Z] \Big/ \sim, \quad (2.1)$$

where we run through all morphisms $Z \rightarrow \mathcal{X}$ with Z a $\mathbb{K}$ -variety, and $\bigoplus'$ denotes the set of *locally finite sums*, that is, possibly infinite sums $\sum_{Z \rightarrow \mathcal{X}} n_Z \cdot [Z]$, such that for each open substack $\mathcal{U} \subset \mathcal{X}$ of finite type, there are only finitely many Z such that $n_Z \neq 0$ and $Z \times_{\mathcal{X}} \mathcal{U} \neq \emptyset$. The relation $\sim$ is generated by $[Z] \sim [Z'] + [Z \setminus Z']$ for closed subschemes $Z' \subset Z$.

One can define multiplication on $K_{\text{var}}(\mathcal{X})$ by taking the fibre product over $\mathcal{X}$, making it into a commutative ring, possibly non-unital when $\mathcal{X}$ is not an algebraic space. It is also a (possibly non-unital) commutative $K_{\text{var}}(\mathbb{K})$ -algebra, where $K_{\text{var}}(\mathbb{K}) = K_{\text{var}}(\text{Spec } \mathbb{K})$, with the action given by the product.

Let $\mathbb{L} = [\mathbb{A}^1] \in K_{\text{var}}(\mathbb{K})$, and define *rings of motives* over $\mathcal{X}$,

$$M(\mathcal{X}) = K_{\text{var}}(\mathcal{X}) \otimes_{K_{\text{var}}(\mathbb{K})} K_{\text{var}}(\mathbb{K})[\mathbb{L}^{-1}] / (\mathbb{L} - 1)\text{-torsion}, \quad (2.2)$$

$$\hat{M}(\mathcal{X}) = K_{\text{var}}(\mathcal{X}) \otimes_{K_{\text{var}}(\mathbb{K})} K_{\text{var}}(\mathbb{K})[\mathbb{L}^{-1}, (\mathbb{L}^k - 1)^{-1}], \quad (2.3)$$

where we invert $\mathbb{L}^k - 1$ for all $k \geq 1$ in (2.3).

2.1.4. Motives of algebraic spaces and stacks. Let $\mathcal{X}$ be as above. For an algebraic space Z and a morphism $Z \rightarrow \mathcal{X}$ of finite type, one can also assign a class $[Z] \in K_{\text{var}}(\mathcal{X})$, extending the usual definition for varieties, since Z can be stratified by varieties.

Furthermore, as in Joyce [30] or Ben-Bassat–Brav–Bussi–Joyce [7, §5.3], for any finite type morphism of algebraic stacks $\mathcal{Z} \rightarrow \mathcal{X}$, where $\mathcal{Z}$ has affine stabilizers, one can assign a class $[\mathcal{Z}] \in \hat{M}(\mathcal{X})$, which agrees with the usual one when $\mathcal{Z}$ is a variety, and satisfies the relation $[\mathcal{Z}] = [\mathcal{Z}'] + [\mathcal{Z} \setminus \mathcal{Z}']$ for closed substacks $\mathcal{Z}' \subset \mathcal{Z}$. In particular, the class $[\mathcal{X}] \in \hat{M}(\mathcal{X})$ is the multiplicative unit of the ring $\hat{M}(\mathcal{X})$.

For an algebraic stack $\mathcal{X}$ over $\mathbb{K}$ of finite type, with affine stabilizers, one thus have a class $[\mathcal{X}] \in \hat{M}(\mathbb{K})$, called the *motive* of $\mathcal{X}$.

2.1.5. Pullbacks and pushforwards. Let $\mathcal{X}, \mathcal{Y}$ be algebraic stacks over $\mathbb{K}$ with affine stabilizers, and let $f: \mathcal{X} \rightarrow \mathcal{Y}$ be a morphism.

There is a pullback map

$$f^*: \hat{M}(\mathcal{Y}) \longrightarrow \hat{M}(\mathcal{X}), \quad (2.4)$$

which is an $\hat{M}(\mathbb{K})$ -algebra homomorphism, given on generators by $f^*[Z] = [Z \times_{\mathcal{Y}} \mathcal{X}]$, where the right-hand side is defined as in §2.1.4. Pulling back respects composition of morphisms. If, moreover, f is representable, then there are pullback maps

$$f^*: K_{\text{var}}(\mathcal{Y}) \longrightarrow K_{\text{var}}(\mathcal{X}), \quad (2.5)$$

$$f^*: M(\mathcal{Y}) \longrightarrow M(\mathcal{X}), \quad (2.6)$$

which are $K_{\text{var}}(\mathbb{K})$ - and $M(\mathbb{K})$ -algebra homomorphisms, respectively, and are defined similarly.

On the other hand, if f is of finite type, then there are pushforward maps

$$f_!: K_{\text{var}}(\mathcal{X}) \longrightarrow K_{\text{var}}(\mathcal{Y}), \quad (2.7)$$

$$f_!: M(\mathcal{X}) \longrightarrow M(\mathcal{Y}), \quad (2.8)$$

$$f_!: \hat{M}(\mathcal{X}) \longrightarrow \hat{M}(\mathcal{Y}), \quad (2.9)$$

which are $K_{\text{var}}(\mathbb{K})$ -, $M(\mathbb{K})$ -, and $\hat{M}(\mathbb{K})$ -module homomorphisms, respectively, given on generators by $f_![Z] = [Z]$. Pushing forward respects composition of morphisms.

In particular, when $\mathcal{X}$ is of finite type, pushing forward along the structure morphism $\mathcal{X} \rightarrow \text{Spec } \mathbb{K}$ is sometimes called *motivic integration*, and denoted by

$$\int_{\mathcal{X}} (-): \hat{M}(\mathcal{X}) \longrightarrow \hat{M}(\mathbb{K}). \quad (2.10)$$

2.1.6. Base change and projection formulae. Suppose we have a 2-pullback diagram

$$\begin{array}{ccc} \mathcal{X}' & \xrightarrow{f'} & \mathcal{Y}' \\ g' \downarrow & \ulcorner & \downarrow g \\ \mathcal{X} & \xrightarrow{f} & \mathcal{Y}, \end{array} \quad (2.11)$$

where $\mathcal{X}, \mathcal{Y}, \mathcal{X}', \mathcal{Y}'$ are algebraic stacks over $\mathbb{K}$ with affine stabilizers, and f is of finite type. Then we have the *base change formula*

$$g^* \circ f_! = f'_! \circ g'^* \quad (2.12)$$

on $\hat{M}(-)$, as in [30, Theorem 3.5]. Moreover, if g is representable, then this also holds for $K_{\text{var}}(-)$ and $M(-)$.

Let $f: \mathcal{X} \rightarrow \mathcal{Y}$ be as above. We have the *projection formula*

$$f_!(a \cdot f^*(b)) = f_!(a) \cdot b \quad (2.13)$$

for all $a \in \hat{M}(\mathcal{X})$ and $b \in \hat{M}(\mathcal{Y})$, which can be verified directly on generators. Moreover, if f is representable, then this also holds for $K_{\text{var}}(-)$ and $M(-)$.

2.1.7. Motives of principal bundles. Following Serre [49, §4], an algebraic group G over $\mathbb{K}$ is *special* if all principal G -bundles over a $\mathbb{K}$ -scheme are Zariski locally trivial. For example, the groups $\text{GL}(n)$ and $\mathbb{G}_a$ are special; semidirect products of special groups are special; disconnected groups are not special.

For a special group G and a principal G -bundle $\pi: \mathcal{Y} \rightarrow \mathcal{X}$, where $\mathcal{X}$ is an algebraic stack over $\mathbb{K}$ with affine stabilizers, we have the relation

$$\pi_! \circ \pi^* = [G] \cdot \text{id} \quad (2.14)$$

in $K_{\text{var}}(\mathcal{X})$, $M(\mathcal{X})$, and $\hat{M}(\mathcal{X})$, which follows from the definition of special groups. In particular, we have $[\mathcal{Y}] = [G] \cdot [\mathcal{X}]$ in $\hat{M}(\mathcal{X})$.

The relation (2.14) is not necessary true if G is not special. For example, consider the principal $\mathbb{Z}_2$ -bundle $\mathbb{G}_m \rightarrow \mathbb{G}_m$ given by $t \mapsto t^2$. Then the equality cannot hold, since $[\mathbb{G}_m] \neq 2 \cdot [\mathbb{G}_m]$.

2.1.8. The Euler characteristic. As in Joyce [30, Example 6.3], there is a ring map

$$\chi: M(\mathbb{K}) \longrightarrow \mathbb{Z}, \quad (2.15)$$

sending each generator $[Z]$ to its Euler characteristic, and sending $\mathbb{L}$ to 1. This extends naturally to a map $\chi: \hat{M}(\mathbb{K}) \rightarrow \mathbb{Q} \cup \{\infty\}$, sending $1/(1 + \mathbb{L} + \cdots + \mathbb{L}^{k-1})$ to $1/k$ for each $k \geq 1$, and sending elements not in $M(\mathbb{K})[(1 + \mathbb{L} + \cdots + \mathbb{L}^{k-1})^{-1} : k \geq 1]$ to ∞ .

For an algebraic stack $\mathcal{X}$ over $\mathbb{K}$ of finite type, with affine stabilizers, we have a map

$$\int_{\mathcal{X}} (-) d\chi: \hat{M}(\mathcal{X}) \longrightarrow \mathbb{Q} \cup \{\infty\}, \quad (2.16)$$

defined by pushing forward along $\mathcal{X} \rightarrow \text{Spec } \mathbb{K}$, and then taking the Euler characteristic. We have $\int_{\mathcal{X}} a d\chi \in \mathbb{Z}$ for all $a \in M(\mathcal{X})$.

2.1.9. Constructible functions. The ring of *constructible functions* on an algebraic stack is a coarser version of the rings of motives considered above, recording only the Euler characteristic of the motives. This was studied by Joyce [28; 30].

For an algebraic stack $\mathcal{X}$ over $\mathbb{K}$, a *constructible function* on $\mathcal{X}$ is a map of sets

$$a: |\mathcal{X}| \longrightarrow \mathbb{Z}, \quad (2.17)$$

where $|\mathcal{X}|$ is the underlying topological space of $\mathcal{X}$, as in Laumon–Moret-Bailly [43, Chapter 5], such that for any $c \in \mathbb{Z}$, the preimage $a^{-1}(c)$ is a locally constructible subset of $|\mathcal{X}|$. The abelian group of constructible functions on $\mathcal{X}$ is denoted by $\text{CF}(\mathcal{X})$.

There is an Euler characteristic map

$$\chi: \tilde{M}(\mathcal{X}) \longrightarrow \text{CF}(\mathcal{X}), \quad (2.18)$$

where $\tilde{M}(\mathcal{X}) \subset \hat{M}(\mathcal{X})$ is the $M(\mathbb{K})$ -subalgebra generated by classes $[\mathcal{Z}]$ for representable morphisms $\mathcal{Z} \rightarrow \mathcal{X}$, and χ is given by taking the fibrewise Euler characteristic, as in Joyce [30, Definition 3.2], where this map was denoted by $\pi_{\mathcal{X}}^{\text{stk}}$.

One can also define pullback and pushforward maps on $\text{CF}(-)$ for representable morphisms, where pushing forward needs the morphism to be of finite type. They are compatible with the map χ , and satisfy the base change and projection formulae as in §2.1.6, for representable morphisms.

2.1.10. The ring of monodromic motives. Let $\hat{\mu} = \lim \mu_n$ be the projective limit of the groups μ_n of roots of unity. For a $\mathbb{K}$ -scheme Z , a *good $\hat{\mu}$ -action* on Z is one that factors through μ_n for some n , such that each orbit is contained in an affine open subscheme.

Let $\mathcal{X}$ be an algebraic stack over $\mathbb{K}$ with affine stabilizers. Define the *monodromic Grothendieck ring of varieties* over $\mathcal{X}$ to be the abelian group

$$K_{\text{var}}^{\hat{\mu}}(\mathcal{X}) = \bigoplus'_{Z \rightarrow \mathcal{X}} \mathbb{Z} \cdot [Z]^{\hat{\mu}} / \sim, \quad (2.19)$$

where $\bigoplus'$ denotes the set of locally finite sums as in §2.1.3, and we sum over all morphisms $Z \rightarrow \mathcal{X}$ with Z a $\mathbb{K}$ -variety with a good $\hat{\mu}$ -action that is compatible with the trivial $\hat{\mu}$ -action on $\mathcal{X}$. The relation $\sim$ is generated by $[Z]^{\hat{\mu}} \sim [Z']^{\hat{\mu}} + [Z \setminus Z']^{\hat{\mu}}$ for $\hat{\mu}$ -invariant closed subschemes $Z' \subset Z$, and $[Z \times V]^{\hat{\mu}} \sim [Z \times \mathbb{A}^n]^{\hat{\mu}}$ for a $\hat{\mu}$ -representation V of dimension n , where the projections to $\mathcal{X}$ factor through Z , and $\hat{\mu}$ acts on $\mathbb{A}^n$ trivially. See Looijenga [45, §5] and Ben-Bassat–Brav–Bussi–Joyce [7, §5].

Using the $K_{\text{var}}(\mathbb{K})$ -module structure on $K_{\text{var}}^{\hat{\mu}}(\mathcal{X})$, we define *rings of monodromic*

motives over $\mathcal{X}$,

$$M^{\hat{\mu}}(\mathcal{X}) = K_{\text{var}}^{\hat{\mu}}(\mathcal{X}) \otimes_{K_{\text{var}}(\mathbb{K})} K_{\text{var}}(\mathbb{K}) [\mathbb{L}^{-1}] / (\mathbb{L} - 1)\text{-torsion}, \quad (2.20)$$

$$\hat{M}^{\hat{\mu}}(\mathcal{X}) = K_{\text{var}}^{\hat{\mu}}(\mathcal{X}) \otimes_{K_{\text{var}}(\mathbb{K})} K_{\text{var}}(\mathbb{K}) [\mathbb{L}^{-1}, (\mathbb{L}^k - 1)^{-1}] / \approx, \quad (2.21)$$

where the ring structures and the relation $\approx$ are defined below.

We consider multiplication on $K_{\text{var}}^{\hat{\mu}}(\mathcal{X})$ denoted by ‘ $\odot$ ’ in [7, Definition 5.3]; see there for the definition. We will denote this by ‘ $\cdot$ ’. This makes $K_{\text{var}}^{\hat{\mu}}(\mathcal{X})$ into a ring, possibly non-unital when $\mathcal{X}$ is not an algebraic space. Note that this is *not* given by the fibre product, although the latter does define a different ring structure. The relation $\approx$ is defined as in [7, Definitions 5.5 and 5.13], denoted by ‘ $I_{\mathcal{X}}^{\text{st}, \hat{\mu}}$ ’ there, and is imposed so that the map Υ in §2.1.11 below respects the tensor product.

There is an element

$$\mathbb{L}^{1/2} = 1 - [\mu_2]^{\hat{\mu}} \in K_{\text{var}}^{\hat{\mu}}(\mathbb{K}), \quad (2.22)$$

where $\hat{\mu}$ acts on μ_2 non-trivially. It satisfies $(\mathbb{L}^{1/2})^2 = \mathbb{L}$. We also write $\mathbb{L}^{-1/2} = \mathbb{L}^{-1} \cdot \mathbb{L}^{1/2} \in M^{\hat{\mu}}(\mathbb{K})$.

There are the inclusion maps

$$\iota^{\hat{\mu}}: K_{\text{var}}(\mathcal{X}) \longrightarrow K_{\text{var}}^{\hat{\mu}}(\mathcal{X}), \quad \iota^{\hat{\mu}}: M(\mathcal{X}) \longrightarrow M^{\hat{\mu}}(\mathcal{X}), \quad \iota^{\hat{\mu}}: \hat{M}(\mathcal{X}) \longrightarrow \hat{M}^{\hat{\mu}}(\mathcal{X}), \quad (2.23)$$

given on generators by $[Z] \mapsto [Z]$, with the trivial $\hat{\mu}$ -action on Z . They are $K_{\text{var}}(\mathbb{K})$ -, $M(\mathbb{K})$ -, and $\hat{M}(\mathbb{K})$ -algebra homomorphisms, respectively.

One can define pullback and pushforward maps on $K_{\text{var}}^{\hat{\mu}}(-)$, $M^{\hat{\mu}}(-)$, and $\hat{M}^{\hat{\mu}}(-)$, similar to the case of $K_{\text{var}}(-)$, $M(-)$, and $\hat{M}(-)$. They satisfy the base change and projection formulae in §2.1.6, and the principal bundle relation in §2.1.7.

There is also the Euler characteristic map $\chi: M^{\hat{\mu}}(\mathbb{K}) \rightarrow \mathbb{Z}$, which is a ring homomorphism, from which one can define analogues of operations in §§2.1.8 and 2.1.9 for $M^{\hat{\mu}}(-)$ and $\hat{M}^{\hat{\mu}}(-)$. In particular, we have $\chi(\mathbb{L}^{1/2}) = -1$.

2.1.11. Motives of double covers. For a principal μ_2 -bundle $\mathcal{P} \rightarrow \mathcal{X}$, there is a class

$$\Upsilon(\mathcal{P}) = \mathbb{L}^{-1/2} \cdot ([\mathcal{X}] - [\mathcal{P}]^{\hat{\mu}}) \in \hat{M}^{\hat{\mu}}(\mathcal{X}), \quad (2.24)$$

where $\hat{\mu}$ acts on $\mathcal{P}$ via the μ_2 -action, and the class $[\mathcal{P}]^{\hat{\mu}}$ is defined similarly to §2.1.4 when $\mathcal{P}$ is not a variety. See [7, Definitions 5.5 and 5.13] for more details.

Note that Υ commutes with pullbacks by definition. Also, we have the relation

$$\Upsilon(\mathcal{P}_1 \otimes \mathcal{P}_2) = \Upsilon(\mathcal{P}_1) \cdot \Upsilon(\mathcal{P}_2) \quad (2.25)$$

for principal μ_2 -bundles $\mathcal{P}_1, \mathcal{P}_2 \rightarrow \mathcal{X}$, where $\mathcal{P}_1 \otimes \mathcal{P}_2$ is also a principal μ_2 -bundle. See [7, Definition 5.5 and 5.13].

2.2 Descent of motives

2.2.1. We now discuss descent properties of the rings of motives defined above. While constructible functions $\text{CF}(-)$ descend under any reasonable topology, descent for rings of motives such as $\hat{M}(-)$ and $\hat{M}(-)$ is more subtle. For example, pulling back along the double cover $\mathbb{G}_m \rightarrow \mathbb{G}_m, t \mapsto t^2$ is *not* injective on motives, since the class of the trivial double cover $\mathbb{G}_m \times \mu_2 \rightarrow \mathbb{G}_m$ and the non-trivial double cover $\mathbb{G}_m \rightarrow \mathbb{G}_m$ gets identified after pulling back. Therefore, rings of motives do *not* satisfy étale descent.

However, we show in [Theorem 2.2.3](#) below that these rings of motives do satisfy descent under the Nisnevich topology.

2.2.2. **The Nisnevich topology.** Recall that a family of morphisms of algebraic spaces $(f_i: X_i \rightarrow X)_{i \in I}$ is a *Nisnevich cover*, if each f_i is étale, and for each field-valued point $x \in X$, there exists $i \in I$ and a point $x' \in X_i$, such that $f_i(x') = x$, and f induces an isomorphism on residue fields at x' and x .

Let $\mathcal{X}$ be an algebraic stack. We define a *Nisnevich cover* of $\mathcal{X}$ to be a representable étale cover $(f_i: \mathcal{X}_i \rightarrow \mathcal{X})_{i \in I}$ such that its base change to any algebraic space is a Nisnevich cover of algebraic spaces. See, for example, [17, Definition 1.2 ff.].

For example, for an integer $n > 1$, the morphism $*$ $\rightarrow$ $[*/\mu_n]$ is *not* a Nisnevich cover, since its base change $\mathbb{G}_m \rightarrow \mathbb{G}_m, t \mapsto t^n$ is not a Nisnevich cover.

Algebraic spaces over $\mathbb{K}$ (assumed locally of finite type) admit Nisnevich covers by affine $\mathbb{K}$ -varieties, as can be deduced from Knutson [38, II, Theorem 6.4].

2.2.3. **Theorem.** *Let $\mathcal{X}$ be an algebraic stack over $\mathbb{K}$ with affine stabilizers, and let $(f_i: \mathcal{X}_i \rightarrow \mathcal{X})_{i \in I}$ be a Nisnevich cover. Then the map*

$$(f_i^*)_{i \in I}: \hat{M}(\mathcal{X}) \longrightarrow \text{eq} \left(\prod_{i \in I} \hat{M}(\mathcal{X}_i) \rightrightarrows \prod_{i, j \in I} \hat{M}(\mathcal{X}_i \times_{\mathcal{X}} \mathcal{X}_j) \right) \quad (2.26)$$

is an isomorphism, where the right-hand side is the equalizer of the two maps induced by pulling back along projections from each $\mathcal{X}_i \times_{\mathcal{X}} \mathcal{X}_j$ to $\mathcal{X}_i$ and $\mathcal{X}_j$, respectively.

The same also holds for $\hat{M}^{\text{h}}(-)$ in place of $\hat{M}(-)$. Moreover, if $\mathcal{X}$ is an algebraic space, then the same holds for $K_{\text{var}}(-)$, $M(-)$, $K_{\text{var}}^{\text{h}}(-)$, and $M^{\text{h}}(-)$.

Proof. We write down the proof for $\hat{M}(-)$, and the other cases are similar.

We first consider the case when $\mathcal{X}$ is an algebraic space. In this case, one can stratify $\mathcal{X}$ into locally closed pieces $S_k \subset \mathcal{X}$, such that the map $\coprod_i \mathcal{X}_i \rightarrow \mathcal{X}$ admits a section s_k over each S_k . After a base change to each S_k , we can assume that $\coprod_i \mathcal{X}_i \rightarrow \mathcal{X}$ admits a global section, in which case the result is clear.

For the general case, by Kresch [42, Proposition 3.5.9], $\mathcal{X}$ can be stratified by quotient stacks of the form $[U/G]$, where U is a quasi-projective $\mathbb{K}$ -variety acted on by $G \simeq \text{GL}(n)$ for some n . Therefore, we may assume that $\mathcal{X} = [U/G]$ is of this form. Let $\pi: U \rightarrow [U/G]$ be the projection. Then for all $a \in \hat{M}([U/G])$, we have $a = [G]^{-1} \cdot \pi_! \circ \pi^*(a)$, so that π^* is injective. Its image consists of elements $\tilde{a} \in \hat{M}(U)$ such that $\pi^* \circ \pi_!(\tilde{a}) = [G] \cdot \tilde{a}$. We call such elements *G-invariant*. In other words,

we may identify $\hat{M}([U/G])$ with the subring of $\hat{M}(U)$ consisting of G -invariant elements. Writing $U_i = U \times_{\mathcal{X}} \mathcal{X}_i$, it suffices to show that $\hat{M}(U) \xrightarrow{\sim} \text{eq}(\prod_{i \in I} \hat{M}(U_i) \rightrightarrows \prod_{i,j \in I} \hat{M}(U_i \times_U U_j))$, since taking G -invariant elements on both sides gives the desired result. We are now reduced to the already known case of algebraic spaces. $\square$

2.2.4. Quotient stacks and fundamental stacks. We now discuss classes of algebraic stacks that can be covered by quotient stacks, which will be used in the sequel. See §2.2.5 below for examples of stacks satisfying these properties.

A *quotient stack* over $\mathbb{K}$ is an algebraic stack of the form $[U/G]$, where U is an algebraic space over $\mathbb{K}$ acted on by an algebraic group $G \simeq \text{GL}(n)$ for some n . Equivalently, one can allow G to be any linear algebraic group, since if one chooses an embedding $G \hookrightarrow \text{GL}(n)$, then $[U/G] \simeq [(U \times^G \text{GL}(n))/\text{GL}(n)]$.

A *fundamental stack* over $\mathbb{K}$ is a quotient stack of the form $[U/G]$, where U is an affine $\mathbb{K}$ -variety acted on by an algebraic group $G \simeq \text{GL}(n)$ for some n . Equivalently, one can allow G to be any reductive group, by a similar argument as above; see the proof of [2, Corollary 4.14]. This terminology is from Alper–Hall–Rydh [3].

An algebraic stack over $\mathbb{K}$ is *Nisnevich locally a quotient stack*, if it admits a Nisnevich cover by quotient stacks. This class of stacks is also discussed in Choudhury–Deshmukh–Hogadi [17], where they are called *cd-quotient stacks*.

An algebraic stack over $\mathbb{K}$ is *Nisnevich locally fundamental*, if it admits a Nisnevich cover by fundamental stacks. This implies being Nisnevich locally a quotient stack.

Similarly, we define stacks that are *étale locally a quotient stack* or *étale locally fundamental*, by requiring representable étale covers with the corresponding properties.

These properties are satisfied by a large class of stacks, which we discuss below.

2.2.5. Local structure theorems. A series of local structure theorems for algebraic stacks by Alper–Hall–Rydh [2; 3] can be used to produce covers of algebraic stacks by quotient and fundamental stacks.

Theorem. *Let $\mathcal{X}$ be an algebraic stack with affine stabilizers.*

- (i) *If closed points of $\mathcal{X}$ have reductive stabilizers, then $\mathcal{X}$ is étale locally fundamental.*
- (ii) *If $\mathcal{X}$ admits a good moduli space in the sense of Alper [1], then $\mathcal{X}$ is Nisnevich locally fundamental.*

The first result follows from [2, Theorem 1.1], and is stated in Alper–Halpern-Leistner–Heinloth [4, Remarks 2.6 and 2.7]. The second result is [3, Theorem 6.1].

2.3 Motivic vanishing cycles for schemes

2.3.1. Idea. The motivic nearby and vanishing cycle maps considered below are based on the idea of *Milnor fibres*, which is perhaps more straightforward to explain in the analytic setting.

For this purpose, let X be a complex manifold, with a smooth metric $d: X \times X \rightarrow \mathbb{R}_{\geq 0}$, and let $f: X \rightarrow \mathbb{C}$ be a holomorphic function. Let $x \in X$ be a point such that $f(x) = 0$. Let $0 < \delta \ll \varepsilon \ll 1$ be small positive numbers, and consider the map

$$X_{\delta, \varepsilon}^\times(x) = B_\varepsilon(x) \cap f^{-1}(D_\delta^\times) \xrightarrow{f} D_\delta^\times, \quad (2.27)$$

where $D_\delta^\times = \{z \in \mathbb{C} \mid 0 < |z| < \delta\}$. This map is a topological fibration, and its fibre $\mathrm{MF}_f(x)$ is called the *Milnor fibre* of f at x . The cohomology of $\mathrm{MF}_f(x)$ carries the action of the *monodromy operator* induced by this fibration.

We will consider below the motivic analogue of the above construction, with X a smooth variety over a field $\mathbb{K}$. The Milnor fibre is then replaced by the *motivic Milnor fibre* of Denef–Loeser [20–22], which is a monodromic motive on X_0 . This construction is closely related to Donaldson–Thomas theory; see Behrend [6], Joyce–Song [35, §4], and Kontsevich–Soibelman [40, §4] for more details.

2.3.2. The motivic Milnor fibre. Let X be a smooth, irreducible $\mathbb{K}$ -variety, and let $f: X \rightarrow \mathbb{A}^1$ be a morphism. Write $X_0 = f^{-1}(0)$. Following Denef–Loeser [20–22], we define the *motivic Milnor fibre* of f , which is an element

$$\mathrm{MF}_f \in \mathcal{M}^{\hat{\mu}}(X_0), \quad (2.28)$$

as follows.

If f is constant, define $\mathrm{MF}_f = 0$. Otherwise, choose a resolution $\pi: \tilde{X} \rightarrow X$ of f , in the sense that $\tilde{X}$ is a smooth, irreducible $\mathbb{K}$ -variety, π is a proper morphism that restricts to an isomorphism on $\pi^{-1}(X \setminus X_0)$, and $\pi^{-1}(X_0)$ is a simple normal crossings divisor in $\tilde{X}$. See, for example, Kollár [39] for the existence and properties of such resolutions.

Let $(E_i)_{i \in J}$ be the irreducible components of $\pi^{-1}(X_0)$, and write N_i for the multiplicity of E_i in the divisor of $f \circ \pi$ on $\tilde{X}$. For a non-empty subset $I \subset J$, write $E_I^\circ = \bigcap_{i \in I} E_i \setminus \bigcup_{i \notin I} E_i$. Let $m_I = \gcd_{i \in I} N_i$, and define an m_I -fold cover $\tilde{E}_I^\circ \rightarrow E_I^\circ$ as follows. For each open set $U \subset \tilde{X}$ such that $f \circ \pi = uv^{m_I}$ on U for $u: U \rightarrow \mathbb{A}^1 \setminus \{0\}$ and $v: U \rightarrow \mathbb{A}^1$, define the restriction of $\tilde{E}_I^\circ$ on $E_I^\circ \cap U$ as

$$\tilde{E}_I^\circ|_{E_I^\circ \cap U} = \{(z, y) \in \mathbb{A}^1 \times (E_I^\circ \cap U) \mid z^{m_I} = u^{-1}\}. \quad (2.29)$$

Since E_I° can be covered by such open sets U , (2.29) can be glued together to obtain a cover $\tilde{E}_I^\circ \rightarrow E_I^\circ$, with a natural μ_{m_I} -action given by scaling the z -coordinate, which induces a $\hat{\mu}$ -action on $\tilde{E}_I^\circ$. The motivic Milnor fibre MF_f is then given by

$$\mathrm{MF}_f = \sum_{\emptyset \neq I \subset J} (1 - \mathbb{L})^{|I|-1} [\tilde{E}_I^\circ]^{\hat{\mu}}. \quad (2.30)$$

It can be shown [21, Definition 3.8] that this is independent of the choice of the resolution π .

2.3.3. Nearby and vanishing cycles. Let X be a $\mathbb{K}$ -variety, and let $f: X \rightarrow \mathbb{A}^1$ a morphism. Write $X_0 = f^{-1}(0)$.

Define the *nearby cycle map* of f , denoted by

$$\Psi_f: M(X) \longrightarrow M^{\hat{\mu}}(X_0), \quad (2.31)$$

to be the unique $M(\mathbb{K})$ -linear map such that for any smooth, irreducible $\mathbb{K}$ -variety Z and any proper morphism $\rho: Z \rightarrow X$, we have

$$\Psi_f([Z]) = (\rho_0)_! (MF_{f \circ \rho}) \in M^{\hat{\mu}}(X_0), \quad (2.32)$$

where $\rho_0: Z_0 \rightarrow X_0$ is the restriction of ρ to $Z_0 = (f \circ \rho)^{-1}(0)$, and $MF_{f \circ \rho} \in M^{\hat{\mu}}(Z_0)$ is the motivic Milnor fibre of $f \circ \rho$. It follows from Bittner [9, Claim 8.2] that the map Ψ_f is well-defined.

Define the *vanishing cycle map* of f to be the map

$$\Phi_f = \Psi_f - \iota^{\hat{\mu}} \circ i^*: M(X) \longrightarrow M^{\hat{\mu}}(X_0), \quad (2.33)$$

where $i: X_0 \hookrightarrow X$ is the inclusion, and $\iota^{\hat{\mu}}: M(X_0) \hookrightarrow M^{\hat{\mu}}(X_0)$ is as in §2.1.10.

2.3.4. For algebraic spaces. As in Bittner [9, Theorem 8.4], the nearby and vanishing cycle maps are compatible with pulling back along smooth morphisms. In particular, they define morphisms $\Psi, \Phi: M(-) \rightarrow M^{\hat{\mu}}((-)_0)$ of sheaves on the category of $\mathbb{K}$ -varieties over $\mathbb{A}^1$, with the Nisnevich topology. Since algebraic spaces admit Nisnevich covers by affine $\mathbb{K}$ -varieties, as mentioned in §2.2.2, these morphisms of sheaves induce maps on their evaluations on algebraic spaces over $\mathbb{K}$.

In other words, for an algebraic space X over $\mathbb{K}$ and a morphism $f: X \rightarrow \mathbb{A}^1$, we have defined nearby and vanishing cycle maps

$$\Psi_f, \Phi_f: M(X) \longrightarrow M^{\hat{\mu}}(X_0). \quad (2.34)$$

We state some of their properties below.

2.3.5. Theorem. *Let X, Y be algebraic spaces over $\mathbb{K}$.*

- (i) *Let $g: Y \rightarrow X$ be a proper morphism, and $f: X \rightarrow \mathbb{A}^1$ a morphism. Then we have a commutative diagram*

$$\begin{array}{ccc} M(Y) & \xrightarrow{g_!} & M(X) \\ \Psi_{f \circ g} \downarrow & & \downarrow \Psi_f \\ M^{\hat{\mu}}(Y_0) & \xrightarrow{g_!} & M^{\hat{\mu}}(X_0). \end{array} \quad (2.35)$$

- (ii) *Let $g: Y \rightarrow X$ be a smooth morphism, and $f: X \rightarrow \mathbb{A}^1$ a morphism. Then we*

have a commutative diagram

$$\begin{array}{ccc}
M(X) & \xrightarrow{g^*} & M(Y) \\
\Psi_f \downarrow & & \downarrow \Psi_{f \circ g} \\
M^{\hat{\mu}}(X_0) & \xrightarrow{g^*} & M^{\hat{\mu}}(Y_0) .
\end{array} \tag{2.36}$$

Proof. The case when X and Y are $\mathbb{K}$ -varieties was proved in Bittner [9, Theorem 8.4]. The verification of (ii) for algebraic spaces is completely formal, by passing to Nisnevich covers by $\mathbb{K}$ -varieties.

We now prove (i) for algebraic spaces. Again, passing to a Nisnevich cover, we may assume that X is a $\mathbb{K}$ -variety. We claim that $K_{\text{var}}(Y)$ is spanned over $\mathbb{Z}$ by classes $[Z]$ of proper morphisms $Z \rightarrow Y$, where Z is a smooth $\mathbb{K}$ -variety. Indeed, let $u: U \rightarrow Y$ be an arbitrary morphism, where U is an integral $\mathbb{K}$ -variety. By Nagata compactification, as in Conrad–Lieblich–Olsson [19, Theorem 1.2.1], u can be factored as a dense open immersion $U \hookrightarrow V$ followed by a proper morphism $V \rightarrow Y$, where V is an integral algebraic space over $\mathbb{K}$. By Chow’s lemma for algebraic spaces, as in Knutson [38, IV, Theorem 3.1], there exists a $\mathbb{K}$ -variety W and a projective birational morphism $W \rightarrow V$. Applying a resolution of singularities, we may assume that W is smooth. Now $W \rightarrow Y$ is proper, and the difference $[W] - [U]$ is a sum of lower dimensional classes. An induction on the dimension of U verifies the claim.

Now, let $h: Z \rightarrow Y$ be a proper morphism, where Z is a smooth $\mathbb{K}$ -variety. Passing to a Nisnevich cover of Y by $\mathbb{K}$ -varieties, one can show that $\Psi_{f \circ g}([Z]) = h_*(\text{MF}_{f \circ g \circ h})$. On the other hand, we have $\Psi_f([Z]) = (g \circ h)_*(\text{MF}_{f \circ g \circ h})$ by definition. This completes the proof since such classes $[Z]$ span $K_{\text{var}}(Y)$, so they also span $M(Y)$ over $M(\mathbb{K})$. $\square$

2.4 Motivic vanishing cycles for stacks

2.4.1. Assumptions on the stack. From now on, we assume that $\mathcal{X}$ is an algebraic stack over $\mathbb{K}$ that is Nisnevich locally a quotient stack in the sense of §2.2.4.

Note that this assumption is weaker than that in Ben-Bassat–Brav–Bussi–Joyce [7, §5], where the stack was assumed to be Zariski locally a quotient stack.

2.4.2. Theorem. *Let $\mathcal{X}$ be as in §2.4.1, and let $f: \mathcal{X} \rightarrow \mathbb{A}^1$ be a morphism. Write $\mathcal{X}_0 = f^{-1}(0)$. Then there is a unique $\hat{M}(\mathbb{K})$ -linear map*

$$\Psi_f: \hat{M}(\mathcal{X}) \longrightarrow \hat{M}^{\hat{\mu}}(\mathcal{X}_0) , \tag{2.37}$$

called the nearby cycle map of f , such that for any $\mathbb{K}$ -variety Y and any smooth morphism

ism $g: Y \rightarrow \mathcal{X}$, we have a commutative diagram

$$\begin{array}{ccc} \hat{M}(\mathcal{X}) & \xrightarrow{g^*} & \hat{M}(Y) \\ \Psi_f \downarrow & & \downarrow \Psi_{f \circ g} \\ \hat{M}^{\hat{\mu}}(\mathcal{X}_0) & \xrightarrow{g^*} & \hat{M}^{\hat{\mu}}(Y_0) . \end{array} \quad (2.38)$$

where the right-hand map is defined in §2.3.3.

We then define the vanishing cycle map of f to be the map

$$\Phi_f = \Psi_f - \iota^{\hat{\mu}} \circ i^* : \hat{M}(\mathcal{X}) \longrightarrow \hat{M}^{\hat{\mu}}(\mathcal{X}_0) , \quad (2.39)$$

where $i: \mathcal{X}_0 \hookrightarrow \mathcal{X}$ and $\iota^{\hat{\mu}}: \hat{M}(\mathcal{X}_0) \hookrightarrow \hat{M}^{\hat{\mu}}(\mathcal{X}_0)$ are the inclusions.

Proof. Let $(j_i: \mathcal{X}_i \rightarrow \mathcal{X})_{i \in I}$ be a Nisnevich cover, where each $\mathcal{X}_i \simeq [U_i/G_i]$, with U_i an algebraic space over $\mathbb{K}$, acted on by a group $G_i \simeq \mathrm{GL}(n_i)$ for some n_i . Let $\pi_i: U_i \rightarrow \mathcal{X}_i$ be the projection.

First, note that the condition on Ψ_f implies that the same condition holds when Y is an algebraic space, with the right-hand map in (2.38) defined in §2.3.4. This can be seen by passing to a Nisnevich cover of Y by $\mathbb{K}$ -varieties, and applying Theorem 2.2.3 to this cover.

To define the map Ψ_f , by Theorem 2.2.3, it is enough to define it on each $\mathcal{X}_i$, and then verify that they agree on overlaps. Let $a \in \hat{M}(\mathcal{X})$ be an element. We define the element $\Psi_f(a) \in \hat{M}^{\hat{\mu}}(\mathcal{X}_0)$ by giving its pullbacks $\Psi_f(a)_i = j_i^* \circ \Psi_f(a) \in \hat{M}^{\hat{\mu}}(\mathcal{X}_{i,0})$ for each i , where $\mathcal{X}_{i,0} = \mathcal{X}_i \times_{\mathcal{X}} \mathcal{X}_0$. The condition on Ψ_f forces

$$\begin{aligned} \Psi_f(a)_i &= j_i^* \circ \Psi_f(a) = [G_i]^{-1} \cdot (\pi_i)_! \circ \pi_i^* \circ j_i^* \circ \Psi_f(a) \\ &= [G_i]^{-1} \cdot (\pi_i)_! \circ \Psi_{f \circ j_i \circ \pi_i} \circ \pi_i^* \circ j_i^*(a) , \end{aligned} \quad (2.40)$$

where $[G_i] \in \hat{M}(\mathbb{K})$ is the class of G_i and is invertible in $\hat{M}(\mathbb{K})$, and we applied (2.14) to π_i , using the fact that G_i is special. This shows that if the map Ψ_f exists, then it is unique.

To check that the elements $\Psi_f(a)_i$ agree on overlaps, let $1, 2 \in I$ be two indices, and form the pullback squares

$$\begin{array}{ccccc} U'' & \xrightarrow{\pi_2''} & U_1' & \xrightarrow{j_2''} & U_1 \\ \pi_1'' \downarrow & \ulcorner & \pi_1' \downarrow & \ulcorner & \downarrow \pi_1 \\ U_2' & \xrightarrow{\pi_2'} & \mathcal{X}_{1,2} & \xrightarrow{j_2'} & \mathcal{X}_1 \\ j_1'' \downarrow & \ulcorner & j_1' \downarrow & \ulcorner & \downarrow j_1 \\ U_2 & \xrightarrow{\pi_2} & \mathcal{X}_2 & \xrightarrow{j_2} & \mathcal{X} , \end{array} \quad (2.41)$$

where U_1' , U_2' , and U'' are algebraic spaces. We need to show that $(j_2')^*(\Psi_f(a)_1) =$

$(j'_1)^*(\Psi_f(a)_2)$. We have

$$\begin{aligned}
& (j'_2)^*(\Psi_f(a)_1) \\
&= [G_1]^{-1} \cdot (j'_2)^* \circ (\pi_1)_! \circ \Psi_{f \circ j_1 \circ \pi_1} \circ (j_1 \circ \pi_1)^*(a) \\
&= [G_1]^{-1} \cdot (\pi'_1)_! \circ (j'_2)^* \circ \Psi_{f \circ j_1 \circ \pi_1} \circ (j_1 \circ \pi_1)^*(a) \\
&= [G_1]^{-1} \cdot (\pi'_1)_! \circ \Psi_{f \circ j_1 \circ \pi_1 \circ j'_2} \circ (j_1 \circ \pi_1 \circ j'_2)^*(a) \\
&= [G_1]^{-1} \cdot [G_2]^{-1} \cdot (\pi'_1)_! \circ (\pi'_2)_! \circ (\pi'_2)^* \circ \Psi_{f \circ j_1 \circ \pi_1 \circ j'_2} \circ (j_1 \circ \pi_1 \circ j'_2)^*(a) \\
&= [G_1]^{-1} \cdot [G_2]^{-1} \cdot (\pi'_1 \circ \pi'_2)_! \circ \Psi_{f \circ j_1 \circ \pi_1 \circ j'_2 \circ \pi'_2} \circ (j_1 \circ \pi_1 \circ j'_2 \circ \pi'_2)^*(a), \quad (2.42)
\end{aligned}$$

where we applied (2.12) in the second step, Theorem 2.3.5 (ii) in the third and fifth steps, and §2.1.7 in the fourth step. This expression is now symmetric in the indices 1 and 2, so the element $\Psi_f(a)$ is well-defined.

It now remains to show that the map Ψ_f satisfies the required condition. Let Y be a $\mathbb{K}$ -variety and $\pi: Y \rightarrow \mathcal{X}$ a smooth morphism. For each $i \in I$, write $Y_i = Y \times_{\mathcal{X}} \mathcal{X}_i$. Then $(k_i: Y_i \rightarrow Y)_{i \in I}$ is a Nisnevich cover by algebraic spaces, and it suffices to show that

$$k_i^* \circ g^* \circ \Psi_f = k_i^* \circ \Psi_{f \circ g} \circ g^* \quad (2.43)$$

for each i . Consider the diagram

$$\begin{array}{ccccc}
V_i & \xrightarrow{\rho_i} & Y_i & \xrightarrow{k_i} & Y \\
g'_i \downarrow & \lrcorner & g_i \downarrow & \lrcorner & \downarrow g \\
U_i & \xrightarrow{\pi_i} & \mathcal{X}_i & \xrightarrow{j_i} & \mathcal{X},
\end{array} \quad (2.44)$$

where all squares are pullback squares. In particular, ρ_i is a principal G_i -bundle. For any $a \in \hat{M}(\mathcal{X})$, we have

$$\begin{aligned}
& k_i^* \circ g^* \circ \Psi_f(a) \\
&= g_i^*(\Psi_f(a)_i) \\
&= [G_i]^{-1} \cdot g_i^* \circ (\pi_i)_! \circ \Psi_{f \circ j_i \circ \pi_i} \circ (j_i \circ \pi_i)^*(a) \\
&= [G_i]^{-1} \cdot (\rho_i)_! \circ (g'_i)^* \circ \Psi_{f \circ j_i \circ \pi_i} \circ (j_i \circ \pi_i)^*(a) \\
&= [G_i]^{-1} \cdot (\rho_i)_! \circ \Psi_{f \circ j_i \circ \pi_i \circ g'_i} \circ (j_i \circ \pi_i \circ g'_i)^*(a) \\
&= [G_i]^{-1} \cdot (\rho_i)_! \circ \Psi_{f \circ g \circ k_i \circ \rho_i} \circ (g \circ k_i \circ \rho_i)^*(a) \\
&= [G_i]^{-1} \cdot (\rho_i)_! \circ \rho_i^* \circ \Psi_{f \circ g \circ k_i} \circ (g \circ k_i)^*(a) \\
&= \Psi_{f \circ g \circ k_i} \circ (g \circ k_i)^*(a) \\
&= k_i^* \circ \Psi_{f \circ g} \circ g^*(a), \quad (2.45)
\end{aligned}$$

where we applied (2.12) in the third step, Theorem 2.3.5 (ii) in the fourth, sixth, and eighth steps, and §2.1.7 in the seventh step. This proves the desired identity (2.43). $\square$

2.4.3. The motivic Milnor fibre. Let $\mathcal{X}$ be as in §2.4.1, and let $f: \mathcal{X} \rightarrow \mathbb{A}^1$ be a morphism. Write $\mathcal{X}_0 = f^{-1}(0)$. The *motivic Milnor fibre* of f is the element

$$\mathrm{MF}_f = \Psi_f([\mathcal{X}]) \in \hat{M}^{\hat{\mu}}(\mathcal{X}_0). \quad (2.46)$$

When $\mathcal{X}$ is smooth, this is closely related to the construction of Ben-Bassat–Brav–Bussi–Joyce [7, §5.4], which we will further discuss and generalize in §2.5.4 below. The main difference is that the latter construction starts from the critical locus of f instead of $\mathcal{X}$, and can be generalized to stacks glued from such critical loci; it uses Φ_f instead of Ψ_f , and introduces a twist by a power of $\mathbb{L}^{1/2}$ to make gluing possible.

We relate this to the description of the motivic Milnor fibre for schemes in §2.3.2. Suppose that we are given a *resolution* of f , which is a representable proper morphism $\pi: \tilde{\mathcal{X}} \rightarrow \mathcal{X}$, such that it restricts to an isomorphism on $\pi^{-1}(\mathcal{X} \setminus \mathcal{X}_0)$, and $\pi^{-1}(\mathcal{X}_0)$ is a simple normal crossings divisor in $\tilde{\mathcal{X}}$, in the sense that it is so after pulling back along smooth morphisms from schemes. Let $(\mathcal{E}_i)_{i \in J}$ be the family of irreducible components of $\pi^{-1}(\mathcal{X}_0)$, and define $\mathcal{E}_I^\circ$ and $\tilde{\mathcal{E}}_I^\circ$ for non-empty $I \subset J$ similarly to §2.3.2, where $\tilde{\mathcal{E}}_I^\circ$ carries a natural $\hat{\mu}$ -action. We then claim that

$$\mathrm{MF}_f = \sum_{\emptyset \neq I \subset J} (1 - \mathbb{L})^{|I|-1} [\tilde{\mathcal{E}}_I^\circ]^{\hat{\mu}}. \quad (2.47)$$

Indeed, this can be shown by a similar argument as in the proof of Theorem 2.4.2, by first passing to a Nisnevich cover by quotient stacks, then using the relation (2.14) to further reduce to the case of algebraic spaces, and finally passing to a Nisnevich cover again to reduce to the case of affine varieties.

2.4.4. Theorem. *Let $\mathcal{X}, \mathcal{Y}$ be algebraic stacks as in §2.4.1.*

- (i) *Let $g: \mathcal{Y} \rightarrow \mathcal{X}$ be a proper morphism, and $f: \mathcal{X} \rightarrow \mathbb{A}^1$ a morphism. Then we have a commutative diagram*

$$\begin{array}{ccc} \hat{M}(\mathcal{Y}) & \xrightarrow{g!} & \hat{M}(\mathcal{X}) \\ \Psi_{f \circ g} \downarrow & & \downarrow \Psi_f \\ \hat{M}^{\hat{\mu}}(\mathcal{Y}_0) & \xrightarrow{g!} & \hat{M}^{\hat{\mu}}(\mathcal{X}_0). \end{array} \quad (2.48)$$

- (ii) *Let $g: \mathcal{Y} \rightarrow \mathcal{X}$ be a smooth morphism, and $f: \mathcal{X} \rightarrow \mathbb{A}^1$ a morphism. Then we have a commutative diagram*

$$\begin{array}{ccc} \hat{M}(\mathcal{X}) & \xrightarrow{g^*} & \hat{M}(\mathcal{Y}) \\ \Psi_f \downarrow & & \downarrow \Psi_{f \circ g} \\ \hat{M}^{\hat{\mu}}(\mathcal{X}_0) & \xrightarrow{g^*} & \hat{M}^{\hat{\mu}}(\mathcal{Y}_0). \end{array} \quad (2.49)$$

In particular, we have $\mathrm{MF}_{f \circ g} = g^(\mathrm{MF}_f)$.*

Proof. For (i), we first restrict to the case when g is representable. By Theorem 2.4.2, the map Ψ_f is determined by pullbacks along smooth morphisms from $\mathbb{K}$ -varieties to $\mathcal{X}$, so we may assume that $\mathcal{X}$ is a $\mathbb{K}$ -variety, and $\mathcal{Y}$ is an algebraic space that is proper over $\mathcal{X}$. This case is covered by Theorem 2.3.5 (i).

For the general case, similarly, we may assume that $\mathcal{X} = X$ is a $\mathbb{K}$ -variety. It suffices to show that $g_! \circ \Psi_{f \circ g}([Z]) = \Psi_f \circ g_!([Z])$ for smooth $\mathbb{K}$ -varieties Z mapping to $\mathcal{Y}$, as these classes span $M(\mathcal{Y})$ over $M(\mathbb{K})$. Since $\mathcal{Y}$ is proper over X and has affine stabilizers, it has finite inertia, and admits a coarse space $\pi_{\mathcal{Y}}: \mathcal{Y} \rightarrow \bar{Y}$ by the Keel–Mori theorem [18; 36]. The morphism $\pi_{\mathcal{Y}}$ is a proper universal homeomorphism.

By Rydh’s compactification theorem for representable morphisms of Deligne–Mumford stacks [48, Theorem B], we may choose a relative compactification $\mathcal{Z}$ of Z over $\mathcal{Y}$, such that there is a dense open immersion $i: Z \hookrightarrow \mathcal{Z}$ and a proper representable morphism $h: \mathcal{Z} \rightarrow \mathcal{Y}$. In particular, $\mathcal{Z}$ also has finite inertia, and admits a coarse space $\pi_{\mathcal{Z}}: \mathcal{Z} \rightarrow \bar{Z}$, which can be seen as a relative compactification of Z over $\bar{Y}$. We have a commutative diagram

$$\begin{array}{ccccc} Z & \xhookrightarrow{i} & \mathcal{Z} & \xrightarrow{\pi_{\mathcal{Z}}} & \bar{Z} \\ & & \downarrow h & & \downarrow \bar{h} \\ & & \mathcal{Y} & \xrightarrow{\pi_{\mathcal{Y}}} & \bar{Y} \\ & & \searrow g & & \swarrow \bar{g} \\ & & & X, & \end{array} \quad (2.50)$$

where $\bar{g}$ and $\bar{h}$ are the induced morphisms, and all morphisms except i are proper. It is then enough to show that

$$(\pi_{\mathcal{Z}})_! \circ \Psi_{f \circ g \circ h}([Z]) = \Psi_{f \circ \bar{g} \circ \bar{h}} \circ (\pi_{\mathcal{Z}})_!([Z]), \quad (2.51)$$

since the compatibility with pushing forward along h and $\bar{g} \circ \bar{h}$ is covered by the previous case.

We now apply Bergh–Rydh’s *divisorialification theorem* [8, Theorem A] to a desingularization of the pair $(\mathcal{Z}, \mathcal{Z} \setminus Z)$ (see, for example, [24]), which gives a representable proper morphism $\tilde{\mathcal{Z}} \rightarrow \mathcal{Z}$ that is an isomorphism on the preimage of Z , such that $\tilde{\mathcal{Z}} \setminus Z = \mathcal{D}$ is a simple normal crossings divisor on $\tilde{\mathcal{Z}}$, with smooth irreducible components $\mathcal{D}_i \subset \tilde{\mathcal{Z}}$, and for each $x \in \tilde{\mathcal{Z}}$, writing $I_x = \{i \in I \mid x \in \mathcal{D}_i\}$, étale locally around x , one has $\tilde{\mathcal{Z}} \sim \prod_{i \in I_x} [\mathbb{A}^1 / \mu_{n_i}] \times \mathbb{A}^{d-|I_x|}$, where $d = \dim \tilde{\mathcal{Z}}$, each μ_{n_i} acts on $\mathbb{A}^1$ by scaling, and $\mathcal{D}_i$ corresponds to the locus where the i -th factor is zero; the number n_i is the order of the generic stabilizer of $\mathcal{D}_i$.

From now on, we assume that $\mathcal{Z} = \tilde{\mathcal{Z}}$, since again, pushing forward along the representable morphism $\tilde{\mathcal{Z}} \rightarrow \mathcal{Z}$ and the corresponding morphism of coarse spaces is already dealt with.

Now, choose a resolution $\pi: \tilde{\mathcal{Z}} \rightarrow \mathcal{Z}$ for the morphism $\mathcal{Z} \rightarrow \mathbb{A}^1$, which is a composition of blow-ups along smooth centres. Then $\tilde{\mathcal{Z}}$ still has the same local description as before. The local description implies that the coarse space of $\tilde{\mathcal{Z}}$, denoted $\bar{Z}$, is

a smooth algebraic space, and can be seen as a resolution for the morphism $\bar{Z} \rightarrow \mathbb{A}^1$.

For each $i \in I$, let $\tilde{\mathcal{D}}_i \subset \tilde{Z}$ be the strict transform of $\mathcal{D}_i$, which is a smooth divisor, and let $(\mathcal{E}_j \subset \tilde{Z})_{j \in J}$ be the family of irreducible components of $\tilde{Z}_0$. Then by construction, all the divisors $\tilde{\mathcal{D}}_i, \mathcal{E}_j \subset \tilde{Z}$ have simple normal crossings, and $\tilde{Z} \setminus \bigcup_{i \in I} \tilde{\mathcal{D}}_i$ is an algebraic space. Let $\tilde{D}_i, E_j \subset \tilde{Z}$ be the corresponding divisors in the coarse spaces. For $I' \subset I$, write $\mathcal{D}_{I'} = \bigcap_{i \in I'} \mathcal{D}_i$ and $\tilde{\mathcal{D}}_{I'} = \bigcap_{i \in I'} \tilde{\mathcal{D}}_i$, with the convention that $\mathcal{D}_\emptyset = \tilde{Z}$ and $\tilde{\mathcal{D}}_\emptyset = \tilde{Z}$. Then, each $\mathcal{D}_{I'}$ can be seen as a resolution for the morphism $\mathcal{D}_{I'} \rightarrow \mathbb{A}^1$. By §2.4.3, we have

$$\begin{aligned} (\pi_Z)_! \circ \Psi_{f \circ g \circ h}([Z]) &= \sum_{I' \subset I} (-1)^{|I'|} \cdot (\pi_Z)_! \circ \Psi_{f \circ g \circ h}([\mathcal{D}_{I'}]) \\ &= \sum_{I' \subset I} (-1)^{|I'|} \cdot \sum_{\emptyset \neq J' \subset J} (1 - \mathbb{L})^{|J'| - 1} [\tilde{\mathcal{E}}_{J'}^\circ \cap \tilde{\mathcal{D}}_{I'}]^\sharp \\ &= \sum_{\emptyset \neq J' \subset J} (1 - \mathbb{L})^{|J'| - 1} [\tilde{\mathcal{E}}_{J'}^\circ \setminus \bigcup_{i \in I} \tilde{\mathcal{D}}_i]^\sharp \\ &= \sum_{\emptyset \neq J' \subset J} (1 - \mathbb{L})^{|J'| - 1} [\tilde{E}_{J'}^\circ \setminus \bigcup_{i \in I} \tilde{D}_i]^\sharp \\ &= \Psi_{f \circ \bar{g} \circ \bar{h}} \circ (\pi_Z)_!([Z]), \end{aligned}$$

where the second last step used the fact that each $\tilde{\mathcal{E}}_{J'}^\circ \setminus \bigcup_{i \in I} \tilde{\mathcal{D}}_i$ is an algebraic space.

For (ii), similarly, the case when g is representable follows from Theorem 2.3.5 (ii). For the general case, we may assume that $\mathcal{X}$ is a $\mathbb{K}$ -variety. Since $\Psi_{f \circ g}$ is determined by pullbacks along smooth morphisms from schemes to $\mathcal{Y}$, we can also assume that $\mathcal{Y}$ is a $\mathbb{K}$ -variety, and the result follows from Theorem 2.3.5 (ii).

The final statement follows from applying (ii) to the element $[\mathcal{X}] \in \hat{M}(\mathcal{X})$. $\square$

2.5 The motivic Behrend function

2.5.1. Shifted symplectic and d-critical stacks. Let $\mathfrak{X}$ be a (-1) -shifted symplectic stack over $\mathbb{K}$ (see §1.1.8), and let $\mathcal{X} = t_0(\mathfrak{X})$ be its classical truncation. Assume that $\mathcal{X}$ is an algebraic stack over $\mathbb{K}$.

Ben-Bassat–Brav–Bussi–Joyce [7, §3.3] define a *d-critical structure* on $\mathcal{X}$ induced from the shifted symplectic structure on $\mathfrak{X}$, so that $\mathcal{X}$ is a *d-critical stack*. See there and Joyce [31] for the precise definitions. For our purposes, it suffices to know the following properties:

- (i) For a smooth $\mathbb{K}$ -variety U and a morphism $f: U \rightarrow \mathbb{A}^1$, the critical locus $\text{Crit}(f) \subset U$ carries a canonical d-critical structure.
- (ii) d-critical structures can be pulled back along smooth morphisms of algebraic stacks over $\mathbb{K}$.
- (iii) If a $\mathbb{K}$ -scheme X carries a d-critical structure, then it can be covered by open subschemes called *critical charts*, each of which with the induced d-critical structure has the form $\text{Crit}(f)$ as in (i). We denote such a critical chart by $i: \text{Crit}(f) \hookrightarrow X$.

2.5.2. Orientations. Let $\mathfrak{X}$ be an s -shifted symplectic stack over $\mathbb{K}$, where s is odd. An *orientation* of $\mathfrak{X}$ is a line bundle $K_{\mathfrak{X}}^{1/2} \rightarrow \mathfrak{X}$, together with an isomorphism $(K_{\mathfrak{X}}^{1/2})^{\otimes 2} \simeq K_{\mathfrak{X}}$, where $K_{\mathfrak{X}}$ is the *canonical bundle* of $\mathfrak{X}$, defined as the determinant line bundle of the cotangent complex of $\mathfrak{X}$.

Note the unfortunate clash of terminology with the unrelated notion of *orientation* in the sense of Pantev–Toën–Vaquié–Vezzosi [47, Definition 2.4]. The latter notion will not be used in this paper except in the proof of Theorem 3.1.5.

Now let $s = -1$, and let $\mathcal{X}$ be the associated d-critical stack of $\mathfrak{X}$, as in §2.5.1. Ben-Bassat–Brav–Bussi–Joyce [7, Theorem 3.18] show that the restriction $K_{\mathfrak{X}}|_{\mathcal{X}^{\text{red}}}$ is determined by the d-critical structure on $\mathcal{X}$, where $\mathcal{X}^{\text{red}}$ is the reduction of $\mathcal{X}$. We denote this restriction simply by $K_{\mathcal{X}}$, and call it the *canonical bundle* of the d-critical stack $\mathcal{X}$. Similarly, we define an *orientation* of a d-critical stack $\mathcal{X}$ to be a line bundle $K_{\mathcal{X}}^{1/2} \rightarrow \mathcal{X}^{\text{red}}$, together with an isomorphism $(K_{\mathcal{X}}^{1/2})^{\otimes 2} \simeq K_{\mathcal{X}}$.

The d-critical scheme $\text{Crit}(f)$ in §2.5.1 (i) has a canonical orientation given by $K_{\text{Crit}(f)}^{1/2} = K_U|_{\text{Crit}(f)^{\text{red}}}$.

As in [31, Lemma 2.58], for a smooth morphism $g: \mathcal{Y} \rightarrow \mathcal{X}$ of d-critical stacks, compatible with the d-critical structures, an orientation $K_{\mathcal{X}}^{1/2}$ of $\mathcal{X}$ induces an orientation of $\mathcal{Y}$ given by $K_{\mathcal{Y}}^{1/2} = g^*(K_{\mathcal{X}}^{1/2}) \otimes \det \mathbb{L}_{\mathcal{Y}/\mathcal{X}}|_{\mathcal{Y}^{\text{red}}}$.

2.5.3. For schemes. Let X be an oriented d-critical $\mathbb{K}$ -scheme. Its *motivic Behrend function* $v_X^{\text{mot}} \in M^{\text{fl}}(X)$ is defined by the following property:

- For any critical chart $i: \text{Crit}(f) \hookrightarrow X$, where $f: U \rightarrow \mathbb{A}^1$ and U is a smooth $\mathbb{K}$ -variety, we have

$$i^*(v_X^{\text{mot}}) = \mathbb{L}^{-\dim U/2} \cdot \Phi_f([U]) \cdot \Upsilon(i^*(K_X^{1/2}) \otimes K_U^{-1}|_{\text{Crit}(f)^{\text{red}}}), \quad (2.52)$$

in $M^{\text{fl}}(\text{Crit}(f))$, where Φ_f is the vanishing cycle map defined in §2.3.3, and $\Phi_f([U])$ is supported on $\text{Crit}(f)$. The map Υ is as in §2.1.11, and the part inside $\Upsilon(\dots)$ is a line bundle on $\text{Crit}(f)^{\text{red}}$ whose square is trivial, so it can be seen as a μ_2 -bundle.

This is well-defined due to Bussi–Joyce–Meinhardt [14, Theorem 5.10].

For X as above, and a smooth morphism $g: Y \rightarrow X$ of relative dimension d , where Y is equipped with the induced oriented d-critical structure, we have the relation

$$g^*(v_X^{\text{mot}}) = \mathbb{L}^{d/2} \cdot v_Y^{\text{mot}}, \quad (2.53)$$

which follows from [7, Theorem 5.14].

2.5.4. For stacks. Let $\mathcal{X}$ be an oriented d-critical stack over $\mathbb{K}$, and assume that $\mathcal{X}$ is Nisnevich locally a quotient stack in the sense of §2.2.4.

We define the *motivic Behrend function* of $\mathcal{X}$, slightly generalizing the construction of Ben-Bassat–Brav–Bussi–Joyce [7, Theorem 5.14], who only considered stacks that are *Zariski locally a quotient stack*.

Theorem. *Let $\mathcal{X}$ be as above. Then there exists a unique element*

$$v_{\mathcal{X}}^{\text{mot}} \in \hat{M}^{\hat{\mu}}(\mathcal{X}), \quad (2.54)$$

called the motivic Behrend function of $\mathcal{X}$, such that for any $\mathbb{K}$ -variety Y and any smooth morphism $f: Y \rightarrow \mathcal{X}$ of relative dimension d , we have

$$f^*(v_{\mathcal{X}}^{\text{mot}}) = \mathbb{L}^{d/2} \cdot v_Y^{\text{mot}} \quad (2.55)$$

in $\hat{M}^{\hat{\mu}}(Y)$, where Y is equipped with the induced oriented d -critical structure.

In particular, we write $v_{\mathfrak{X}}^{\text{mot}} = v_{\mathcal{X}}^{\text{mot}}$ if the d -critical structure on $\mathcal{X}$ comes from a (-1) -shifted symplectic stack $\mathfrak{X}$ with $\mathcal{X} \simeq \mathfrak{t}_0(\mathfrak{X})$.

Proof. We first show that the theorem holds when $\mathcal{X} = X$ is an algebraic space. Indeed, this follows formally from [Theorem 2.2.3](#) and the relation [\(2.53\)](#) for schemes, since X has a Nisnevich cover by affine varieties.

Also, note that if the element $v_{\mathcal{X}}^{\text{mot}}$ exists, then the relation [\(2.55\)](#) must also hold for smooth morphisms from algebraic spaces Y to $\mathcal{X}$, by passing to a Nisnevich cover of Y by affine varieties.

Now, the proof of [\[7, Theorem 5.14\]](#) can be repeated word-by-word to show that the theorem is true when $\mathcal{X} \simeq [S/G]$ is a quotient stack, where S is an algebraic space over $\mathbb{K}$ and $G = \text{GL}(n)$ for some n .

For the general case, let $(j_i: \mathcal{X}_i \hookrightarrow \mathcal{X})_{i \in I}$ be a Nisnevich cover by quotient stacks. The condition on $v_{\mathcal{X}}^{\text{mot}}$ forces that $j_i^*(v_{\mathcal{X}}^{\text{mot}}) = v_{\mathcal{X}_i}^{\text{mot}}$ for all i . We show that the elements $v_{\mathcal{X}_i}^{\text{mot}}$ agree on overlaps. Indeed, let $1, 2 \in I$ be two indices, and let $\mathcal{X}_{1,2} = \mathcal{X}_1 \times_{\mathcal{X}} \mathcal{X}_2$. Then $\mathcal{X}_{1,2}$ is also a quotient stack, so the theorem holds for $\mathcal{X}_{1,2}$. Let $j'_i: \mathcal{X}_{1,2} \rightarrow \mathcal{X}_i$ be the projections, where $i = 1, 2$. Then we have $(j'_i)^*(v_{\mathcal{X}_{1,2}}^{\text{mot}}) = v_{\mathcal{X}_i}^{\text{mot}}$ for $i = 1, 2$, since the left-hand side satisfies the characterizing property of $v_{\mathcal{X}_{1,2}}^{\text{mot}}$. By [Theorem 2.2.3](#), it then follows that the elements $v_{\mathcal{X}_i}^{\text{mot}}$ for $i \in I$ glue to a unique element $v_{\mathcal{X}}^{\text{mot}}$, and a standard argument verifies that it satisfies the relation [\(2.55\)](#). $\square$

2.5.5. Compatibility with smooth pullbacks. We now show that the smooth pullback relation [\(2.55\)](#) holds for all smooth morphisms of d -critical stacks.

Theorem. *Let $\mathcal{X}, \mathcal{Y}$ be oriented d -critical stacks over $\mathbb{K}$ that are Nisnevich locally quotient stacks, and let $f: \mathcal{Y} \rightarrow \mathcal{X}$ be a smooth morphism of relative dimension d which is compatible with the oriented d -critical structures. Then we have the relation*

$$f^*(v_{\mathcal{X}}^{\text{mot}}) = \mathbb{L}^{d/2} \cdot v_{\mathcal{Y}}^{\text{mot}}. \quad (2.56)$$

Proof. It is straightforward to verify that the element $\mathbb{L}^{-d/2} \cdot f^*(v_{\mathcal{X}}^{\text{mot}})$ satisfies the characterizing property of $v_{\mathcal{Y}}^{\text{mot}}$. $\square$

2.5.6. The numerical Behrend function. Let $\mathcal{X}$ be an algebraic stack over $\mathbb{K}$ that is Nisnevich locally a quotient stack, equipped with an oriented d -critical structure. The

Behrend function of $\mathcal{X}$ is the constructible function

$$v_{\mathcal{X}} = \chi(\nu_{\mathcal{X}}^{\text{mot}}) \in \text{CF}(\mathcal{X}) , \quad (2.57)$$

where χ denotes taking the pointwise Euler characteristic, as in §2.1.9.

In fact, one can still define $v_{\mathcal{X}}$ even if $\mathcal{X}$ is only étale locally a quotient stack, and without the orientability assumption. Indeed, the relation (2.56) implies that the Behrend function is compatible with smooth morphisms preserving the d -critical structure (not necessarily orientations), up to a sign $(-1)^d$, where d is the relative dimension. This is because changing the orientation only affects the term $\Upsilon(\cdots)$ in (2.52), which always has Euler characteristic 1. Now, to define $v_{\mathcal{X}}$, one can pass to a smooth cover of $\mathcal{X}$ by $\mathbb{K}$ -varieties, and apply smooth descent of constructible functions.

When $\mathbb{K} = \mathbb{C}$, the Behrend function $v_{\mathcal{X}}$ agrees with the original definitions by Behrend [6] and Joyce–Song [35, §4.1]. This follows from the compatibility of both versions with smooth pullbacks, Theorem 2.5.5 and [35, Theorem 4.3], and the case of critical loci on smooth varieties, which follows from [22, Theorem 3.10] and [35, Theorem 4.7].

3 Graded and filtered points

We discuss the *stack of graded points* and the *stack of filtered points* of algebraic stacks and derived algebraic stacks, following Halpern-Leistner [26], and study their interactions with shifted symplectic structures. Then, in §3.2, we use the theory of étale local structures of algebraic stacks developed by Alper–Hall–Rydth [2] to give local descriptions of these stacks.

3.1 Definition and deformation theory

3.1.1. Definition. Let $\mathfrak{X}$ be a derived algebraic stack over $\mathbb{K}$, such that its classical truncation $\mathcal{X} = t_0(\mathfrak{X})$ is a classical algebraic stack over $\mathbb{K}$.

Following Halpern-Leistner [26], define the *stack of graded points* and the *stack of filtered points* of $\mathfrak{X}$, respectively, to be the derived mapping stacks over $\mathbb{K}$,

$$\mathrm{Grad}(\mathfrak{X}) = \mathfrak{Map}([*/\mathbb{G}_m], \mathfrak{X}), \quad (3.1)$$

$$\mathrm{Filt}(\mathfrak{X}) = \mathfrak{Map}([\mathbb{A}^1/\mathbb{G}_m], \mathfrak{X}), \quad (3.2)$$

where $\mathbb{G}_m$ acts on $\mathbb{A}^1$ by scaling. The morphisms

$$(-)^{-1} \curvearrowright [*/\mathbb{G}_m] \xrightleftharpoons[\mathrm{pr}]{0} [\mathbb{A}^1/\mathbb{G}_m] \xrightleftharpoons[1]{0} *, \quad (3.3)$$

where pr is induced by the projection $\mathbb{A}^1 \rightarrow *$, induce morphisms of stacks

$$\mathrm{op} \curvearrowright \mathrm{Grad}(\mathfrak{X}) \xrightleftharpoons[\mathrm{fi}]{\mathrm{gr}} \mathrm{Filt}(\mathfrak{X}) \xrightleftharpoons[\mathrm{ev}_1]{\mathrm{ev}_0} \mathfrak{X}, \quad (3.4)$$

where the notations ‘op’, ‘gr’, ‘fi’, and ‘tot’ stand for the *opposite graded point*, the *associated graded point*, the *split filtration*, and the *total point*, respectively. We will use the morphisms tot and ev_1 to regard $\mathrm{Grad}(\mathfrak{X})$ and $\mathrm{Filt}(\mathfrak{X})$ as stacks over $\mathfrak{X}$.

By Halpern-Leistner [26, Proposition 1.1.2 and Lemma 1.3.8], if $\mathfrak{X}$ is a (classical) algebraic stack over $\mathbb{K}$ with affine stabilizers, then so are $\mathrm{Grad}(\mathfrak{X})$ and $\mathrm{Filt}(\mathfrak{X})$. In this case, the morphism gr is of finite type, and induces a bijection $\pi_0(\mathrm{Filt}(\mathfrak{X})) \xrightarrow{\sim} \pi_0(\mathrm{Grad}(\mathfrak{X}))$.

It is also shown in [26, Lemma 1.2.1] that taking graded or filtered points commutes with taking the classical truncation of a derived algebraic stack.

3.1.2. For quotient stacks. Let $\mathcal{X} = [U/G]$ be a quotient stack, where U is an algebraic space over $\mathbb{K}$, acted on by a reductive group G . The stacks of graded and filtered points of $\mathcal{X}$ can be described very explicitly, following Halpern-Leistner [26, §1.4].

Let $\lambda: \mathbb{G}_m \rightarrow G$ be a *cocharacter*, that is, a morphism of algebraic groups. Define

the *Levi subgroup* and the *parabolic subgroup* of G associated to λ by

$$L_\lambda = \{g \in G \mid g = \lambda(t) g \lambda(t)^{-1} \text{ for all } t\}, \quad (3.5)$$

$$P_\lambda = \{g \in G \mid \lim_{t \rightarrow 0} \lambda(t) g \lambda(t)^{-1} \text{ exists}\}, \quad (3.6)$$

respectively. Define the *fixed locus* and the *attractor* associated to λ by

$$U^{\lambda,0} = \text{Map}^{\mathbb{G}_m}(*, U), \quad (3.7)$$

$$U^{\lambda,+} = \text{Map}^{\mathbb{G}_m}(\mathbb{A}^1, U), \quad (3.8)$$

where $\text{Map}^{\mathbb{G}_m}(-, -)$ denotes the $\mathbb{G}_m$ -equivariant mapping space; $\mathbb{G}_m$ acts on U via λ , and on $\mathbb{A}^1$ by scaling. These are algebraic spaces by Drinfeld–Gaitsgory [23, Proposition 1.3.4 and Theorem 1.5.2] or Halpern-Leistner [26, Proposition 1.4.1]. There is a closed immersion $U^{\lambda,0} \hookrightarrow U$, an unramified morphism $U^{\lambda,+} \rightarrow U$ given by evaluation at 1, and an affine morphism $U^{\lambda,+} \rightarrow U^{\lambda,0}$ given by evaluation at 0.

The G -action on U induce a P_λ -action on $U^{\lambda,+}$ and an L_λ -action on $U^{\lambda,0}$. Moreover, by [26, Theorem 1.4.8], we have

$$\text{Grad}(\mathcal{X}) \simeq \bigsqcup_{\lambda: \mathbb{G}_m \rightarrow G} [U^{\lambda,0}/L_\lambda], \quad (3.9)$$

$$\text{Filt}(\mathcal{X}) \simeq \bigsqcup_{\lambda: \mathbb{G}_m \rightarrow G} [U^{\lambda,+}/P_\lambda], \quad (3.10)$$

where the disjoint union is over all conjugacy classes of cocharacters λ . The morphism $\text{tot}: \text{Grad}(\mathcal{X}) \rightarrow \mathcal{X}$ is induced by the $(L_\lambda \hookrightarrow G)$ -equivariant morphism $U^{\lambda,0} \rightarrow U$; $\text{ev}_1: \text{Filt}(\mathcal{X}) \rightarrow \mathcal{X}$ induced by the $(P_\lambda \hookrightarrow G)$ -equivariant morphism $U^{\lambda,+} \rightarrow U$; and $\text{gr}: \text{Filt}(\mathcal{X}) \rightarrow \text{Grad}(\mathcal{X})$ induced by the $(P_\lambda \rightarrow L_\lambda)$ -equivariant morphism $U^{\lambda,+} \rightarrow U^{\lambda,0}$.

3.1.3. Deformation theory. One can express the tangent complexes of $\text{Grad}(\mathcal{X})$ and $\text{Filt}(\mathcal{X})$ in terms of that of $\mathcal{X}$. Concretely, as in Halpern-Leistner [26, Lemma 1.2.2], we have

$$\mathbb{T}_{\text{Grad}(\mathcal{X})} \simeq \text{tot}^*(\mathbb{T}_{\mathcal{X}})_0, \quad (3.11)$$

$$\mathbb{T}_{\text{Filt}(\mathcal{X})} \simeq q_* \circ \text{ev}^*(\mathbb{T}_{\mathcal{X}}), \quad (3.12)$$

where $(-)_0$ denotes taking the weight 0 part with respect to the natural $\mathbb{G}_m$ -action, $\text{ev}: [\mathbb{A}^1/\mathbb{G}_m] \times \text{Filt}(\mathcal{X}) \rightarrow \mathcal{X}$ is the evaluation morphism, and $q: [\mathbb{A}^1/\mathbb{G}_m] \times \text{Filt}(\mathcal{X}) \rightarrow \text{Filt}(\mathcal{X})$ is the projection.

Moreover, for shifted symplectic stacks $\mathcal{X}$, we have a more refined description of the tangent complexes in Theorem 3.1.5 below.

3.1.4. Oriented Lagrangian correspondences. Let $\mathfrak{X}, \mathfrak{Y}$ be oriented s -shifted symplectic stacks over $\mathbb{K}$, as in §2.5.2, where s is odd, and let

$$\mathfrak{X} \xleftarrow{f} \mathfrak{L} \xrightarrow{g} \mathfrak{Y} \quad (3.13)$$

be an s -shifted Lagrangian correspondence, in the sense of [16, §2.4]. We thus have an exact triangle

$$\mathbb{T}_{\mathfrak{L}} \longrightarrow f^*(\mathbb{T}_{\mathfrak{X}}) \oplus g^*(\mathbb{T}_{\mathfrak{Y}}) \longrightarrow \mathbb{L}_{\mathfrak{L}}[s] \longrightarrow \mathbb{T}_{\mathfrak{L}}[1] \quad (3.14)$$

of perfect complexes on $\mathfrak{L}$. An orientation of the shifted Lagrangian correspondence (3.13) is an isomorphism $K_{\mathfrak{L}} \simeq f^*(K_{\mathfrak{X}}^{1/2}) \otimes g^*(K_{\mathfrak{Y}}^{1/2})$, such that it squares to the canonical isomorphism $K_{\mathfrak{L}}^{\otimes 2} \simeq f^*(K_{\mathfrak{X}}) \otimes g^*(K_{\mathfrak{Y}})$ induced by the exact triangle (3.14).

3.1.5. Theorem. *Let $\mathfrak{X}$ be an s -shifted symplectic stack over $\mathbb{K}$, with symplectic structure ω . Then we have an induced s -shifted symplectic structure $\text{tot}^*(\omega)$ on $\text{Grad}(\mathfrak{X})$, and an s -shifted Lagrangian correspondence*

$$\text{Grad}(\mathfrak{X}) \xleftarrow{\text{gr}} \text{Filt}(\mathfrak{X}) \xrightarrow{\text{ev}_1} \mathfrak{X}. \quad (3.15)$$

Moreover, if s is odd and $\mathfrak{X}$ has an orientation $K_{\mathfrak{X}}^{1/2}$, then $\text{Grad}(\mathfrak{X})$ has an induced orientation $K_{\text{Grad}(\mathfrak{X})}^{1/2}$, and the Lagrangian correspondence is oriented.

Proof. To prove that (3.15) is an s -shifted Lagrangian correspondence, by Calaque [15, Theorem 4.8], it is enough to show that the cospan

$$[*/\mathbb{G}_m] \xrightarrow{0} [\mathbb{A}^1/\mathbb{G}_m] \xleftarrow{1} * \quad (3.16)$$

is a 0-oriented cospan, in the sense of [15, §4.2] and [16, §2.5]. Indeed, $*$ carries a natural 0-orientation, and the 0-orientation on $[\mathbb{A}^1/\mathbb{G}_m]$ is given by the evaluation map $\mathbb{R}\Gamma(\mathcal{O}_{[\mathbb{A}^1/\mathbb{G}_m]}) \rightarrow \mathbb{K}$. To see that this is indeed a 0-orientation, we check the condition in [47, Definition 2.4]. For $A \in \text{CdgA}_{\mathbb{K}}^{\leq 0}$ and a perfect complex $\mathcal{E} \in \text{Perf}(\text{Spec } A \times [*/\mathbb{G}_m])$, one has $p_*(\mathcal{E}^\vee)^\vee \simeq p_*(\mathcal{E})$ on $\text{Spec } A$, where $p: \text{Spec } A \times [*/\mathbb{G}_m] \rightarrow \text{Spec } A$ is the projection, since both sides are the weight 0 part of the induced $\mathbb{G}_m$ -action on $\pi^*(\mathcal{E})$, where $\pi: \text{Spec } A \rightarrow \text{Spec } A \times [*/\mathbb{G}_m]$ is the projection.

To see that (3.16) is a 0-oriented cospan, we check the condition in [16, Lemma 2.5.5]. For any $A \in \text{CdgA}_{\mathbb{K}}^{\leq 0}$ and $\mathcal{E} \in \text{Perf}(\text{Spec } A \times [\mathbb{A}^1/\mathbb{G}_m])$, we need to show that the induced 2-commutative diagram

$$\begin{array}{ccc} q_*(\mathcal{E}) & \longrightarrow & p_* \circ 0^*(\mathcal{E}) \\ \downarrow & & \downarrow \\ 1^*(\mathcal{E}) & \longrightarrow & q_*(\mathcal{E}^\vee)^\vee \end{array} \quad (3.17)$$

in $\text{Perf}(A)$ is cartesian, where p and q are the projections from $\text{Spec } A \times [*/\mathbb{G}_m]$ and $\text{Spec } A \times [\mathbb{A}^1/\mathbb{G}_m]$ to $\text{Spec } A$, respectively. Indeed, as in Halpern-Leistner [27, Propo-

ition 1.1.2 ff.], such an object $\mathcal{E}$ can be seen as a filtered object in $\text{Perf}(A)$, that is, a sequence of maps

$$\cdots \longrightarrow E_{\geq 1} \longrightarrow E_{\geq 0} \longrightarrow E_{\geq -1} \longrightarrow \cdots \quad (3.18)$$

in $\text{Perf}(A)$, where all but finitely many arrows are isomorphisms, such that $E_{\geq n} = 0$ for $n \gg 0$. Write $E_n = \text{cofib}(E_{\geq n+1} \rightarrow E_{\geq n})$, and write $E = \text{colim}_{n \rightarrow -\infty} E_{\geq n}$. Then $0^*(\mathcal{E}) \simeq \bigoplus_n E_n$, with the natural $\mathbb{G}_m$ -action having weight n on E_n . One can deduce from [27, Proposition 1.1.2 ff.] that we have natural identifications

$$q_*(\mathcal{E}) \simeq E_{\geq 0}, \quad (3.19)$$

$$p_* \circ 0^*(\mathcal{E}) \simeq E_0, \quad (3.20)$$

$$1^*(\mathcal{E}) \simeq E, \quad (3.21)$$

$$q_*(\mathcal{E}^\vee)^\vee \simeq ((E^\vee)_{\geq 0})^\vee \simeq E_{\leq 0}, \quad (3.22)$$

where $E_{\leq 0} = \text{cofib}(E_{\geq 1} \rightarrow E)$, and the arrows in the diagram (3.17) are the natural ones. This implies that (3.17) is cartesian.

For the final statement, observe that

$$\begin{aligned} \text{tot}^*(K_{\mathfrak{X}}) &\simeq \det(\text{tot}^*(\mathbb{L}_{\mathfrak{X}})^0) \otimes \det(\text{tot}^*(\mathbb{L}_{\mathfrak{X}})^+) \otimes \det(\text{tot}^*(\mathbb{L}_{\mathfrak{X}})^-) \\ &\simeq K_{\text{Grad}(\mathfrak{X})} \otimes \det(\text{tot}^*(\mathbb{L}_{\mathfrak{X}})^+) \otimes \det((\text{tot}^*(\mathbb{L}_{\mathfrak{X}})^+)^{\vee}[-s]) \\ &\simeq K_{\text{Grad}(\mathfrak{X})} \otimes \det(\text{tot}^*(\mathbb{L}_{\mathfrak{X}})^+)^2, \end{aligned} \quad (3.23)$$

where $(-)^0$, $(-)^+$, $(-)^-$ denote the parts with zero, positive, and negative weights, respectively, with respect to the natural $\mathbb{G}_m$ -action. Therefore, we may define

$$K_{\text{Grad}(\mathfrak{X})}^{1/2} = \text{tot}^*(K_{\mathfrak{X}}^{1/2}) \otimes \det(\text{tot}^*(\mathbb{L}_{\mathfrak{X}})^+)^{-1}, \quad (3.24)$$

and this gives an orientation on $\text{Grad}(\mathfrak{X})$. To see that the s -shifted Lagrangian correspondence is oriented, consider the 2-cartesian diagram

$$\begin{array}{ccc} \mathbb{T}_{\text{Filt}(\mathfrak{X})} & \longrightarrow & \text{gr}^*(\mathbb{T}_{\text{Grad}(\mathfrak{X})}) \\ \downarrow & \ulcorner & \downarrow \\ \text{ev}_1^*(\mathbb{T}_{\mathfrak{X}}) & \longrightarrow & \mathbb{L}_{\text{Filt}(\mathfrak{X})}[s] \end{array} \quad (3.25)$$

in $\text{Perf}(\text{Filt}(\mathfrak{X}))$, witnessing the s -shifted Lagrangian correspondence structure. Write $\mathcal{E} = \text{ev}^*(\mathbb{T}_{\mathfrak{X}})$, where $\text{ev}: [\mathbb{A}^1/\mathbb{G}_m] \times \text{Filt}(\mathfrak{X}) \rightarrow \mathfrak{X}$ is the evaluation morphism. As in the argument above, $\mathcal{E}$ can be seen as a filtered object in $\text{Perf}(\text{Filt}(\mathfrak{X}))$, and the terms in (3.25) can be identified with $E_{\geq 0}$, E_0 , E , and $E_{\leq 0}$, respectively. In particular, one has $K_{\text{Filt}(\mathfrak{X})} \simeq \text{gr}^*(K_{\text{Grad}(\mathfrak{X})}^{1/2}) \otimes \text{ev}_1^*(K_{\mathfrak{X}}^{1/2})$, as both sides can be identified with $\det(E_{\geq 0})^{-1}$. $\square$

3.1.6. Lemma. *Let $\mathfrak{X}$ be an s -shifted symplectic stack over $\mathbb{K}$. Then we have*

$$\text{fi}^*(\mathbb{T}_{\text{Filt}(\mathfrak{X})}) \simeq \text{op}^* \circ \text{fi}^*(\mathbb{L}_{\text{Filt}(\mathfrak{X})}[s]) \quad (3.26)$$

on $\text{Grad}(\mathcal{X})$.

Proof. By Halpern-Leistner [26, Lemma 1.2.3], we have $\text{fi}^*(\mathbb{T}_{\text{Filt}(\mathcal{X})}) \simeq \text{tot}^*(\mathbb{T}_{\mathcal{X}})_{\geq 0}$, where $(-)_{\geq 0}$ denotes taking the part with non-negative weights with respect to the natural $\mathbb{G}_m$ -action. Consequently, we have $\text{op}^* \circ \text{fi}^*(\mathbb{T}_{\text{Filt}(\mathcal{X})}) \simeq \text{tot}^*(\mathbb{T}_{\mathcal{X}})_{\leq 0}$; its dual shifted by s becomes $\text{tot}^*(\mathbb{L}_{\mathcal{X}}[s])_{\geq 0} \simeq \text{tot}^*(\mathbb{T}_{\mathcal{X}})_{\geq 0}$. $\square$

3.2 Local structure

3.2.1. In this subsection, we consider the stacks of graded and filtered points of algebraic stacks admitting étale covers by quotient stacks in the sense of §2.2.4–2.2.5, and give local descriptions of these stacks.

3.2.2. Theorem. *Let $\mathcal{X}$ be an algebraic stack over $\mathbb{K}$, and let $(\mathcal{X}_i \rightarrow \mathcal{X})_{i \in I}$ be a representable étale cover, where each $\mathcal{X}_i \simeq [S_i/G_i]$, with S_i an algebraic space over $\mathbb{K}$ and G_i a reductive group. Then there are commutative diagrams*

$$\begin{array}{ccccc} [S_i^{\lambda,0}/L_{i,\lambda}] & \longleftarrow & [S_i^{\lambda,+}/P_{i,\lambda}] & \longrightarrow & [S_i/G_i] \\ \downarrow & & \downarrow & & \downarrow \\ \text{Grad}(\mathcal{X}) & \xleftarrow{\text{gr}} & \text{Filt}(\mathcal{X}) & \xrightarrow{\text{ev}_1} & \mathcal{X}, \end{array} \quad (3.27)$$

where all vertical arrows are representable and étale, $\lambda: \mathbb{G}_m \rightarrow G_i$ is a cocharacter, and the left-hand square is a pullback square. Moreover, the families

$$([S_i^{\lambda,0}/L_{i,\lambda}] \rightarrow \text{Grad}(\mathcal{X}))_{i \in I, \lambda: \mathbb{G}_m \rightarrow G_i}, \quad (3.28)$$

$$([S_i^{\lambda,+}/P_{i,\lambda}] \rightarrow \text{Filt}(\mathcal{X}))_{i \in I, \lambda: \mathbb{G}_m \rightarrow G_i} \quad (3.29)$$

are representable étale covers of $\text{Grad}(\mathcal{X})$ and $\text{Filt}(\mathcal{X})$, respectively.

Proof. By Halpern-Leistner [26, Corollary 1.1.7], one has $\text{Grad}(\mathcal{X}_i) \simeq \text{Grad}(\mathcal{X}) \times_{\mathcal{X}} \mathcal{X}_i$ for all i . Therefore, the family $(\text{Grad}(\mathcal{X}_i) \rightarrow \text{Grad}(\mathcal{X}))_{i \in I}$ is a representable étale cover. By Lemma 3.2.4 below, so is the family $(\text{Filt}(\mathcal{X}_i) \rightarrow \text{Filt}(\mathcal{X}))_{i \in I}$. The rest of the theorem follows from the description of $\text{Grad}(\mathcal{X}_i)$ and $\text{Filt}(\mathcal{X}_i)$ in §3.1.2. That the left-hand square in (3.27) is a pullback square follows from Lemma 3.2.4 below. $\square$

3.2.3. Lemma. *Let U be an algebraic space over $\mathbb{K}$ acted on by $\mathbb{G}_m$, V a finite-dimensional $\mathbb{G}_m$ -representation with only positive weights, and $f: U \rightarrow V$ a $\mathbb{G}_m$ -equivariant étale morphism. If there exists a point $u_0 \in U(\mathbb{K})$ such that $f(u_0) = 0$ and $\lim_{t \rightarrow 0} f(t \cdot u) = u_0$ for all $u \in U$, then f is an isomorphism.*

Proof. It is enough to show that f is bijective. It is surjective since it is open and $\mathbb{G}_m$ -equivariant. To see that it is injective, suppose that $f(u_1) = f(u_2) = v \neq 0$ for

$u_1, u_2 \in U(\mathbb{K})$. Consider the morphism $g: \mathbb{A}^1 \rightarrow V$ given by $g(t) = t \cdot v$, and form the pullback diagram

$$\begin{array}{ccc} U' & \xrightarrow{g'} & U \\ f' \downarrow & \ulcorner & \downarrow f \\ \mathbb{A}^1 & \xrightarrow{g} & V. \end{array} \quad (3.30)$$

For $i = 1, 2$, define $h_i: \mathbb{A}^1 \rightarrow U'$ by $h_i(t) = (t, t \cdot u_i)$. Then h_1, h_2 are sections of the étale morphism f' , and agree at 0, so they agree on an open neighbourhood of 0, and hence everywhere. This implies that $u_1 = u_2$. $\square$

3.2.4. Lemma. *Let $\mathcal{X}, \mathcal{Y}$ be algebraic stacks over $\mathbb{K}$ with affine stabilizers, and let $f: \mathcal{X} \rightarrow \mathcal{Y}$ be a representable étale morphism. Then there is a 2-pullback diagram*

$$\begin{array}{ccc} \mathrm{Filt}(\mathcal{Y}) & \longrightarrow & \mathrm{Filt}(\mathcal{X}) \\ \mathrm{gr} \downarrow & \ulcorner & \downarrow \mathrm{gr} \\ \mathrm{Grad}(\mathcal{Y}) & \longrightarrow & \mathrm{Grad}(\mathcal{X}). \end{array} \quad (3.31)$$

Proof. Note first that all the stacks in (3.31) are locally of finite type. Let T be an affine $\mathbb{K}$ -scheme of finite type. Then the groupoid of T -points of $\mathrm{Grad}(\mathcal{Y}) \times_{\mathrm{Grad}(\mathcal{X})} \mathrm{Filt}(\mathcal{X})$ can be identified with the groupoid of 2-commutative diagrams

$$\begin{array}{ccc} T \times [*/\mathbb{G}_m] & \longrightarrow & \mathcal{Y} \\ T \times 0 \downarrow & & \downarrow f \\ T \times [\mathbb{A}^1/\mathbb{G}_m] & \longrightarrow & \mathcal{X}, \end{array} \quad (3.32)$$

where the vertical edges are fixed. It is then enough to show that each such diagram has a unique lifting $T \times [\mathbb{A}^1/\mathbb{G}_m] \rightarrow \mathcal{Y}$, up to a unique 2-isomorphism.

Write $\mathcal{U} = (T \times [\mathbb{A}^1/\mathbb{G}_m]) \times_{\mathcal{X}} \mathcal{Y}$, and let $g: \mathcal{U} \rightarrow T \times [\mathbb{A}^1/\mathbb{G}_m]$ be the projection. Then g is representable and étale, and has a section s over the closed substack $T \times [*/\mathbb{G}_m]$. It is enough to show that s can be extended uniquely to a section of g . Indeed, write $U = \mathcal{U} \times_{T \times [\mathbb{A}^1/\mathbb{G}_m]} (T \times \mathbb{A}^1)$, so that $\mathcal{U} \simeq [U/\mathbb{G}_m]$. The induced morphism $\bar{g}: U \rightarrow T \times \mathbb{A}^1$ is a $\mathbb{G}_m$ -equivariant étale morphism of algebraic spaces over T , with a section $\bar{s}$ over $T \times \{0\}$. Let $U^0 \subset U$ be the image of $\bar{s}$, and let $U^+ \subset U$ be the attracted locus of U^0 under the $\mathbb{G}_m$ -action. Then $U^+ \subset U$ is open. We claim that $\bar{g}|_{U^+}$ is an isomorphism onto $T \times \mathbb{A}^1$, which implies the required unique extension property. Indeed, since $\bar{g}|_{U^+}$ is étale, it is enough to show that it is bijective, which can be checked on $\mathbb{K}$ -points of T , and we are reduced to Lemma 3.2.3 with $V = \mathbb{A}^1$. $\square$

4 The main results

We now present the main results of this paper, in three different versions. First, in §4.1, we prove a local version of the main theorem, using the theory of motivic nearby and vanishing cycles for stacks developed in §2.4. Then, in §4.2, we glue the local versions together to prove the global version of our main result, Theorem 4.2.2. Finally, in §4.3, we take Euler characteristics in the main identity, and obtain integral identities involving the numerical Behrend functions.

4.1 The local version

4.1.1. Theorem. *Suppose that we are given the following data:*

- A finite-dimensional $\mathbb{G}_m$ -representation V over $\mathbb{K}$. Let

$$V = \bigoplus_{k \in \mathbb{Z}} V_k \quad (4.1)$$

be the decomposition into weight spaces. Write $V_+ = \bigoplus_{k > 0} V_k$.

- A $\mathbb{K}$ -variety U acted on by $\mathbb{G}_m$, and a $\mathbb{G}_m$ -equivariant étale morphism $\iota: U \rightarrow V$. Write $U^0 = U^{\mathbb{G}_m}$, and for $u_0 \in U^0$, define

$$U^+(u_0) = \{u \in U \mid \lim_{t \rightarrow 0} t \cdot u = u_0\}. \quad (4.2)$$

We have a canonical isomorphism $U^+(u_0) \simeq V_+$ by Lemma 3.2.3.

- A $\mathbb{G}_m$ -invariant function $f: U \rightarrow \mathbb{A}^1$, with $f(0) = 0$.

Then we have the identities

$$\int_{u \in U^+(u_0)} \Psi_f([U])(u) = \mathbb{L}^{\dim V_+} \cdot \Psi_f([U^0])(u_0), \quad (4.3)$$

$$\int_{u \in U^+(u_0)} \Phi_f([U])(u) = \mathbb{L}^{\dim V_+} \cdot \Phi_f([U^0])(u_0). \quad (4.4)$$

Moreover, these hold as identities in $M^{\text{fl}}(U^0)$, where we vary $u_0 \in U^0$.

This theorem can be seen as a generalization of the integral identity conjectured by Kontsevich–Soibelman [40, Conjecture 4], and proved by Lê [44], who restricted to the case when the $\mathbb{G}_m$ -action on V only has weights $-1, 0$, and 1 . Compare also Joyce–Song [35, Theorem 5.11], where a similar identity involving Euler characteristics is proved, with the same restriction on the weights.

The rest of this subsection is devoted to the proof of Theorem 4.1.1. In the following, we first provide preliminaries on weighted projective spaces and weighted

blow-ups, and prove some preparatory results. Then, in [Lemma 4.1.8](#), we establish a weaker version of the theorem, using the theory of motivic nearby cycles for stacks developed in [§2.4](#). Finally, in [§4.1.9](#), we show that the weaker version implies the stronger version.

4.1.2. Weighted projective spaces. Let V be a finite-dimensional $\mathbb{G}_m$ -representation over $\mathbb{K}$, with only positive weights. The *weighted projective space* of V is defined to be the quotient stack

$${}^w\mathbb{P}(V) = [(V \setminus \{0\})/\mathbb{G}_m]. \quad (4.5)$$

This is a proper Deligne–Mumford stack over $\mathbb{K}$, since we have the identification

$${}^w\mathbb{P}(V) \simeq \left[\mathbb{P}(V) \Big/ \prod_{k=1}^{\dim V} \mu_{n_k} \right], \quad (4.6)$$

where $\mathbb{P}(V)$ is the usual projective space, and using a basis of eigenvectors of V , each n_k is the weight of the k -th coordinate, and μ_{n_k} acts by scaling the k -th coordinate.

By [§2.1.7](#), the motive of ${}^w\mathbb{P}(V)$ is given by

$$[{}^w\mathbb{P}(V)] = \frac{\mathbb{L}^{\dim V} - 1}{\mathbb{L} - 1}, \quad (4.7)$$

and is independent of the choice of weights on V .

We also consider the coarse space ${}^{cw}\mathbb{P}(V)$ of ${}^w\mathbb{P}(V)$, which is also given by

$${}^{cw}\mathbb{P}(V) = \text{Proj } \mathbb{K}[V], \quad (4.8)$$

where $\mathbb{K}[V]$ is the free polynomial algebra on V , with $\mathbb{Z}$ -grading given by the weights of V . It is an integral normal projective $\mathbb{K}$ -variety.

4.1.3. Weighted blow-ups. Let V be a finite-dimensional $\mathbb{G}_m$ -representation over $\mathbb{K}$, with only positive weights. Let U be a smooth $\mathbb{K}$ -scheme, $U_0 \subset U$ a reduced closed subscheme, and let $p: U \rightarrow V$ be a smooth morphism such that $U_0 = p^{-1}(0)$.

The *weighted blow-up* of U along U_0 , with weights given by those of V , is defined to be the quotient stack

$${}^w\text{Bl}_{U_0}(U) = \left[\left\{ (t, v, u) \in \mathbb{A}^1 \times (V \setminus \{0\}) \times U \mid p(u) = t \cdot v \right\} / \mathbb{G}_m \right], \quad (4.9)$$

where $t \cdot (-)$ denotes the $\mathbb{G}_m$ -action naturally extended to $t \in \mathbb{A}^1$, and $\mathbb{G}_m$ acts with weight -1 on $\mathbb{A}^1$ and trivially on U .

The natural projection ${}^w\text{Bl}_{U_0}(U) \rightarrow U$ is proper. It restricts to an isomorphism over $U \setminus U_0$, and has fibres ${}^w\mathbb{P}(V)$ over U_0 . In particular, we have the relation

$$[{}^w\text{Bl}_{U_0}(U)] = \frac{\mathbb{L}^{\dim V} - 1}{\mathbb{L} - 1} \cdot [U_0] + [U \setminus U_0] \quad (4.10)$$

of motives on U .

4.1.4. Lemma. *Let U be a separated algebraic space of finite type over $\mathbb{K}$, acted on by a torus $T \simeq \mathbb{G}_m^n$ for some n , such that points in U have finite stabilizers. Let $\mathcal{X} = [U/T]$ be the quotient stack.*

Then $\mathcal{X}$ has finite inertia, and admits a coarse space $\pi: \mathcal{X} \rightarrow X$ which is a proper universal homeomorphism. Moreover, we have an isomorphism

$$\pi_! = (\pi^*)^{-1}: \hat{M}(\mathcal{X}) \xrightarrow{\sim} \hat{M}(X), \quad (4.11)$$

and similarly for $\hat{M}^{\dagger}(-)$.

Proof. The inertia $\mathcal{I}_{\mathcal{X}}$ is a closed substack of $H \times \mathcal{X}$ for some finite group $H \subset T$, and is thus finite over $\mathcal{X}$. It then follows from the Keel–Mori theorem [18] that $\mathcal{X}$ admits a coarse space $\pi: \mathcal{X} \rightarrow X$, and that π is a proper universal homeomorphism.

To prove (4.11), stratify U by locally closed subspaces $U_i \subset U$, where each U_i is the locus of points with stabilizer $H_i \subset T$, and let $\mathcal{X}_i \subset \mathcal{X}$ and $X_i \subset X$ be the corresponding strata. Then $\mathcal{X}_i \simeq (U_i/(T/H_i)) \times [*/H_i]$, so that $X_i \simeq U_i/(T/H_i)$ and $\mathcal{X}_i \rightarrow X_i$ is a trivial gerbe with stabilizer H_i . Note that the motive of $[*/\mu_k]$ is 1 for all $k > 0$, since it is the quotient $[\mathbb{G}_m/\mathbb{G}_m]$ with a weight k action, and $\mathbb{G}_m$ is a special group; see §2.1.7. It follows that the motive of each $[*/H_i]$ is 1, and the result follows. $\square$

4.1.5. Lemma. *In the situation of Theorem 4.1.1, the locus in U where the morphism ι preserves $\mathbb{G}_m$ -stabilizers is open.*

Proof. For each $n > 1$, let $\zeta_n \in \mathbb{G}_m$ be a primitive n -th root of unity. It is enough to show that the locus of $u \in U$ such that $\zeta_n \cdot u \neq u$ and $\iota(\zeta_n \cdot u) = \iota(u)$ is closed. The latter condition is equivalent to $\iota(u) \in V_{(n)}$, where $V_{(n)} = \bigoplus_{k \in \mathbb{Z}} V_{kn} \subset V$. Write $U_{(n)} = \iota^{-1}(V_{(n)})$, which is étale over $V_{(n)}$, with a μ_n -action on its fibres, induced from the $\mathbb{G}_m$ -action on U . The locus where this action is trivial is open in $U_{(n)}$, proving the claim. $\square$

4.1.6. Lemma. *In the situation of Theorem 4.1.1, suppose that U is affine, and ι preserves $\mathbb{G}_m$ -stabilizers and sends closed $\mathbb{G}_m$ -orbits to closed $\mathbb{G}_m$ -orbits. Then the affine GIT quotient $U//\mathbb{G}_m$ is smooth.*

Proof. By Alper [1, Theorem 5.1], since ι is étale and preserves $\mathbb{G}_m$ -stabilizers, the induced morphism $\bar{\iota}: U//\mathbb{G}_m \rightarrow V//\mathbb{G}_m$ is étale at $[u] \in (U//\mathbb{G}_m)(\mathbb{K})$ for points $u \in U(\mathbb{K})$ such that the $\mathbb{G}_m$ -orbits of u and $\iota(u)$ are closed. By the assumption on closed orbits, it is enough to require that the $\mathbb{G}_m$ -orbit of u is closed. Since every S -equivalence class in U contains a closed orbit, the morphism $\bar{\iota}$ is étale, and it is enough to check that $V//\mathbb{G}_m$ is smooth, which is standard. $\square$

4.1.7. Lemma. *Let $f: X \rightarrow Y$ be a morphism of integral $\mathbb{K}$ -varieties. If f is bijective on $\mathbb{K}$ -points and Y is normal, then f is an isomorphism.*

Proof. By generic flatness, f is flat over a dense open subset $U \subset Y$ with fibres $\text{Spec } \mathbb{K}$, and hence étale, hence an isomorphism $f^{-1}(U) \xrightarrow{\sim} U$. It follows that f is birational.

Now, a version of Zariski's main theorem [25, IV-3, Corollary 8.12.10] implies that f is an open immersion, hence an isomorphism. $\square$

4.1.8. Lemma. *In the situation of Theorem 4.1.1, write $V_- = \bigoplus_{k < 0} V_k$, and for $u_0 \in U^0$, define*

$$U^-(u_0) = \{u \in U \mid \lim_{t \rightarrow \infty} t \cdot u = u_0\}. \quad (4.12)$$

Then we have the identity

$$\int_{u \in U^+(u_0)} \Psi_f([U])(u) - \int_{u \in U^-(u_0)} \Psi_f([U])(u) = (\mathbb{L}^{\dim V_+} - \mathbb{L}^{\dim V_-}) \cdot \Psi_f([U^0])(u_0). \quad (4.13)$$

Moreover, this holds as an identity of monodromic motives on U^0 .

Proof. Write

$$U^+ = \{u \in U \mid \lim_{t \rightarrow 0} t \cdot u \text{ exists}\}, \quad (4.14)$$

$$U^- = \{u \in U \mid \lim_{t \rightarrow \infty} t \cdot u \text{ exists}\}. \quad (4.15)$$

Shrinking U if necessary, we may assume that $U^\pm = \iota^{-1}(V_\pm \times V_0)$, since we may remove the points in $\iota^{-1}(V_\pm \times V_0)$ such that the limits do not exist. The morphism ι now sends closed $\mathbb{G}_m$ -orbits to closed $\mathbb{G}_m$ -orbits.

By Lemma 4.1.5, we may also assume that ι preserves $\mathbb{G}_m$ -stabilizers, by replacing U with a $\mathbb{G}_m$ -invariant open neighbourhood of U^0 .

Since U is smooth, by Sumihiro [50, Corollary 2], U admits a $\mathbb{G}_m$ -invariant affine open cover. We may thus assume that U is affine and connected, hence integral.

Let $U_\Theta = U \setminus U^-$, and let $U_\Theta^+ = U^+ \setminus U^0 \subset U_\Theta$. Consider the weighted blow-up

$$\pi_\Theta : \tilde{U}_\Theta = {}^w\text{Bl}_{U_\Theta^+}(U_\Theta) \longrightarrow U_\Theta, \quad (4.16)$$

with weight k along the V_{-k} -direction for $k > 0$, and write $\tilde{f}_\Theta = f \circ \pi_\Theta$. Explicitly, as in §4.1.3, we may write

$$W_\Theta = \{(t, v_-, u) \in \mathbb{A}^1 \times (V_- \setminus \{0\}) \times U_\Theta \mid \iota(u)_- = t^{-1} \cdot v_-\}, \quad (4.17)$$

$$\tilde{U}_\Theta = [W_\Theta / \mathbb{G}_m], \quad (4.18)$$

where $\iota(u)_-$ is the projection of $\iota(u)$ to V_- , and $\mathbb{G}_m$ acts on W_Θ by $s \cdot (t, v_-, u) = (s^{-1}t, s^{-1} \cdot v_-, u)$. Note that W_Θ is smooth over $\mathbb{A}^1 \times (V_- \setminus \{0\})$, and hence over $\mathbb{K}$. For

any $u \in U_\Theta^+$, by [Theorem 2.4.4 \(i\)](#), we have

$$\begin{aligned}
& \int_{[v_-] \in {}^w\mathbb{P}(V_-)} \Psi_{\check{f}_\Theta}([\tilde{U}_\Theta])([v_-], u) \\
&= \Psi_f([\tilde{U}_\Theta])(u) \\
&= \Psi_f({}^w\mathbb{P}(V_-) \times U_\Theta^+ + [U_\Theta \setminus U_\Theta^+])(u) \\
&= ([{}^w\mathbb{P}(V_-)] - 1) \cdot \Psi_f([U_\Theta^+])(u) + \Psi_f([U_\Theta])(u) \\
&= \left(\frac{\mathbb{L}^{\dim V_-} - 1}{\mathbb{L} - 1} - 1 \right) \cdot \Psi_f([U_\Theta^+])(u) + \Psi_f([U])(u), \tag{4.19}
\end{aligned}$$

and this holds as an identity of monodromic motives on U_Θ^+ .

Define $p^+ : U_\Theta^+ \rightarrow U^0$ by $p^+(u) = \lim_{t \rightarrow 0} t \cdot u$. Then $f(u) = f(p^+(u))$ for all $u \in U_\Theta^+$, and by [Theorem 2.3.5 \(ii\)](#), we have

$$\Psi_f([U_\Theta^+])(u) = \Psi_f([U^0])(p^+(u)) \tag{4.20}$$

for all $u \in U_\Theta^+$. Again, this holds as an identity of monodromic motives on U_Θ^+ , where the right-hand side means $(p^+)^* \circ \Psi_f([U^0])$.

Now, consider the quotient stack

$$\check{U}_\Theta = [W_\Theta / \mathbb{G}_m^2], \tag{4.21}$$

where $\mathbb{G}_m^2$ acts on W_Θ by $(s_1, s_2) \cdot (t, v_-, u) = (s_1^{-1}t, s_1^{-1}s_2 \cdot v_-, s_2 \cdot u)$. There is, by definition, a principal $\mathbb{G}_m$ -bundle $\pi_\Theta : \check{U}_\Theta \rightarrow \check{U}_\Theta$. There is a morphism $\check{f}_\Theta : \check{U}_\Theta \rightarrow \mathbb{A}^1$ induced by $\check{f}_\Theta$.

Let $U // \mathbb{G}_m$ be the affine GIT quotient, and consider the reduced closed subscheme

$$\tilde{U} \subset {}^w\mathbb{P}(V_+) \times {}^w\mathbb{P}(V_-) \times (U // \mathbb{G}_m) \tag{4.22}$$

consisting of points $([\iota(u)_+], [\iota(u)_-], [u])$ and $([v_+], [v_-], [u_0])$ for $u \in U$, $v_\pm \in V_\pm \setminus \{0\}$, and $u_0 \in U^0$. There is a morphism $f : \tilde{U} \rightarrow \mathbb{A}^1$ induced by f .

Consider the projection $\check{\pi}_\Theta : \check{U}_\Theta \rightarrow \tilde{U}$ given by $(t, v_-, u) \mapsto ([\iota(u)_+], [v_-], [u])$. One can check that fibres of the composition $W_\Theta \rightarrow \tilde{U}$ are single $\mathbb{G}_m^2$ -orbits. We thus have an induced morphism $W_\Theta // \mathbb{G}_m^2 \xrightarrow{\sim} \tilde{U}$, which is an isomorphism by [Lemma 4.1.7](#). Here, we used the fact that $\tilde{U}$ is normal by [Lemma 4.1.6](#), and the fact that W_Θ is integral since it is smooth and connected. In other words, the morphism $\check{\pi}_\Theta$ is a coarse space map. In particular, it is proper by [Lemma 4.1.4](#).

Since the projection $\pi_\Theta : \check{U}_\Theta \rightarrow \check{U}_\Theta$ is smooth and $\check{\pi}_\Theta$ is proper, by [Theorem 2.4.4](#) and [Lemma 4.1.4](#), for any $u \in U_\Theta^+$ and $[v_-] \in {}^w\mathbb{P}(V_-)$, we have

$$\begin{aligned}
\Psi_{\check{f}_\Theta}([\tilde{U}_\Theta])([0, v_-, u]) &= \Psi_{\check{f}_\Theta}([\check{U}_\Theta])([0, v_-, u]) \\
&= \Psi_{\check{f}}([\tilde{U}])([\iota(u)_+], [v_-], [p^+(u)]), \tag{4.23}
\end{aligned}$$

where $[u] = [p^+(u)]$ in $U//\mathbb{G}_m$. Moreover, this holds as an identity of monodromic motives on ${}^w\mathbb{P}(V_-) \times U_\Theta^+$.

Combining (4.19), (4.20), and (4.23), we obtain the identity

$$\begin{aligned} \Psi_f([U])(u) = & \int_{[v_-] \in {}^w\mathbb{P}(V_-)} \Psi_{\tilde{f}}([\tilde{U}])([l(u)_+], [v_-], [p^+(u)]) \\ & + \left(1 - \frac{\mathbb{L}^{\dim V_-} - 1}{\mathbb{L} - 1}\right) \cdot \Psi_f([U^0])(p^+(u)), \end{aligned} \quad (4.24)$$

where $u \in U_\Theta^+$ and $[v_-] \in {}^w\mathbb{P}(V_-)$. Integrating over $u \in U^+(u_0) \setminus \{u_0\}$, we obtain

$$\begin{aligned} \int_{u \in U^+(u_0) \setminus \{u_0\}} \Psi_f([U])(u) = & (\mathbb{L} - 1) \cdot \int_{([v_+], [v_-]) \in {}^w\mathbb{P}(V_+) \times {}^w\mathbb{P}(V_-)} \Psi_{\tilde{f}}([\tilde{U}])([v_+], [v_-], [u_0]) \\ & + (\mathbb{L}^{\dim V_+} - 1) \cdot \left(1 - \frac{\mathbb{L}^{\dim V_-} - 1}{\mathbb{L} - 1}\right) \cdot \Psi_f([U^0])(u_0). \end{aligned} \quad (4.25)$$

Subtracting the analogous identity for integrating over $U_-(u_0) \setminus \{u_0\}$, we arrive at the desired identity (4.13). $\square$

4.1.9. Proof of Theorem 4.1.1. Consider the $\mathbb{G}_m$ -representation $V' = V \times \mathbb{A}^1$, with the $\mathbb{G}_m$ -action on V as given, and on $\mathbb{A}^1$ by scaling. Let $U' = U \times \mathbb{A}^1$, with the $\mathbb{G}_m$ -action on U as given, and on $\mathbb{A}^1$ by scaling, and let $f' = f \circ \text{pr}_1 : U' \rightarrow \mathbb{A}^1$, where $\text{pr}_1 : U' \rightarrow U$ is the projection. Let $u'_0 = (u_0, 0) \in U'^0 = U^0 \times \{0\}$. By Theorem 2.3.5 (ii), we have $\Psi_{f'}([U']) = \text{pr}_1^* \circ \Psi_f([U])$, and similarly, $\Psi_{f'}([U'^0]) = \text{pr}_1^* \circ \Psi_f([U^0])$.

Applying Lemma 4.1.8 to this new set of data, and simplifying the expression by the observations above, we obtain

$$\mathbb{L} \cdot \int_{u \in U^+(u_0)} \Psi_f([U])(u) - \int_{u \in U^-(u_0)} \Psi_f([U])(u) = (\mathbb{L}^{\dim V_+ + 1} - \mathbb{L}^{\dim V_-}) \cdot \Psi_f([U^0])(u_0). \quad (4.26)$$

Subtracting the original identity (4.13) from this, and dividing by $\mathbb{L} - 1$, we obtain the desired identity (4.3).

Finally, (4.4) follows from (4.3) by the definition of Φ_f . $\square$

4.2 The global version

4.2.1. Assumptions on the stack. In the following, we assume that $\mathfrak{X}$ is an oriented (-1) -shifted symplectic stack over $\mathbb{K}$, with classical truncation $\mathfrak{X} = \mathfrak{t}_0(\mathfrak{X})$.

We assume that $\mathfrak{X}$ is an algebraic stack that is Nisnevich locally fundamental in the sense of §2.2.4. For example, as in §2.2.5, this is satisfied if $\mathfrak{X}$ admits a good moduli

space, or can be covered by open substacks with good moduli spaces.

4.2.2. Theorem. *Let $\mathfrak{X}, \mathcal{X}$ be as in §4.2.1. Consider the (-1) -shifted Lagrangian correspondence*

$$\mathrm{Grad}(\mathfrak{X}) \xleftarrow{\mathrm{gr}} \mathrm{Filt}(\mathfrak{X}) \xrightarrow{\mathrm{ev}_1} \mathfrak{X} \quad (4.27)$$

given by Theorem 3.1.5. Then we have the identity

$$\mathrm{gr}_! \circ \mathrm{ev}_1^*(v_{\mathfrak{X}}^{\mathrm{mot}}) = \mathbb{L}^{\mathrm{vdim} \mathrm{Filt}(\mathfrak{X})/2} \cdot v_{\mathrm{Grad}(\mathfrak{X})}^{\mathrm{mot}} \quad (4.28)$$

in $\hat{M}^{\hat{\mu}}(\mathrm{Grad}(\mathcal{X}))$, where $\mathrm{vdim} \mathrm{Filt}(\mathfrak{X})$ is the virtual dimension of $\mathrm{Filt}(\mathfrak{X})$, seen as a function $\pi_0(\mathrm{Grad}(\mathfrak{X})) \simeq \pi_0(\mathrm{Filt}(\mathfrak{X})) \rightarrow \mathbb{Z}$.

We will prove the theorem in two steps. First, in Lemma 4.2.3, we show that the theorem holds for a stack if it holds for a Nisnevich cover of the stack, reducing it to the case of fundamental stacks. Then, we deduce the case of fundamental stacks from the local version, Theorem 4.1.1.

4.2.3. Lemma. *Let $\mathfrak{X}, \mathcal{X}$ be as in §4.2.1. Let $(\mathcal{X}_i \rightarrow \mathcal{X})_{i \in I}$ be a Nisnevich cover, and write $\mathfrak{X}_i = \mathcal{X}_i \times_{\mathcal{X}} \mathfrak{X}$, with the induced (-1) -shifted symplectic structure and orientation. Then, if Theorem 4.2.2 holds for each $\mathfrak{X}_i$, then it holds for $\mathfrak{X}$.*

Proof. For each i , consider the diagram

$$\begin{array}{ccccc} \mathrm{Grad}(\mathfrak{X}_i) & \xleftarrow{\mathrm{gr}} & \mathrm{Filt}(\mathfrak{X}_i) & \xrightarrow{\mathrm{ev}_1} & \mathfrak{X}_i \\ \downarrow & & \downarrow & & \downarrow \\ \mathrm{Grad}(\mathcal{X}) & \xleftarrow{\mathrm{gr}} & \mathrm{Filt}(\mathcal{X}) & \xrightarrow{\mathrm{ev}_1} & \mathcal{X} \end{array} \quad (4.29)$$

where the left-hand square is a pullback square by Lemma 3.2.4. Therefore, there is a commutative diagram

$$\begin{array}{ccccc} \hat{M}^{\hat{\mu}}(\mathrm{Grad}(\mathfrak{X}_i)) & \xleftarrow{\mathrm{gr}_!} & \hat{M}^{\hat{\mu}}(\mathrm{Filt}(\mathfrak{X}_i)) & \xleftarrow{\mathrm{ev}_1^*} & \hat{M}^{\hat{\mu}}(\mathfrak{X}_i) \\ \uparrow & & \uparrow & & \uparrow \\ \hat{M}^{\hat{\mu}}(\mathrm{Grad}(\mathcal{X})) & \xleftarrow{\mathrm{gr}_!} & \hat{M}^{\hat{\mu}}(\mathrm{Filt}(\mathcal{X})) & \xleftarrow{\mathrm{ev}_1^*} & \hat{M}^{\hat{\mu}}(\mathcal{X}) \end{array} \quad (4.30)$$

where the vertical maps are the pullback maps.

By Halpern-Leistner [26, Corollary 1.1.7], one has $\mathrm{Grad}(\mathcal{X}_i) \simeq \mathrm{Grad}(\mathcal{X}) \times_{\mathcal{X}} \mathcal{X}_i$ for all i . Therefore, the family $(\mathrm{Grad}(\mathcal{X}_i) \rightarrow \mathrm{Grad}(\mathcal{X}))_{i \in I}$ is a Nisnevich cover. By Theorem 2.2.3, it is enough to check the identity (4.28) after pulling back to each $\mathrm{Grad}(\mathcal{X}_i)$. But this follows from the identity (4.28) for each $\mathfrak{X}_i$, the commutativity of (4.30), the relation (2.56) establishing the compatibility of the motivic Behrend function with smooth pullbacks, and the fact that the rank of the tangent complex of $\mathrm{Filt}(\mathfrak{X}_i)$ agrees with that of $\mathrm{Filt}(\mathfrak{X})$ on the corresponding components, which follows from (3.12). $\square$

4.2.4. Lemma. Suppose we have a pullback diagram of d -critical stacks

$$\begin{array}{ccc} \mathcal{Y}' & \xrightarrow{g'} & \mathcal{Y} \\ f' \downarrow & \ulcorner & \downarrow f \\ \mathcal{X}' & \xrightarrow{g} & \mathcal{X}, \end{array} \quad (4.31)$$

where all morphisms are smooth and compatible with the d -critical structures.

Let $K_{\mathcal{X}}^{1/2} \rightarrow \mathcal{X}$ and $K_{\mathcal{Y}}^{1/2} \rightarrow \mathcal{Y}$ be orientations, not necessarily compatible with f . Let $K_{\mathcal{X}'}^{1/2} \rightarrow \mathcal{X}'$ and $K_{\mathcal{Y}'}^{1/2} \rightarrow \mathcal{Y}'$ be the orientations induced by $K_{\mathcal{X}}^{1/2}$ and $K_{\mathcal{Y}}^{1/2}$, respectively, as in §2.5.2. Then we have

$$g'^* \circ \Upsilon(K_{\mathcal{Y}}^{1/2} \otimes f^*(K_{\mathcal{X}}^{-1/2}) \otimes \det(\mathbb{L}_{\mathcal{Y}/\mathcal{X}})^{-1}) = \Upsilon(K_{\mathcal{Y}'}^{1/2} \otimes f'^*(K_{\mathcal{X}'}^{-1/2}) \otimes \det(\mathbb{L}_{\mathcal{Y}'/\mathcal{X}'}))^{-1} \quad (4.32)$$

in $\hat{M}^{\text{fl}}(\mathcal{Y}')$, where Υ is the map from §2.1.11, and the parts in $\Upsilon(\cdots)$ are line bundles with trivial square, and can be seen as μ_2 -bundles.

Proof. These line bundles have trivial square by Joyce [31, Lemma 2.58]. We have

$$\begin{aligned} & g'^*(K_{\mathcal{Y}}^{1/2} \otimes f^*(K_{\mathcal{X}}^{-1/2}) \otimes \det(\mathbb{L}_{\mathcal{Y}/\mathcal{X}})^{-1}) \\ & \simeq g'^*(K_{\mathcal{Y}}^{1/2}) \otimes f'^* \circ g^*(K_{\mathcal{X}}^{-1/2}) \otimes \det(g^*(\mathbb{L}_{\mathcal{Y}/\mathcal{X}}))^{-1} \\ & \simeq K_{\mathcal{Y}'}^{1/2} \otimes \det(\mathbb{L}_{\mathcal{Y}'/\mathcal{Y}})^{-1} \otimes f'^*(K_{\mathcal{X}'}^{-1/2}) \otimes f'^* \circ \det(\mathbb{L}_{\mathcal{X}'/\mathcal{X}}) \otimes \det(\mathbb{L}_{\mathcal{Y}'/\mathcal{X}'})^{-1} \\ & \simeq K_{\mathcal{Y}'}^{1/2} \otimes f'^*(K_{\mathcal{X}'}^{-1/2}) \otimes \det(\mathbb{L}_{\mathcal{Y}'/\mathcal{X}'})^{-1}, \end{aligned}$$

and applying Υ gives the desired identity. $\square$

4.2.5. Proof of Theorem 4.2.2. By Lemma 4.2.3, we may assume that $\mathcal{X}$ is fundamental. Let $\mathcal{X} \simeq [S/G]$, where S is an affine $\mathbb{K}$ -variety, and $G = \text{GL}(n)$ for some n . The classical truncation of the correspondence (4.27) can be written as

$$\coprod_{\lambda: \mathbb{G}_m \rightarrow G} [S^{\lambda,0}/L_{\lambda}] \xleftarrow{\text{gr}} \coprod_{\lambda: \mathbb{G}_m \rightarrow G} [S^{\lambda,+}/P_{\lambda}] \xrightarrow{\text{ev}_1} [S/G], \quad (4.33)$$

with notations as in §3.1.2. The assumption on G implies that all the groups L_{λ} and P_{λ} are special groups.

We fix a cocharacter $\lambda: \mathbb{G}_m \rightarrow G$, and prove the identity on the component $[S^{\lambda,+}/P_{\lambda}]$. We may assume that $S^{\lambda,+} \neq \emptyset$.

By Joyce [31, Remark 2.47], shrinking S if necessary, we may assume that there exists a smooth affine $\mathbb{K}$ -scheme U acted on by G , and a G -invariant function $f: U \rightarrow \mathbb{A}^1$, such that $\mathcal{X}$ is isomorphic as a d -critical stack to the critical locus $[\text{Crit}(f)/G]$,

and $S \simeq \text{Crit}(f)$. We now have a commutative diagram

$$\begin{array}{ccccc}
 U^{\lambda,0} & \xleftarrow{p} & U^{\lambda,+} & \xrightarrow{i} & U \\
 \pi^0 \downarrow & & \downarrow \pi^+ & & \downarrow \pi \\
 [U^{\lambda,0}/L_\lambda] & \xleftarrow{\text{gr}} & [U^{\lambda,+}/P_\lambda] & \xrightarrow{\text{ev}_1} & [U/G] .
 \end{array} \tag{4.34}$$

Let $0 \in S^{\lambda,0}$ be a $\mathbb{K}$ -point, and let $V = \mathbb{T}_U|_0$ be the tangent space. Consider the $\mathbb{G}_m$ -actions on U and V via the cocharacter λ . By Luna [46, Lemma in §III.1], shrinking U if necessary, we may choose a $\mathbb{G}_m$ -equivariant étale morphism $\iota: U \rightarrow V$ such that $\iota(0) = 0$. Applying Theorem 4.1.1 gives the identity

$$p_! \circ i^* \circ \Phi_f([U]) = \mathbb{L}^{\dim V_+^\lambda} \cdot \Phi_f([U^{\lambda,0}]) , \tag{4.35}$$

where $V_+^\lambda \subset V$ is the subspace where $\mathbb{G}_m$ acts with positive weights. Note that $\Phi_f(U)$ is supported on S by its definition. Let $K_S^{1/2}$ be the orientation of the d-critical scheme S induced from that of $\mathfrak{X}$. One computes that

$$\begin{aligned}
 & \text{gr}_! \circ \text{ev}_1^*(\nu_{\mathfrak{X}}^{\text{mot}}) \\
 &= [P_\lambda]^{-1} \cdot \text{gr}_! \circ \pi_!^+ \circ (\pi^+)^* \circ \text{ev}_1^*(\nu_{\mathfrak{X}}^{\text{mot}}) \\
 &= [P_\lambda]^{-1} \cdot \pi_!^0 \circ p_! \circ i^* \circ \pi^* (\nu_{\mathfrak{X}}^{\text{mot}}) \\
 &= \mathbb{L}^{\dim G/2} \cdot [P_\lambda]^{-1} \cdot \pi_!^0 \circ p_! \circ i^* (\nu_S^{\text{mot}}) \\
 &= \mathbb{L}^{\dim G/2 - \dim V/2} \cdot [P_\lambda]^{-1} \cdot \pi_!^0 \circ p_! \circ i^* (\Phi_f([U]) \cdot \Upsilon(K_S^{1/2} \otimes K_U^{-1}|_S)) \\
 &= \mathbb{L}^{\dim G/2 - \dim V/2} \cdot [P_\lambda]^{-1} \cdot \\
 & \quad \pi_!^0 \circ p_! \left(i^* \circ \Phi_f([U]) \cdot i^* \circ \pi^* \circ \Upsilon(K_{\mathfrak{X}}^{1/2} \otimes K_{[U/G]}^{-1}|_{\mathfrak{X}}) \right) \\
 &= \mathbb{L}^{\dim G/2 - \dim V/2} \cdot [P_\lambda]^{-1} \cdot \\
 & \quad \pi_!^0 \circ p_! \left(i^* \circ \Phi_f([U]) \cdot (\pi^+)^* \circ \text{ev}_1^* \circ \Upsilon(K_{\mathfrak{X}}^{1/2} \otimes K_{[U/G]}^{-1}|_{\mathfrak{X}}) \right) \\
 &= \mathbb{L}^{\dim G/2 - \dim V/2} \cdot [P_\lambda]^{-1} \cdot \\
 & \quad \pi_!^0 \circ p_! \left(i^* \circ \Phi_f([U]) \cdot (\pi^+)^* \circ \text{gr}^* \circ \Upsilon(K_{\text{Grad}(\mathfrak{X})}^{1/2} \otimes K_{[U^{\lambda,0}/L_\lambda]}^{-1}|_{[S^{\lambda,0}/L_\lambda]}) \right) \\
 &= \mathbb{L}^{\dim G/2 - \dim V/2} \cdot [P_\lambda]^{-1} \cdot \\
 & \quad \pi_!^0 \circ p_! \left(i^* \circ \Phi_f([U]) \cdot p^* \circ (\pi^0)^* \circ \Upsilon(K_{\text{Grad}(\mathfrak{X})}^{1/2} \otimes K_{[U^{\lambda,0}/L_\lambda]}^{-1}|_{[S^{\lambda,0}/L_\lambda]}) \right) \\
 &= \mathbb{L}^{\dim G/2 - \dim V/2} \cdot [P_\lambda]^{-1} \cdot \\
 & \quad \pi_!^0 \left(p_! \circ i^* \circ \Phi_f([U]) \cdot (\pi^0)^* \circ \Upsilon(K_{\text{Grad}(\mathfrak{X})}^{1/2} \otimes K_{[U^{\lambda,0}/L_\lambda]}^{-1}|_{[S^{\lambda,0}/L_\lambda]}) \right) \\
 &= \mathbb{L}^{\dim G/2 - \dim V/2 + \dim V_+^\lambda} \cdot [P_\lambda]^{-1} \cdot \\
 & \quad \pi_!^0 \left(\Phi_f([U^{\lambda,0}]) \cdot (\pi^0)^* \circ \Upsilon(K_{\text{Grad}(\mathfrak{X})}^{1/2} \otimes K_{[U^{\lambda,0}/L_\lambda]}^{-1}|_{[S^{\lambda,0}/L_\lambda]}) \right)
 \end{aligned}$$

$$\begin{aligned}
&= \mathbb{L}^{\dim G/2 - \dim V/2 + \dim V_+^\lambda} \cdot [P_\lambda]^{-1} \cdot \\
&\quad \pi_!^0 \left(\Phi_f([U^{\lambda,0}]) \cdot \Upsilon(K_{S^{\lambda,0}}^{1/2} \otimes K_{U^{\lambda,0}}^{-1}|_{S^{\lambda,0}}) \right) \\
&= \mathbb{L}^{\dim G/2 - \dim V/2 + \dim V_+^\lambda - \dim V_0^\lambda/2} \cdot [P_\lambda]^{-1} \cdot \pi_!^0(v_{S^{\lambda,0}}^{\text{mot}}) \\
&= \mathbb{L}^{(\dim G - \dim L_\lambda)/2 + (\dim V_+^\lambda - \dim V_-^\lambda)/2} \cdot [P_\lambda]^{-1} \cdot \pi_!^0 \circ (\pi^0)^*(v_{\text{Grad}(\mathfrak{X})}^{\text{mot}}) \\
&= \mathbb{L}^{(\dim G - \dim L_\lambda)/2 + (\dim V_+^\lambda - \dim V_-^\lambda)/2} \cdot [P_\lambda]^{-1} \cdot [L_\lambda] \cdot v_{\text{Grad}(\mathfrak{X})}^{\text{mot}} \\
&= \mathbb{L}^{(\dim V_+^\lambda - \dim V_-^\lambda)/2} \cdot v_{\text{Grad}(\mathfrak{X})}^{\text{mot}}. \tag{4.36}
\end{aligned}$$

Here, the first step uses (2.14); the third uses (2.55); the fourth uses (2.52); the fifth uses Lemma 4.2.4, where the morphism f there is taken to be an isomorphism; the seventh uses the fact that the shifted Lagrangian correspondence (4.27) is oriented; the ninth uses (2.13); the tenth is the key step, and uses (4.35); the eleventh is analogous to the fifth; the twelfth uses (2.52) again; the thirteenth uses (2.55) again; the fourteenth uses (2.14) again; and the final step uses the relation $[P_\lambda] = [L_\lambda] \cdot \mathbb{L}^{(\dim G - \dim L_\lambda)/2}$.

Finally, we verify that $\text{vdim Filt}^\lambda(\mathfrak{X}) = \dim V_+^\lambda - \dim V_-^\lambda$, where $\text{Filt}^\lambda(\mathfrak{X}) \subset \text{Filt}(\mathfrak{X})$ is the closed and open substack corresponding to the cocharacter λ . Indeed, let $\mathfrak{X}' = [\text{Crit}(f)/G]$ as a derived critical locus, with the natural (-1) -shifted symplectic structure, so $\mathfrak{t}_0(\mathfrak{X}') \simeq \mathfrak{X}$. For $x \in S^{\lambda,0}(\mathbb{K})$, by Lemma 3.1.6, one has

$$\begin{aligned}
\text{rank}(\mathbb{L}_{\text{Filt}^\lambda(\mathfrak{X})|_x}) &= \text{rank}^{[0,1]}(\mathbb{L}_{\text{Filt}^\lambda(\mathfrak{X})|_x}) - \text{rank}^{[0,1]}(\mathbb{L}_{\text{Filt}^{-\lambda}(\mathfrak{X})|_x}) \\
&= \text{rank}^{[0,1]}(\mathbb{L}_{\text{Filt}^\lambda(\mathfrak{X})|_x}) - \text{rank}^{[0,1]}(\mathbb{L}_{\text{Filt}^{-\lambda}(\mathfrak{X})|_x}) \\
&= \text{rank}(\mathbb{L}_{\text{Filt}^\lambda(\mathfrak{X}')|_x}), \tag{4.37}
\end{aligned}$$

where $\text{rank}^{[0,1]} = \dim H^0 - \dim H^1$. We have a presentation

$$\mathbb{L}_{\mathfrak{X}'|_x} \simeq \left(\mathfrak{g} \longrightarrow \mathbb{T}_U|_x \longrightarrow \mathbb{L}_U|_x \longrightarrow \mathfrak{g}^\vee \right) \tag{4.38}$$

with degrees in $[-1, 2]$, where $\mathfrak{g}$ is the Lie algebra of G . By Halpern-Leistner [26, Lemma 1.2.3], we have $\text{fi}^*(\mathbb{L}_{\text{Filt}^\lambda(\mathfrak{X})}) \simeq \text{tot}^*(\mathbb{L}_{\mathfrak{X}})_{\leq 0}$, where $(-)_{\leq 0}$ denotes the part of non-positive weights with respect to the natural $\mathbb{G}_m$ -action. This now gives

$$\mathbb{L}_{\text{Filt}^\lambda(\mathfrak{X}')|_x} \simeq \left(\mathfrak{p}_\lambda \longrightarrow \mathbb{T}_{U^{\lambda,-}}|_x \longrightarrow \mathbb{L}_{U^{\lambda,+}}|_x \longrightarrow \mathfrak{p}_{-\lambda}^\vee \right), \tag{4.39}$$

where $\mathfrak{p}_\lambda$ is the Lie algebra of P_λ , and $-\lambda$ is the opposite cocharacter of λ . Note that $\dim P_\lambda = \dim P_{-\lambda}$ and that $\dim U^{\lambda,\pm} = \dim V_\pm^\lambda + \dim V_0^\lambda$. It follows that $\text{vdim Filt}(\mathfrak{X})$, which is equal to the rank of (4.39) by (4.37), is $\dim V_+^\lambda - \dim V_-^\lambda$. $\square$

4.3 The numerical version

4.3.1. Assumptions on the stack. In the following, we assume that $\mathfrak{X}$ is a (-1) -shifted symplectic stack over $\mathbb{K}$, with classical truncation $\mathfrak{X} = \mathfrak{t}_0(\mathfrak{X})$. Note that we no longer assume that $\mathfrak{X}$ is oriented.

We assume that $\mathcal{X}$ is an algebraic stack that is étale locally fundamental in the sense of §2.2.4. For example, as mentioned in §2.2.5, this is satisfied if $\mathcal{X}$ has affine stabilizers and has reductive stabilizers at closed points.

4.3.2. For a graded point $\gamma \in \text{Grad}(\mathcal{X})(\mathbb{K})$, write

$$\mathbb{P}(\text{gr}^{-1}(\gamma)) = \left([* / \mathbb{G}_m] \times_{\text{Grad}(\mathcal{X})} \text{Filt}(\mathcal{X}) \right) \setminus \{ \text{fi}(\gamma) \}, \quad (4.40)$$

where the map $[* / \mathbb{G}_m] \rightarrow \text{Grad}(\mathcal{X})$ is given by the tautological $\mathbb{G}_m$ -action on γ . The $\mathbb{K}$ -point $\text{fi}(\gamma)$ is closed in the fibre product, which can be seen from the étale local description in Theorem 3.2.2. The space $\mathbb{P}(\text{gr}^{-1}(\gamma))$ can be seen as the projectivized space of filtrations of a given associated graded point.

4.3.3. **Theorem.** *Let $\mathfrak{X}, \mathcal{X}$ be as in §4.3.1. Let $\gamma \in \text{Grad}(\mathcal{X})(\mathbb{K})$ be a graded point, and let $\bar{\gamma} = \text{op}(\gamma)$ be its opposite graded point.*

Then we have the numerical identities

$$v_{\mathcal{X}}(\text{tot}(\gamma)) = (-1)^{\text{rank}^{[0,1]}(\mathbb{L}_{\text{Filt}(\mathcal{X})} |_{\text{fi}(\gamma)}) - \text{rank}^{[0,1]}(\mathbb{L}_{\text{Filt}(\mathcal{X})} |_{\text{fi}(\bar{\gamma})})} \cdot v_{\text{Grad}(\mathcal{X})}(\gamma), \quad (4.41)$$

$$\begin{aligned} & \int_{\varphi \in \mathbb{P}(\text{gr}^{-1}(\gamma))} v_{\mathcal{X}}(\text{ev}_1(\varphi)) d\chi - \int_{\varphi \in \mathbb{P}(\text{gr}^{-1}(\bar{\gamma}))} v_{\mathcal{X}}(\text{ev}_1(\varphi)) d\chi \\ &= (\dim H^0(\mathbb{L}_{\text{Filt}(\mathcal{X})} |_{\text{fi}(\gamma)}) - \dim H^0(\mathbb{L}_{\text{Filt}(\mathcal{X})} |_{\text{fi}(\bar{\gamma})})) \cdot v_{\mathcal{X}}(\text{tot}(\gamma)), \end{aligned} \quad (4.42)$$

where $\text{rank}^{[0,1]} = \dim H^0 - \dim H^1$.

This theorem is a generalization of Joyce–Song [35, Theorem 5.11], who considered the case when $\mathcal{X}$ is the moduli stack of objects in a 3-Calabi–Yau category.

4.3.4. **Proof of Theorem 4.3.3.** By a similar argument as in the proof of Lemma 4.2.3, passing to a representable étale cover of $\mathcal{X}$ by fundamental stacks, which induces representable étale covers of $\text{Grad}(\mathcal{X})$ and $\text{Filt}(\mathcal{X})$ by Theorem 3.2.2, it is enough to prove the theorem when $\mathcal{X} \simeq [S/G]$ is fundamental, where S is an affine $\mathbb{K}$ -scheme acted on by a reductive group G . Here, we are using étale descent for constructible functions, instead of Nisnevich descent for motives.

As in §4.2.5, shrinking S if necessary, we may assume that there exists a smooth affine $\mathbb{K}$ -scheme U acted on by G , and a G -invariant function $f: U \rightarrow \mathbb{A}^1$, such that $\mathcal{X}$ is isomorphic as a d-critical stack to the critical locus $[\text{Crit}(f)/G]$. Now, $\mathcal{X}$ comes with a natural orientation, and the motivic Behrend function $v_{\mathcal{X}}^{\text{mot}}$ is defined.

Applying Theorem 4.2.2, then evaluating the Euler characteristics at γ , we obtain the identity

$$\int_{\varphi \in \text{gr}^{-1}(\gamma)} v_{\mathcal{X}}(\text{ev}_1(\varphi)) d\chi = (-1)^{\text{vdim}_{\gamma} \text{Filt}(\mathfrak{X})} \cdot v_{\text{Grad}(\mathcal{X})}(\gamma). \quad (4.43)$$

Let $\varphi_0 = \text{fi}(\gamma)$. Then the left-hand side of (4.43) is equal to $v_{\mathcal{X}}(\text{ev}_1(\varphi_0)) = v_{\mathcal{X}}(\text{tot}(\gamma))$, since the integrand is $\mathbb{G}_m$ -invariant and φ_0 is in the closure of all $\mathbb{G}_m$ -orbits. Also, by Lemma 3.1.6, we have

$$\text{vdim}_{\gamma} \text{Filt}(\mathfrak{X}) = \text{rank}^{[0,1]} \mathbb{L}_{\text{Filt}(\mathcal{X})|_{\text{fi}(\gamma)}} - \text{rank}^{[0,1]} \mathbb{L}_{\text{Filt}(\mathcal{X})|_{\text{fi}(\tilde{\gamma})}}. \quad (4.44)$$

This verifies (4.41).

For (4.42), apply Theorem 4.2.2 again, then take the difference of the evaluations at γ and $\tilde{\gamma}$. This gives the identity

$$\begin{aligned} & (\mathbb{L} - 1) \cdot \left[\int_{\varphi \in \mathbb{P}(\text{gr}^{-1}(\gamma))} v_{\mathfrak{X}}^{\text{mot}}(\text{ev}_1(\varphi)) - \int_{\varphi \in \mathbb{P}(\text{gr}^{-1}(\tilde{\gamma}))} v_{\mathfrak{X}}^{\text{mot}}(\text{ev}_1(\varphi)) \right] \\ & + \mathbb{L}^{\dim H^1(\mathbb{L}_{\text{Grad}(\mathcal{X})}|_{\gamma})} \cdot \left(\mathbb{L}^{-\dim H^1(\mathbb{L}_{\text{Filt}(\mathcal{X})}|_{\text{fi}(\gamma)})} - \mathbb{L}^{-\dim H^1(\mathbb{L}_{\text{Filt}(\mathcal{X})}|_{\text{fi}(\tilde{\gamma})})} \right) \cdot v_{\mathfrak{X}}^{\text{mot}}(\text{tot}(\gamma)) \\ & = \left(\mathbb{L}^{\text{rank}(\mathbb{L}_{\text{Filt}(\mathcal{X})}|_{\text{fi}(\gamma)})/2} - \mathbb{L}^{-\text{rank}(\mathbb{L}_{\text{Filt}(\mathcal{X})}|_{\text{fi}(\gamma)})/2} \right) \cdot v_{\text{Grad}(\mathcal{X})}^{\text{mot}}(\gamma) \end{aligned} \quad (4.45)$$

of monodromic motives over $\mathbb{K}$. Here, we used the fact that the stabilizer group G_{γ} of γ in $\text{gr}^{-1}(\gamma)$ is special and has motive $\mathbb{L}^{\dim G_{\gamma}}$, since G_{γ} is a subgroup of the fibre of the projection $P_{\lambda} \rightarrow L_{\lambda}$, and can be obtained by repeated extensions of $\mathbb{G}_a$. All of this can be seen by, for example, equivariantly embedding S into an affine space with a linear G -action.

Starting from (4.45), we divide both sides by $\mathbb{L} - 1$, and then take the Euler characteristic, which sets $\mathbb{L}^{1/2}$ to -1 . We then apply the identity (4.41) to convert $v_{\text{Grad}(\mathcal{X})}(\gamma)$ to $v_{\mathcal{X}}(\text{tot}(\gamma))$. This gives the desired identity (4.42). $\square$

References

- [1] ALPER, J., ‘Good moduli spaces for Artin stacks’, *Ann. Inst. Fourier*, 63/6 (2013), 2349–402. doi: [10.5802/aif.2833](#), arXiv: [0804.2242](#).
- [2] ALPER, J., HALL, J., and RYDH, D., ‘A Luna étale slice theorem for algebraic stacks’, *Ann. Math.* 191/3 (2020), 675–738. doi: [10.4007/annals.2020.191.3.1](#), arXiv: [1504.06467](#).
- [3] ALPER, J., HALL, J., and RYDH, D., ‘The étale local structure of algebraic stacks’, version 3 (13 Apr. 2023), arXiv: [1912.06162v3](#).
- [4] ALPER, J., HALPERN-LEISTNER, D., and HEINLOTH, J., ‘Existence of moduli spaces for algebraic stacks’, *Invent. Math.* 234/3 (2023), 949–1038. doi: [10.1007/s00222-023-01214-4](#), arXiv: [1812.01128](#).
- [5] AMORIM, L., and BEN-BASSAT, O., ‘Perversely categorified Lagrangian correspondences’, *Adv. Theor. Math. Phys.* 21/2 (2017), 289–381. doi: [10.4310/atmp.2017.v21.n2.a1](#), arXiv: [1601.01536](#).
- [6] BEHREND, K., ‘Donaldson–Thomas type invariants via microlocal geometry’, *Ann. Math.* 170/3 (2009), 1307–38. doi: [10.4007/annals.2009.170.1307](#), arXiv: [math/0507523](#).
- [7] BEN-BASSAT, O., BRAV, C., BUSSI, V., and JOYCE, D., ‘A “Darboux theorem” for shifted symplectic structures on derived Artin stacks, with applications’, *Geom. Topol.* 19/3 (2015), 1287–359. doi: [10.2140/gt.2015.19.1287](#), arXiv: [1312.0090](#).
- [8] BERGH, D., and RYDH, D., ‘Functorial destackification and weak factorization of orbifolds’, version 1 (2 May 2019), arXiv: [1905.00872v1](#).
- [9] BITTNER, F., ‘On motivic zeta functions and the motivic nearby fiber’, *Math. Z.* 249/1 (2005), 63–83. doi: [10.1007/s00209-004-0689-1](#), arXiv: [math/0307033](#).
- [10] BRAV, C., BUSSI, V., DUPONT, D., JOYCE, D., and SZENDRŐI, B., ‘Symmetries and stabilization for sheaves of vanishing cycles’, *J. Singul.* 11 (2015), 85–151. doi: [10.5427/jsing.2015.11e](#), arXiv: [1211.3259](#).
- [11] BRAV, C., BUSSI, V., and JOYCE, D., ‘A Darboux theorem for derived schemes with shifted symplectic structure’, *J. Am. Math. Soc.* 32/2 (2019), 399–443. doi: [10.1090/jams/910](#), arXiv: [1305.6302](#).
- [12] BRAV, C., and DYCKERHOFF, T., ‘Relative Calabi–Yau structures. II: Shifted Lagrangians in the moduli of objects’, *Sel. Math., New Ser.* 27/4 (2021), 45. doi: [10.1007/s00029-021-00642-5](#), arXiv: [1812.11913](#).
- [13] BU, C., ‘Enumerative invariants in self-dual categories. I. Motivic invariants’, version 3 (22 Jan. 2024), arXiv: [2302.00038v3](#).
- [14] BUSSI, V., JOYCE, D., and MEINHARDT, S., ‘On motivic vanishing cycles of critical loci’, *J. Algebr. Geom.* 28/3 (2019), 405–38. doi: [10.1090/jag/737](#), arXiv: [1305.6428](#).
- [15] CALAQUE, D., ‘Lagrangian structures on mapping stacks and semi-classical TFTs’, in *Stacks and categories in geometry, topology, and algebra* (Contemp. Math. 643; American Mathematical Society, 2015), 1–23. doi: [10.1090/conm/643/12894](#), arXiv: [1306.3235](#).
- [16] CALAQUE, D., HAUGSENG, R., and SCHEIMBAUER, C., ‘The AKSZ construction in derived algebraic geometry as an extended topological field theory’, version 2 (16 Nov. 2022), arXiv: [2108.02473v2](#).

- [17] CHOUDHURY, U., DESHMUKH, N., and HOGADI, A., ‘The Nisnevich motive of an algebraic stack’, *Ann. K-Theory*, 8/2 (2023), 245–73. doi: [10.2140/akt.2023.8.245](https://doi.org/10.2140/akt.2023.8.245), arXiv: [2012.13304](https://arxiv.org/abs/2012.13304).
- [18] CONRAD, B., ‘The Keel–Mori theorem via stacks’ (27 Nov. 2005), <https://math.stanford.edu/~conrad/papers/coarsespace.pdf>.
- [19] CONRAD, B., LIEBLICH, M., and OLSSON, M., ‘Nagata compactification for algebraic spaces’, *J. Inst. Math. Jussieu*, 11/4 (2012), 747–814. doi: [10.1017/S1474748011000223](https://doi.org/10.1017/S1474748011000223), arXiv: [0910.5008](https://arxiv.org/abs/0910.5008).
- [20] DENEFF, J., and LOESER, F., ‘Motivic Igusa zeta functions’, *J. Algebr. Geom.* 7/3 (1998), 505–37, arXiv: [math/9803040](https://arxiv.org/abs/math/9803040).
- [21] DENEFF, J., and LOESER, F., ‘Geometry on arc spaces of algebraic varieties’, in C. Casacuberta, R. M. Miró-Roig, J. Verdera, and S. Xambó-Descamps (eds.), *European congress of mathematics 2000*, volume I (Prog. Math. 201; Birkhäuser, 2001), 327–48. doi: [10.1007/978-3-0348-8268-2_19](https://doi.org/10.1007/978-3-0348-8268-2_19), arXiv: [math/0006050](https://arxiv.org/abs/math/0006050).
- [22] DENEFF, J., and LOESER, F., ‘Lefschetz numbers of iterates of the monodromy and truncated arcs’, *Topology*, 41/5 (2002), 1031–40. doi: [10.1016/S0040-9383\(01\)00016-7](https://doi.org/10.1016/S0040-9383(01)00016-7), arXiv: [math/0001105](https://arxiv.org/abs/math/0001105).
- [23] DRINFELD, V., and GAITSGORY, D., ‘On a theorem of Braden’, *Transform. Groups*, 19/2 (2014), 313–58. doi: [10.1007/s00031-014-9267-8](https://doi.org/10.1007/s00031-014-9267-8), arXiv: [1308.3786](https://arxiv.org/abs/1308.3786).
- [24] ENCINAS, S., and VILLAMAYOR, O., ‘Good points and constructive resolution of singularities’, *Acta Math.* 181/1 (1998), 109–58. doi: [10.1007/bf02392749](https://doi.org/10.1007/bf02392749).
- [25] GROTHENDIECK, A., ‘Éléments de géométrie algébrique’, *Publ. Math., Inst. Hautes Étud. Sci.* 4 (1960), 1–228; 8 (1961), 1–222; 11 (1962), 349–511; 17 (1963), 137–223; 20 (1964), 101–355; 24 (1965), 1–231; 28 (1966), 1–255; 32 (1967), 1–361, in collab. with J. Dieudonné.
- [26] HALPERN-LEISTNER, D., ‘On the structure of instability in moduli theory’, version 5 (4 Feb. 2022), arXiv: [1411.0627v5](https://arxiv.org/abs/1411.0627v5).
- [27] HALPERN-LEISTNER, D., ‘Derived Θ -stratifications and the D -equivalence conjecture’, version 2 (18 June 2021), arXiv: [2010.01127v2](https://arxiv.org/abs/2010.01127v2).
- [28] JOYCE, D., ‘Constructible functions on Artin stacks’, *J. Lond. Math. Soc.* 74/3 (2006), 583–606. doi: [10.1112/S0024610706023180](https://doi.org/10.1112/S0024610706023180), arXiv: [math/0403305](https://arxiv.org/abs/math/0403305).
- [29] JOYCE, D., ‘Configurations in abelian categories. II. Ringel–Hall algebras’, *Adv. Math.* 210/2 (2007), 635–706. doi: [10.1016/j.aim.2006.07.006](https://doi.org/10.1016/j.aim.2006.07.006), arXiv: [math/0503029](https://arxiv.org/abs/math/0503029).
- [30] JOYCE, D., ‘Motivic invariants of Artin stacks and “stack functions”’, *Q. J. Math.* 58/3 (2007), 345–92. doi: [10.1093/qmath/ham019](https://doi.org/10.1093/qmath/ham019), arXiv: [math/0509722](https://arxiv.org/abs/math/0509722).
- [31] JOYCE, D., ‘A classical model for derived critical loci’, *J. Differ. Geom.* 101/2 (2015), 289–367. doi: [10.4310/jdg/1442364653](https://doi.org/10.4310/jdg/1442364653), arXiv: [1304.4508](https://arxiv.org/abs/1304.4508).
- [32] JOYCE, D., ‘Ringel–Hall style vertex algebra and Lie algebra structures on the homology of moduli spaces’, preliminary version (Mar. 2018), <https://people.maths.ox.ac.uk/joyce/hall.pdf>.
- [33] JOYCE, D., ‘Enumerative invariants and wall-crossing formulae in abelian categories’, version 1 (8 Nov. 2021), arXiv: [2111.04694v1](https://arxiv.org/abs/2111.04694v1).
- [34] JOYCE, D., and SAFRONOV, P., ‘A Lagrangian neighbourhood theorem for shifted symplectic derived schemes’, *Ann. Fac. Sci. Toulouse, Math. (6)*, 28/5 (2019), 831–908. doi: [10.5802/afst.1616](https://doi.org/10.5802/afst.1616), arXiv: [1506.04024](https://arxiv.org/abs/1506.04024).

- [35] JOYCE, D., and SONG, Y., *A theory of generalized Donaldson–Thomas invariants* (Mem. Am. Math. Soc. 1020; American Mathematical Society, 2012). doi: [10.1090/s0065-9266-2011-00630-1](https://doi.org/10.1090/s0065-9266-2011-00630-1), arXiv: [0810.5645](https://arxiv.org/abs/0810.5645).
- [36] KEEL, S., and MORI, S., ‘Quotients by groupoids’, *Ann. Math.* 145/1 (1997), 193–213. doi: [10.2307/2951828](https://doi.org/10.2307/2951828), arXiv: [alg-geom/9508012](https://arxiv.org/abs/alg-geom/9508012).
- [37] KINJO, T., PARK, H., and SAFRONOV, P., ‘Cohomological Hall algebras for 3-Calabi–Yau categories’, in preparation.
- [38] KNUTSON, D., *Algebraic spaces* (Lect. Notes Math. 203; Springer, 1971). doi: [10.1007/bfb0059750](https://doi.org/10.1007/bfb0059750).
- [39] KOLLÁR, J., *Lectures on resolution of singularities* (Ann. Math. Stud. 166; Princeton University Press, 2007). doi: [10.1515/9781400827800](https://doi.org/10.1515/9781400827800).
- [40] KONTSEVICH, M., and SOIBELMAN, Y., ‘Stability structures, motivic Donaldson–Thomas invariants and cluster transformations’, version 1 (16 Nov. 2008), arXiv: [0811.2435v1](https://arxiv.org/abs/0811.2435v1).
- [41] KONTSEVICH, M., and SOIBELMAN, Y., ‘Cohomological Hall algebra, exponential Hodge structures and motivic Donaldson–Thomas invariants’, *Commun. Number Theory Phys.* 5/2 (2011), 231–352. doi: [10.4310/cntp.2011.v5.n2.a1](https://doi.org/10.4310/cntp.2011.v5.n2.a1), arXiv: [1006.2706](https://arxiv.org/abs/1006.2706).
- [42] KRESCH, A., ‘Cycle groups for Artin stacks’, *Invent. Math.* 138/3 (1999), 495–536. doi: [10.1007/s002220050351](https://doi.org/10.1007/s002220050351).
- [43] LAUMON, G., and MORET-BAILLY, L., *Champs algébriques* (Ergeb. Math. Grenzgeb., 3. Folge, 39; Springer, 2000). doi: [10.1007/978-3-540-24899-6](https://doi.org/10.1007/978-3-540-24899-6).
- [44] LÊ, Q. T., ‘Proofs of the integral identity conjecture over algebraically closed fields’, *Duke Math. J.* 164/1 (2015), 157–94. doi: [10.1215/00127094-2869138](https://doi.org/10.1215/00127094-2869138), arXiv: [1206.5334](https://arxiv.org/abs/1206.5334).
- [45] LOOIJENGA, E., ‘Motivic measures’, in *Séminaire Bourbaki. Volume 1999/2000. Exposés 865–879* (Société Mathématique de France, 2002), 267–97, arXiv: [math/0006220](https://arxiv.org/abs/math/0006220).
- [46] LUNA, D., ‘Slices étales’, *Bull. Soc. Math. Fr., Suppl., Mém.* 33 (1973), 81–105.
- [47] PANTEV, T., TOËN, B., VAQUIÉ, M., and VEZZOSI, G., ‘Shifted symplectic structures’, *Publ. Math., Inst. Hautes Étud. Sci.* 117 (2013), 271–328. doi: [10.1007/s10240-013-0054-1](https://doi.org/10.1007/s10240-013-0054-1), arXiv: [1111.3209](https://arxiv.org/abs/1111.3209).
- [48] RYDH, D., ‘Compactification of tame Deligne–Mumford stacks’ (17 May 2011), <https://people.kth.se/~dary/tamecompactification20110517.pdf>.
- [49] SERRE, J.-P., ‘Espaces fibrés algébriques’, in *Séminaire Claude Chevalley. Tome 3. Anneaux de Chow et applications* (Secrétariat Mathématique, 1958), 1–37.
- [50] SUMIHIRO, H., ‘Equivariant completion’, *J. Math. Kyoto Univ.* 14 (1974), 1–28. doi: [10.1215/kjm/1250523277](https://doi.org/10.1215/kjm/1250523277).
- [51] TOËN, B., and VEZZOSI, G., *Homotopical algebraic geometry II: Geometric stacks and applications* (Mem. Am. Math. Soc. 902; American Mathematical Society, 2008). doi: [10.1090/memo/0902](https://doi.org/10.1090/memo/0902), arXiv: [math/0404373](https://arxiv.org/abs/math/0404373).

Mathematical Institute, University of Oxford, Oxford OX2 6GG, United Kingdom.
 Email: bu@maths.ox.ac.uk